Temporal dynamics of seedling emergence among four fire ephemerals: the
interplay of after-ripening and embryo growth with smoke

Siti N. Hidayati¹, David J. Merritt^{2,3}, Shane R. Turner^{2,3}, Kingsley W. Dixon^{2,3,4} and Jeffrey L.
Walck¹

¹Department of Biology, Middle Tennessee State University, Murfreesboro, TN 37132, USA,
²Kings Park Science, Department of Biodiversity, Conservation and Attractions, Kings Park, WA
6005, Australia, ³School of Biological Sciences, The University of Western Australia, Crawley,
WA 6009, Australia, ⁴Department of Environment and Agriculture, Curtin University, Bentley,
WA 6102, Australia

Author for correspondence: Jeffrey L. Walck, Email: Jeffrey.Walck@mtsu.edu

Running title: Temporal dynamics of emergence among fire ephemerals

Abstract

The flora of Mediterranean ecosystems contains families with species having fully- and under-developed embryos in their seeds. After-ripening for physiological dormancy release and smoke influence germination in many species. We investigated how after-ripening and embryo growth interact with smoke to influence the temporal dynamics of seedling emergence among fire ephemerals. Seeds were placed in the field and under standardised (50% RH, 30°C) laboratory conditions to test the effects of summer conditions on physiological dormancy loss. Germination was tested with water or smoke compounds (smoke water, KAR₁) at a simulated autumn/winter temperature (18/7°C). The timing and amount of seedling emergence with smoke was observed for seeds exposed to near-natural conditions. During summer, physiological dormancy was broken in all species, enabling germination at autumn/winter but not summer temperatures; no embryo growth occurred in seeds with under-developed embryos. At the start of the wet season, seedling emergence from seeds with fully-developed embryos occurred earlier than from seeds with under-developed embryos. In a non-consistent manner among our study species, smoke and smoke compounds influenced the rate of embryo growth and amount of germination. Effects of smoke were noticeable in terms of number of emergents in the first emergence season. Among ecologically similar species, we have shown (1) that both thermal and embryo traits exclude germination in the summer, (2) how embryo size influences the timing of seedling emergence in autumn-winter, and (3) a reduced requirement for smoke in the second emergence season after a fire with a shift to reliance on seasonal cues for emergence.

Key words: embryos, ephemerals, fire, KAR₁, Mediterranean climate, seedling emergence, smoke, smoke water

Introduction

Seed dormancy regulates germination until near optimal conditions for seedling emergence and establishment in species-specific ways. The degree of physiological dormancy is well-known to influence germination timing, thereby setting up variation among species in their emergence patterns (Baskin and Baskin, 2014). However, another component of dormancy that has received less attention with regards to germination timing is initial embryo size and the amount of embryo growth that must occur within seeds before radicle emergence (Vivrette, 1995; Verdú, 2006; Vandeloos *et al.*, 2012). Seed embryos may have distinguishable cotyledons and a radicle, but under-developed embryos must elongate within the seed before radicle emergence, in contrast to fully-developed embryos which simply push through the seed coat once key germination conditions are met (Baskin and Baskin, 2014). The two types of embryos also may have a physiological block that must be overcome to enable embryo growth and/or germination. Altogether, the degree of physiological dormancy and timing of dormancy loss interacts with embryo growth to time germination and emergence in response to environmental cues.

The flora of Mediterranean ecosystems contains families with species having fully-developed embryos in their seeds as well as species with under-developed embryos. Physiological dormancy is prominent in seeds of many species, with germination and seedling recruitment only possible during a relatively narrow window during the winter-wet season that is punctuated by the long summer-dry season (Merritt *et al.*, 2007; Carta *et al.*, 2016; Bart *et al.*, 2017). In these habitats, fire is a major periodic disturbance that provides a primary recruitment opportunity for many plants due to increased resource availability (i.e. moisture, nutrients and light), especially for short-lived fire ephemerals that solely rely on seeds stored in the soil seed bank (Rundel and Parsons, 1984; Brown, 1993; Miller and Dixon, 2014). Heat and smoke are the main components of fire influencing germination, and within smoke, KAR₁ is a key compound

that elicits remarkable germination across many taxonomically diverse species (Chiwocha *et al.*, 2009).

During summer months in Mediterranean ecosystems, we speculated that release from physiological dormancy would occur in seeds of our study species regardless of embryo type, but under-developed embryos would not grow since seeds are in a relatively dry state (Turner *et al.*, 2006). When autumn rains begin in concert with a reduction in temperatures, radicles in seeds with fully-developed embryos would readily emerge. In contrast, under-developed embryos would first need time to grow inside the seed before radicle emergence, and thus, germination would be delayed. Concomitantly, under-developed embryos may grow at different rates, leading to non-uniformity in emergence among seeds, and thus spreading emergence over a wider period of time (Hidayati *et al.*, 2012). Germination may also be spread over more than one season, due to variation in dormancy depth amongst individual seeds within a seed population (Hidayati *et al.*, 2012). In addition to germination stimulation, KAR₁ and smoke water have been demonstrated to enhance seedling growth and vigour (Daws *et al.*, 2007; Ghebrehiwot *et al.*, 2009; Nelson *et al.*, 2012; Zhou *et al.*, 2012). Thus, we reasoned that these smoke products would enhance embryo growth prior to germination and allow earlier emergence for treated seeds as compared to non-treated seeds.

In the present study, we investigated the patterns of embryo growth, germination, and seedling emergence as influenced by climatic factors and fire-related cues. Four common, sympatric annual species were studied: *Podotheca angustifolia* (Labill.) Less. (Asteraceae) and *Austrostipa macalpinei* (Reader) S.W.L. Jacobs and J. Everett (Poaceae) with fully-developed embryos and *Trachymene pilosa* Sm. (Apiaceae) and *Levenhookia pusilla* R. Br. (Stylidiaceae) with under-developed embryos. In the winter rainfall region of southwest Western Australia, these species grow and co-occur in fire-prone habitats, such as *Banksia* woodlands and eucalypt

forests (Lamont, 1985; Grant and Loneragan, 1999; Barrett and Tay, 2005), and in less-fire-prone areas, such as granite outcrops (Hopper *et al.*, 1997). The highest population densities of these species are usually found in woodlands or forests following fire, although they do also occur and persist in habitats that have not been recently burnt.

The first aspect of our study was to examine the loss of physiological dormancy over time in relation to growth of embryos within seeds prior to radicle emergence and germination under a range of field and laboratory conditions among our study species. To achieve these aims, we tested embryo growth and germination on water, KAR₁, and smoke water in light and darkness. Second, we followed the emergence of our study species in a tunnel-house trial, and related the emergence observed to ambient temperatures, soil moisture and the application of smoke. Specifically, we documented the onset, median, and time to maximum emergence and the amount of emergence during two winter seasons.

Materials and methods

Plant material

Freshly matured caryopses (*Austrostipa*), true seeds (*Levenhookia*), achenes (*Podotheca*), and mericarps (*Trachymene*) (hereafter, seeds) were collected during the dispersal season: in November 2007 for *Podotheca* and *Trachymene* (hereafter, 2007 cohort) and for all four species in November 2008 (hereafter, 2008 cohort). Due to the ephemeral nature of the populations for each species and the large number of seeds required for experimentation, seeds were obtained from several hundred individuals growing at three sites: recently burnt *Banksia* woodlands near Perth (31°56'59"S, 115°51'38"E) or near Pinjar (31°39'27"S, 115°49'17"E) or in a historically-burnt open *Eucalyptus marginata* (jarrah) forest near Pinjarra (32°36'59"S, 115°52'38"E), Western Australia. Seeds were kept dry in cloth bags at room conditions (21°C) between

collection and initiation of experiments, which begun within the first week following collection. *Trachymene* produces a mix of bristly and tuberculate mericarps, which were combined in approximate equal proportions for experiments since embryo growth and germination did not differ between them (Hidayati and Walck, unpubl. data).

General embryo growth and germination test procedures

Every time embryo growth was recorded prior to radicle emergence, embryos from 25 seeds were excised using a razor blade and measured under a dissecting microscope with a micrometer. The length of the seeds from which the embryos were excised was also measured and embryo lengths were converted to a ratio (percentage) of seed length. If a seed had germinated, embryo length was estimated by determining its fully-elongated length inside the seed (i.e. the point of emergence from the seed); thus, the ratio for a germinated seed was 100%. Seeds in a dry condition (i.e. freshly collected or stored in the field or laboratory) were imbibed at room temperature for 24 h before embryos were excised.

To test germination, seeds were placed on two glass microfibre 1.2 μm filter papers in plastic Petri dishes [9.0 cm (diameter) x 1.5 cm (depth)] moistened with one of three solutions: distilled water, 0.67 μM KAR₁ (3-methyl-2*H*-Furo[2,3-*c*]pyran-2-one) synthesized from pyromeconic acid as described in Flematti *et al.* (2005), or 1:10 (v/v) smoke water produced following Dixon *et al.* (1995). The purpose of using the two smoke solutions was to simulate some of the effects of fire under laboratory conditions; we have found consistently that smoke sensitive species respond readily to the concentrations used (e.g. Merritt *et al.*, 2006; Stevens *et al.*, 2007; Commander *et al.*, 2009; Turner *et al.*, 2009; Hidayati *et al.*, 2012).

Three replications of 25 seeds per dish were used in each treatment. Dishes were wrapped with plastic film to reduce water loss, and those in dark treatments were additionally

wrapped in aluminium foil to exclude all light. Protrusion of the radicle was the criterion for germination. Incubation lasted for 4 weeks in light or in darkness to ensure adequate time for embryo growth and radicle emergence; seeds were not examined while they were in darkness. At the end of 4 weeks, germinated seeds were counted and embryos in non-germinated seeds were examined and determined to be viable if they were firm and white (confirmed by tetrazolium tests, Grabe, 1970), or non-viable if they were soft and grey. Germination percentages were determined based on the number of viable seeds.

A temperature- and light-controlled incubator was used to study embryo growth as well as to test germination. It was set at daily alternating (12/12 h) 18/7°C that approximated the mean daily maximum and minimum air temperatures for Perth from May to September (Bureau of Meteorology, 2009). This temperature is optimum for germination of the study species (Hidayati and Walck, unpubl. data) and simulates the cooler wet season when seedling emergence occurs for most species in southwest Western Australia (Merritt *et al.*, 2007). The 12-h photoperiod corresponded to the high-temperature period with the light source being 30 W cool white (400-700 nm) fluorescent tubes with photon flux density at seed level of 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

Patterns of embryo growth and germination

To understand the timing and amount of embryo growth and germination in relation to physiological dormancy loss, if any, over summer and into autumn, we studied seeds placed on/into the soil in the field and seeds stored under laboratory after-ripening conditions. At the start of the experiment, 2008-cohort seeds were separated into batches and the following occurred: (1) the lengths of embryos and seeds were measured for *Levenhookia* and *Trachymene*, (2) germination tests were performed for all four species in light and in darkness at 18/7°C, and

(3) about 500 seeds for the four species were each placed into fine-mesh nylon bags [5 cm (length) $\times$ 5 cm (width)]. The bags of seeds were stored in the field for *Podotrochea*, *Levenhookia*, and *Trachymene* or in the laboratory for all four species, starting on 9 December 2008.

For field storage, we established a site in natural *Banksia* woodland at Kings Park and Botanic Garden; the soil at this site was classified as Karrakatta sands of the Spearwood Dunes (Bennett, 1988). We used 24 bags per species allowing for extra bags in case some were destroyed or lost and placed one bag into a plastic tray [7 cm (depth) $\times$ 15 cm (length) $\times$ 7 cm (width), with drainage holes] filled with soil from the site. In the tray, the bag was either (1) buried at 5-cm depth or (2) placed on top of the soil surface with ca. 1 mm of soil sprinkled on top. Four trays (2 with a buried bag and 2 with a surface bag) each per species were placed into each of six plots [60 cm (length) $\times$ 60 cm (width)]. The plots were placed at 5 m intervals along an unused sand track in the woodland. In each plot, the trays were randomly positioned (with respect to species and bag placement) alongside each other (in a 2 $\times$ 6 design) and sunk into the ground so that the top was even with the surrounding soil surface. All the trays in a plot were covered with metal screen secured to the ground with metal pins to protect against predators. Part of this woodland was burnt in January 2009, and while the immediate site remained non-burnt, two plots were destroyed by vehicles extinguishing the fire. In January, March, and May 2009, a randomly selected buried and surface bag was removed from a plot. After removal, the bag was opened and the number of germinated seeds counted. For non-germinated seeds, (1) lengths of embryos and seeds were measured for *Levenhookia* and *Trachymene* and (2) germination was tested in light on each solution at 18/7°C for all three species.

For laboratory storage, we placed 20 bags above a non-saturated solution of LiCl (370 g L⁻¹) in an air-tight polycarbonate box [14 cm (depth) $\times$ 28 cm (length) $\times$ 28 cm (width); NHP Fibox, Australia]. The atmosphere inside the box was maintained at ca. 50% relative humidity

(Gold and Hay, 2014). The box was kept in a 30°C incubator, simulating summer temperature, with a 12-h photoperiod. One randomly-selected bag for each species was obtained from the box each month between January 2009 and May 2009 (except for *Austrostipa*, which was removed every other month). Every time a bag was removed, (1) lengths of embryos and seeds were measured for *Levenhookia* and *Trachymene* and (2) germination was tested in light and in darkness on each solution at 18/7°C for all four species.

To determine the start of embryo growth and germination at a fine-interval scale, we measured embryos and recorded germination in seeds of *Levenhookia* and *Trachymene* that had been after-ripened in the laboratory. We placed 50 seeds each of these two species into three Petri dishes on each solution at 18/7°C when they were removed after 5 months dry storage at laboratory conditions (in May 2009). At 4-day intervals up to 28 days (or until all seeds germinated), we randomly-selected 8-9 germinated or non-germinated seeds from each dish (to total 25) and (1) recorded if a seed had germinated and (2) measured the length of the embryo and seed. We determined the relative growth rate of the embryos on each solution over the total measurement period: $[(\log_e E_2 - \log_e E_1)/(t_2 - t_1)]$, where E is the embryo length and t is time between two intervals, 1 (in fresh seeds) and 2 (seeds incubated 28 days for *Levenhookia* or 20 days for *Trachymene*) (Walck *et al.*, 1999). Calculations were made for paired seeds on the basis of ranking by embryo length in each of two sequential measurements.

Patterns of seedling emergence

To document the timing and amount of emergence, we used a tunnel-house, located in Kings Park and Botanic Garden (Perth, Western Australia). This tunnel-house had no heating or air-conditioning and had only the roof and the top-half of its sides covered with fibreglass sheeting forming a rain-out shelter. For each species, 200 seeds each were sown on 23 November 2007

(2007 cohort) or on 9 December 2008 (2008 cohort) into eight 10-cm diameter plastic pots. Seeds were sown on the surface and then lightly covered with about 1 mm of soil. The pots were filled with a soil mixture (4:2:1 composted jarrah sawdust:fine river sand:coarse river sand, pH adjusted to 6.0 with addition of 1 kg of lime and 800 g of dolomite per 1 m³ of soil; 2007 cohort) or with fine river sand (2008 cohort) and placed on benches. Before use, the soil mix, sand, and pots were steam-pasteurized for 1 h at 60–70°C.

The soil mix and sand were watered to maximum capacity weekly during summer and early autumn (dry season, September to March), twice a week in mid-autumn (first 2 weeks in April), then daily from late autumn into winter (wet season, last 2 weeks in April to August) to simulate natural precipitation (Turner *et al.*, 2006; Merritt *et al.*, 2007). One week after the start of daily watering, four randomly selected pots were exposed to 1 h of aerosol smoke in late April of 2007 and 2008 [following Dixon *et al.* (1995)], while the other four served as control pots. Following smoking, the pots were returned to the tunnel-house and daily watering of all pots resumed the next day. Experiments in the tunnel-house ended on 26 June 2009, since no new seedlings had emerged for at least 2 weeks prior to this date.

Scoring was done weekly using a magnifying glass to examine the soil surface for signs of emergence. If seedlings (with an emerged radicle and/or cotyledon) were present, they were counted and gently removed using small forceps. We defined emergence seasons as the period between the commencement of twice weekly watering in April and the point where no additional seedling emergence was observed during the wet season. Emergence percentages were calculated based on the number of seeds sown. Parameters estimated from the cumulative emergence curves during each season using GERMINATOR software (Joosen *et al.*, 2010) were: maximum emergence (defined as Gmax in the software), onset of emergence (i.e. time to 1% of

G_{max}), median emergence (i.e. time to 50% of G_{max}), and the spread of emergence (i.e. time between 10 and 90% of G_{max}).

Temperature and precipitation records

Maximum and minimum daily air temperatures throughout the study (23 November 2007 to 26 June 2009) were determined from regional records at the Perth weather station (#009225; Bureau of Meteorology, 2009), approximately 5 km from Kings Park and Botanic Garden. Data loggers (Tinytag Plus 2, Gemini Data Loggers, Chichester, West Sussex, UK) were placed on top of the soil surface in a pot in the tunnel-house and on the soil surface in the woodland in Kings Park and Botanic Garden and recorded soil temperatures on an hourly basis. Data loggers were located approximately 10 and 50 m from our study pots or plots, respectively. Daily precipitation during the study was obtained from the Perth weather station, as previously described.

Statistical analyses

Generalized linear models (GLMs; binomial error structure, logit link function) or analyses of variances (ANOVAs) followed by protected least significance difference tests (PLSD, $P=0.05$) were used to examine the data (SPSS, 2012; R Core Team, 2014). For GLMs, the main factor terms and their two-way interactions were added sequentially followed by a Wald-test to validate the final model (Fox 2015). All factors in the GLMs were analysed as discrete terms. Germination from seeds was examined using GLM with species, time for storage, location (buried, surface), and solution as factors in the field storage experiment and with species, time for storage, light regime (light, dark), and solution as factors in the laboratory experiment. Relative growth rates for embryos were examined using a Welch's ANOVA with species and solution (water, KAR₁, smoke water) as factors. To analyze patterns of emergence, the effects of species

and smoke treatment on onset, median and spread of emergence were examined by a two-way ANOVA. Emergence was analysed using GLMs with species and smoke treatment as factors.

Results

Patterns of embryo growth and germination

Embryo lengths in fresh *Levenhookia* seeds were 10% the length of the seed and those in *Trachymene* seeds were 11%. Between January 2009 and May 2009, *Levenhookia* embryos remained 10-14% the length of the seed, with no consistent growth, regardless if they were stored in the field (buried in soil or placed on the soil surface) or in laboratory after-ripening conditions. *Trachymene* embryos in field-stored seeds increased linearly up to 32% (while buried) or up to 17% (on soil surface) by May 2009 but were 11-13% under laboratory conditions.

For seeds that were removed from storage in the field, germination was highly species specific and dependent on storage location in the field, time when exhumed and tested, and test solution (all factors and interactions, Wald $X^2 \geq 11.879$, $P \leq 0.018$). No freshly-matured seeds of these species germinated during 4 weeks of incubation at 18/7°C in light on any solution (Fig. 1), and no germinated seeds were observed within bags exhumed in January, March, and May 2009. When removed from buried or surface bags from January to May 2009 and incubated at 18/7°C in light, germination of *Podotheca* and *Trachymene* increased to 73-100% on all solutions. *Levenhookia* germination increased to 56-87% on all solutions when removed from surface bags but to 8, 28, and 76% on water, KAR₁, and smoke water, respectively, from buried bags. Germination of all three species increased progressively with time, but at different rates; for seeds of *Trachymene* > 90% germination was recorded as early as January (2 months old), for seeds of *Podotheca* >80% germination was recorded from March (4 months old), and for seeds of *Levenhookia* >80% germination was not recorded until May (6 months old).

Germination of laboratory-stored seeds was highly species specific and dependent on time when tested, test solution and test light regime (all factors and interactions, Wald $X^2 \geq 21.358$, $P \leq 0.006$). Fresh seeds of *Austrostipa*, *Podotheca*, and *Levenhookia* germinated $\leq 5\%$ during incubation at $18/7^\circ\text{C}$ in light and darkness, regardless of solution (Fig. 2). In contrast, *Trachymene* fresh seeds germinated to $\leq 1\%$ in light on all solutions but to 92, 54, and 5% in darkness on smoke water, KAR₁, and water, respectively. Germination for *Podotheca* and *Trachymene* increased to 69-100% on all solutions in light, but for *Levenhookia* to only 9-22%. In darkness, germination for these three species increased up to 100% on KAR₁ and on smoke water but up to only 9% (*Podotheca*, *Levenhookia*) or 72% (*Trachymene*) on water. Seeds of *Austrostipa* only germinated up to 20% regardless of light regime or solution.

Once incubated for germination, relative growth rates for embryos in laboratory-stored, after-ripened seeds differed among solutions ($F = 7.887$, $P = 0.001$) and between species ($F = 766.446$, $P < 0.0001$), and the response to solutions was species specific (interaction, $F = 8.284$, $P < 0.0001$). Growth rates of *Levenhookia* embryos differed among solutions as follows: smoke water ($0.0415 \pm 0.0063 \text{ mm mm}^{-1} \text{ d}^{-1}$) = KAR₁ ($0.0390 \pm 0.0061 \text{ mm mm}^{-1} \text{ d}^{-1}$) > water ($0.0138 \pm 0.0024 \text{ mm mm}^{-1} \text{ d}^{-1}$). For *Levenhookia* seeds, embryo growth was most evident by day 12 on all solutions (Fig. 3). Germination of *Levenhookia* seeds was evident on day 16 for KAR₁ and on day 20 for smoke water; no seeds germinated on water. Growth rates for *Trachymene* embryos did not differ among solutions: water ($0.1181 \pm 0.0012 \text{ mm mm}^{-1} \text{ d}^{-1}$), KAR₁ ($0.1165 \pm 0.0011 \text{ mm mm}^{-1} \text{ d}^{-1}$), and smoke water ($0.1186 \pm 0.0013 \text{ mm mm}^{-1} \text{ d}^{-1}$). Embryo growth of *Trachymene* seeds started noticeably on day 8 regardless of solution, and germination was obvious on day 12 for seeds treated with both KAR₁ and smoke water, but on day 16 for water (Fig. 3).

Patterns of seedling emergence

For the 2007 cohort during the first germination season, onset, median, and spread of emergence ($F \geq 7.511$, $P \leq 0.018$) and maximum emergence (Wald $X^2 = 72.522$, $P < 0.0001$) varied significantly between species and between treatments. While the responses to smoke for the onset and spread of emergence (interaction, $F \geq 9.460$, $P \leq 0.010$) and maximum emergence (Wald $X^2 = 36.059$, $P < 0.0001$) varied between species, the response for median of emergence was similar between species ($F = 0.106$, $P = 0.751$). During the second germination season, no treatment effect occurred for any of the four emergence parameters ($F \leq 2.940$, $P \geq 0.112$; Wald $X^2 = 3.427$, $P = 0.064$). All four parameters varied between species ($F \geq 14.463$, $P \leq 0.002$; Wald $X^2 = 40.895$, $P < 0.0001$), but for only onset and spread of emergence (interaction, $F \geq 6.670$, $P \leq 0.024$) did species have dissimilar responses between treatments ($F = 3.543$, $P = 0.084$ for median emergence, Wald $X^2 = 0.822$, $P = 0.365$ for maximum emergence). Onset and median emergence was significantly earlier for *Podotheca* than for *Trachymene* during both emergence seasons (Fig. 4, Supplementary Data Fig. S1); seeds of *Podotheca* emerged almost immediately (within 3 d), whilst seeds of *Trachymene* required c. 30 d to emerge. This discrepancy in onset and median emergence times between species was reduced in the second season, but for the most part, seeds of *Podotheca* still emerged significantly earlier than seeds of *Trachymene*. Smoking delayed the onset of emergence for *Trachymene* seeds only during the first season, whereas it slightly delayed median emergence for *Podotheca* in both seasons. There was a smaller difference in the spread of emergence between the two species in both emergence seasons. However, smoke greatly spread emergence of *Podotheca* seeds during the first season and did so to a lesser extent for *Trachymene* in the second season. Smoke treatment induced significantly greater maximum emergence of both species during the first emergence season but not the second season.

For the 2008 cohort, no treatment effect occurred for any of the four emergence parameters ($F \leq 0.480$, $P \geq 0.495$; Wald $X^2 = 2.029$, $P = 0.154$). All four parameters varied

among species ($F \geq 17.237$, $P < 0.0001$; Wald $X^2 = 176.615$, $P < 0.0001$), but for only median emergence did species have dissimilar responses between treatments (interaction, $F = 3.601$, $P = 0.028$; $F \leq 1.537$, $P \geq 0.230$ and Wald $X^2 = 5.281$, $P = 0.152$ for the other three parameters). Onset of emergence was significantly earlier for *Austrostipa* and *Podotheca* compared to *Levenhookia* and *Trachymene* (Fig. 4, Supplementary Data Fig. S2). Median emergence varied among the species as (earliest to latest): *Podotheca* < *Austrostipa* = *Trachymene* < *Levenhookia*. Onset of emergence, median emergence, and total emergence was unaffected by smoke for all species except *Levenhookia*; for *Levenhookia* seeds smoke significantly reduced the median emergence time and significantly increased maximum emergence.

When comparing the first season emergence characteristics of *Podotheca* and *Trachymene* seeds collected in 2007 to seeds collected in 2008, seeds of *Podotheca* in the 2008 cohort had a greatly delayed onset of emergence (*c.* 18 days versus *c.* 3 days), and a much greater maximum emergence (*c.* 92% versus *c.* 32%), than those in the 2007 cohort. Such differences between seed batches collected in 2007 and 2008 were not evident in *Trachymene* seeds.

Temperature and precipitation records

Mean daily maximum and minimum temperatures during summer (December to March) 2007-08 and 2008-09 were 30.7 and 17.6°C, respectively, at the regional level and 40.2 and 17.7°C, respectively, in the tunnel-house (Supplementary Data Fig. S1). Seeds of all species emerged in 2008 and 2009 during April to May when mean daily maximum and minimum temperatures were 24.1 and 10.8°C, respectively, at the regional level and 31.8 and 12.2°C, respectively, in the tunnel-house (Supplementary Data Figs S1, S2). Our daily watering in the tunnel-house commenced at about the same time as the regional wet season during 2008, but it was earlier as compared with regional precipitation in 2009 (Supplementary Data Fig. S1).

Discussion

Fresh seeds of all four study species were dormant upon release in November and after-ripening to overcome physiological dormancy subsequently occurred at summer conditions in all species, during which no embryo growth was observed for the two species (*Levenhookia* and *Trachymene*) with under-developed embryos (Fig. 5). As dormancy loss proceeded (albeit very little in *Austrostipa*), germination increased in light and darkness on all three solutions at a simulated winter temperature of 18/7°C. When seeds were tested for germination at simulated summer temperatures (33/18°C), very low germination occurred, indicating that a rain event during summer would not trigger germination (Hidayati and Walck, unpubl. data). In addition to this thermal suppression of germination, intermittent summer rain would not maintain soil moisture long enough to initiate embryo growth in *Levenhookia* and *Trachymene* seeds – which required at least 8-12 days of high moisture conditions (at winter temperatures). Thus, both of these processes ensure that sporadic summer rainfall does not lead to a false germination event.

An after-ripening requirement for loss of physiological dormancy is reported for a large number of species (Baskin and Baskin, 2014), including many belonging to families well represented in the Australian flora (Merritt *et al.*, 2007). In our study, the degree of physiological dormancy loss was highly dependent on the location of after-ripening. The difference in after-ripening between seeds buried versus placed on the soil surface in the woodland was not as dramatic as that between seeds after-ripened in the woodland versus laboratory. This difference was most evident in *Levenhookia*. For this species, at the end of the after-ripening period, seeds placed on the soil surface in the woodland germinated to high percentages on water and smoke water and moderate percentages on KAR₁, seeds buried in the woodland germinated to high percentages on smoke water and low percentages

on water and KAR₁, and seeds kept in an incubator germinated to low percentages on all three solutions. Thus, after-ripening on was most effective on the soil surface, perhaps due to the warmer and/or dryer conditions compared to burial or the laboratory. Dormancy loss via after-ripening occurs more rapidly at increased temperature (Steadman *et al.*, 2003) and also with alternating wet/dry conditions for some species (Hidayati *et al.*, 2012), as would be experienced for seeds on the soil surface. In contrast, seeds of *Austrostipa* germinated to low percentages, regardless of after-ripening condition in our study. This result is surprising since Baker *et al.* (2005) reported high germination at constant temperatures of 5, 10, and 15°C in the same species following dry storage at 15°C. However, no seeds germinated at an alternating temperature of 25/15°C in their study. We also used an alternating temperature regime and perhaps this was the reason for our very low germination.

At the start of the autumn-winter wet season when seeds were least dormant, seedlings emerged earlier from seeds with fully-developed embryos than from seeds with under-developed embryos (Fig. 5). This pattern was particularly evident for the onset of germination in the tunnel-house: by 30 days for *Podotheca* versus *Trachymene* in the 2007 cohort during the first emergence season and by ca. 10 days for *Austrostipa* and *Podotheca* versus *Levenhookia* and *Trachymene* in the 2008 cohort. A similar pattern was observed in the field. In remnant seed bags kept on the soil surface (following the completion of our experiments), high percentages of *Podotheca* seedlings were observed in late May-early June 2009, 2 weeks before seedlings of *Levenhookia* and *Trachymene* were observed (Hidayati and Walck, unpubl. data). The delay in emergence observed in the tunnel-house and field was due to embryo growth that had to occur inside seeds with under-developed embryos before radicles emerged versus no prior growth for seeds with fully-developed embryos (Fig. 5).

We thought that the timing of induction of embryo growth, and the rate of embryo growth, would differ amongst seeds of species with under-developed embryos, leading to seedling emergence being more spread out during the wet season as compared to fully-developed embryos. Barring differences between smoked and non-smoked treatments, the spread of emergence for the 2007 cohort was not different between embryo types during the first emergence season but was so during the second emergence season. If we consider the germination of seeds in light following after-ripening in the laboratory as a degree of physiological dormancy, then the hierarchy of species (from shallowest to deepest dormancy) would be: *Trachymene* > *Podotheca* > *Levenhookia* $\approx$ *Austrostipa*. Thus, for the 2008 cohort, spread of emergence was related more to degree of (physiological) dormancy than to embryo growth. The two deep-dormant species (*Austrostipa*, *Levenhookia*) had a wider spread than the two shallow-dormant species (*Podotheca*, *Trachymene*). In addition to physiological dormancy break during the summer, additional dormancy release may have occurred during the autumn-winter period to allow continued emergence of seedlings in *Austrostipa* and *Levenhookia*.

We suggested that seedlings from seeds treated with smoke would emerge earlier than those from untreated seeds regardless of embryo type. In our study where differences occurred, smoked seeds emerged later in three instances and earlier in only one case. This is in contrast to a number of studies where the rate of emergence was enhanced in seeds treated with smoke (Dixon *et al.*, 1995; Roche *et al.*, 1997a, b). The effects of smoke were more noticeable in terms of number of emergents rather than timing of emergence: with some smoked treatments having more emergents than non-smoked ones. Our study demonstrated the loss of smoke reactivity after a previous smoke application. In the second emergence season for the 2007 cohort, smoke did not enhance emergence, strongly contrasting to the first season. Apparently, seeds then respond to regular seasonal cues outside of the previous ‘fire’ event. We observed a similarly

reduced requirement for a smoke cue over time in *Hibbertia* species, speculating that it might be a bet-hedging strategy in the event a fire does not occur for some time (Hidayati *et al.*, 2012).

Unlike other studies that have shown enhanced seedling growth (Daws *et al.*, 2007; Nelson *et al.*, 2012; Zhou *et al.*, 2012), KAR₁ and smoke water did not increase embryo growth consistently between our two species with under-developed embryos. While under-developed embryos grew faster in seeds placed on KAR₁ or smoke water than on water for *Levenhookia*, embryos grew at the same rate on all solutions for *Trachymene*. The faster embryo growth for *Levenhookia* resulted in faster germination on KAR₁ and smoke water than on water. These differences again appear to be driven by degrees of physiological dormancy with *Levenhookia* being more deeply dormant than *Trachymene*.

Conclusions

Physiological dormancy was overcome in seeds of four fire ephemerals during summer, enabling them to be non-dormant at the start of the autumn-winter wet season. However, no embryo growth was observed in seeds with under-developed embryos during summer. Once rains started, seedling emergence from seeds of two species with fully-developed embryos occurred earlier than from seeds of two other species with under-developed embryos. In a non-consistent manner among our study species, smoke and smoke compounds influenced the rate of embryo growth and amount of germination, suggesting a strong interaction between seed dormancy status and sensitivity to smoke. Seeds of these species use thermal suppression and/or embryo growth as a means to avoid summer false breaks, which would be lethal since follow-up rains rarely occur in Mediterranean systems. A shift to reliance of regular seasonal cues for emergence and a reduced requirement for smoke occurred in the second germination season after a fire.

Supplementary data. Supplementary data are available online at _____ and consist of the following. Figure S1: Temperatures from a regional weather station (Perth, Western Australia) and from a tunnel-house, regional total daily precipitation, and emergence of seedlings from two fire ephemerals. Figure S2: Temperatures from a regional weather station (Perth, Western Australia) and from the tunnel-house, regional total daily precipitation, and emergence of seedlings from four fire ephemerals.

Acknowledgement. The authors thank Wolfgang Lewandrowski for his assistance with statistical analyses.

Financial support. SNH was supported by the Australian Research Council's Linkage Projects funding scheme LP0455415 and DJM by the Botanic Gardens and Parks Authority – Alcoa of Australia Limited Seed Conservation Partnership. The research was conducted under the auspices of the Millennium Seed Bank Project, Kew, which is supported by the UK Millennium Commission, the Wellcome Trust and Orange plc.

Conflicts of interest. None.

References

- Baker KS, Steadman KJ, Plummer JA, Dixon KW** (2005) Seed dormancy and germination responses of nine Australian fire ephemerals. *Plant and Soil* **277**, 345–358.
- Bart RR, Tague CL, Dennison, PE** (2017) Modeling annual grassland phenology along the central coast of California. *Ecosphere* 8(7):e01875. 10.1002/ecs2.1875.
- Barrett R, Tay EP** (2005) *Perth plants: a field guide to the bushland and coastal flora of Kings Park and Bold Park, Perth, Western Australia*. Perth: Botanic Gardens and Parks Authority.
- Baskin CC, Baskin JM** (2014) *Seeds: ecology, biogeography, and evolution of dormancy and germination*. 2nd edn. San Diego: Academic Press.
- Bennett EM** (1988) *The bushland plants of Kings Park, Western Australia*. Perth: Kings Park Board.
- Brown NAC** (1993) Promotion of germination of fynbos seeds by plant-derived smoke. *New Phytologist* **123**, 575–583.
- Bureau of Meteorology** (2009) Climate statistics for Australian locations. Commonwealth of Australia. Available at: <http://www.bom.gov.au/> [Accessed 10 December 2009].
- Carta A, Hanson S, Müller JV** (2016) Plant regeneration from seeds responds to phylogenetic relatedness and local adaptation in Mediterranean *Romulea* (Iridaceae) species. *Ecology and Evolution* **6**, 4166–4178.
- Chiwocha SDS, Dixon KW, Flematti GR, Ghisalberti EL, Merritt DJ, Nelson DC, Riseborough, JM, Smith SM, Stevens JC** (2009) Karrikins: a new family of plant growth regulators in smoke. *Plant Science* **177**, 252–256.

- 488 **Commander LE, Merritt DJ, Rokich DP, Dixon KW** (2009) Seed biology of Australian arid
 489 zone species: Germination of 18 species used for rehabilitation. *Journal of Arid*
 490 *Environments* **73**, 617–625.
- 491 **Daws MI, Davies J, Pritchard HW, Brown NAC, Van Staden J** (2007) Butenolide from plant-
 492 derived smoke enhances germination and seedling growth of arable weed species. *Plant*
 493 *Growth and Regulation* **51**, 73–82.
- 494 **Dixon KW, Roche S, Pate JS** (1995) The promotive effects of smoke derived from burnt native
 495 vegetation on seed germination of Western Australian plants. *Oecologia* **101**, 185–192.
- 496 **Flematti GR, Ghisalberti EL, Dixon KW, Trengove RD** (2005) Synthesis of the seed
 497 germination stimulant 3-methyl-2H-furo[2,3-*c*]pyran-2-one. *Tetrahedron Letters* **46**,
 498 5719–5721.
- 499 **Fox J** (2015) Applied regression analysis and generalized linear models. Sage Publications, Los
 500 Angeles.
- 501 **Ghebrehiwot HM, Kulkarni MG, Kirkman KP, Van Staden J** (2009) Smoke solutions and
 502 temperature influence the germination and seedling growth of South African mesic
 503 grassland species. *Rangeland Ecology and Management* **62**, 572–578.
- 504 **Gold K, Hay F** (2014) *Equilibrating seeds to specific moisture levels*. Technical Information
 505 Sheet 9. Kew: Millennium Seed Bank Project.
- 506 **Grabe DF** (1970) *Tetrazolium testing handbook for agricultural seeds*. Contribution Number 29
 507 to the Handbook on seed testing. New Jersey: Association of Official Seed Analysts:
 508 North Brunswick.
- 509 **Grant CD, Loneragan WA** (1999) The effects of burning on the understorey composition of 11-
 510 13 year-old rehabilitated bauxite mines in Western Australia. *Plant Ecology* **145**, 291–
 511 305.

- 512 **Hidayati SN, Walck JL, Merritt DJ, Turner SR, Turner DW, Dixon KW** (2012) Sympatric
 513 species of *Hibbertia* (Dilleniaceae) vary in dormancy break and germination
 514 requirements: implications for classifying morphophysiological dormancy in
 515 Mediterranean biomes. *Annals of Botany* **109**, 1111–1123.
- 516 **Hopper SD, Brown AP, Marchant NG** (1997) Plants of Western Australian granite outcrops.
 517 *Journal of the Royal Society of Western Australia* **80**, 141–158.
- 518 **Joosen RVL, Kodde J, Willems LAJ, Ligterink W, van der Plas LHW, Hilhorst HWM**
 519 (2010) GERMINATOR: a software package for high-throughput scoring and curve fitting
 520 of *Arabidopsis* seed germination. *The Plant Journal* **62**, 148–159.
- 521 **Lamont B** (1985) Gradient and zonal analysis of understorey suppression by *Eucalyptus*
 522 *wandoo*. *Vegetatio* **63**, 49–66.
- 523 **Merritt DJ, Turner SR, Clarke S, Dixon KW** (2007) Seed dormancy and germination
 524 stimulation syndromes for Australian temperate species. *Australian Journal of Botany* **55**,
 525 336–344.
- 526 **Merritt DJ, Kristiansen M, Flematti G, Turner SR, Ghisalberti EL, Trengove RD, Dixon**
 527 **KW** (2006) Effects of a butenolide present in smoke on light-mediated germination of
 528 Australian Asteraceae. *Seed Science Research* **16**, 29–35.
- 529 **Miller BP, Dixon KW** (2014) Plants and fire in Kwongan vegetation. pp. 147–169 in Lambers H
 530 (ed), *Plant life on the sandplains in southwest Australia, a global biodiversity hotspot*.
 531 Perth: University of Western Australia Press.
- 532 **Nelson DC, Flematti GR, Ghisalberti EL, Dixon KW, Smith SM** (2012) Regulation of seed
 533 germination and seedling growth by chemical signals from burning vegetation. *Annual*
 534 *Review of Plant Biology* **63**, 107–130.

- 535 **R Core Team** (2014) R: A language and environment for statistical computing. R Foundation for
 536 Statistical Computing, Vienna, Austria, URL: <http://www.R-project.org/>.
- 537 **Roche S, Dixon KW, Pate JS** (1997a) Seed ageing and smoke: partner cues in the amelioration
 538 of seed dormancy in selected Australian native species. *Australian Journal of Botany* **45**,
 539 783–815.
- 540 **Roche S, Koch JM, Dixon KW** (1997b) Smoke enhanced seed germination for mine
 541 rehabilitation in the southwest of Western Australia. *Restoration Ecology* **5**, 191–203.
- 542 **Rundel PW, Parsons DJ** (1984) Post-fire uptake of nutrients by diverse ephemeral herbs in
 543 chamise chaparral. *Oecologia* **61**, 285–288.
- 544 **SPSS** (2012) *IBM SPSS Statistics*. Version 21. Armonk, New York: IBM Corp.
- 545 **Steadman KJ, Crawford AD, Gallagher RS** (2003) Dormancy release in *Lolium rigidum* seeds
 546 is a function of thermal after-ripening time and seed water content. *Functional Plant*
 547 *Biology* **30**, 345–352.
- 548 **Stevens JC, Merritt DJ, Flematti GR, Ghisalberti EL, Dixon KW** (2007) Seed germination of
 549 agricultural weeds is promoted by the butenolide 3-methyl-2H-furo[2,3-c] pyran-2-one
 550 under laboratory and field conditions. *Plant and Soil* **298**, 113–24.
- 551 **Turner SR, Merritt DJ, Ridley EC, Commander LE, Baskin JM, Baskin CC, Dixon KW**
 552 (2006) Ecophysiology of seed dormancy in the Australian endemic species
 553 *Acanthocarpus preissii* (Dasypogonaceae). *Annals of Botany* **98**, 1137–1144.
- 554 **Turner SR, Merritt DJ, Renton MS, Dixon KW** (2009) Seed moisture content affects
 555 afterripening and smoke responsiveness in three sympatric Australian native species from
 556 fire prone environments. *Austral Ecology* **34**, 866–877.
- 557 **Vandelook F, Verdú M, Honnay O** (2012) The role of seed traits in determining the
 558 phylogenetic structure of temperate plant communities. *Annals of Botany* **110**, 629–636.

- 559 **Verdú M** (2006) Tempo, mode and phylogenetic associations of relative embryo size evolution
560 in angiosperms. *Journal of Evolutionary Biology* **19**, 625–634.
- 561 **Vivrette NJ** (1995) Distribution and ecological significance of seed-embryo types in
562 Mediterranean climates in California, Chile, and Australia. pp. 274–288 in Arroyo MTK,
563 Zedler PH, Fox MD (eds), *Ecology and biogeography of Mediterranean ecosystems in*
564 *Chile, California, and Australia*. New York: Springer-Verlag,
- 565 **Walck JL, Baskin JM, Baskin CC** (1999) Relative competitive abilities and growth
566 characteristics of a narrowly endemic and a geographically widespread *Solidago* species
567 (Asteraceae). *American Journal of Botany* **86**, 820–828.
- 568 **Zhou J, Kulkarni MG, Huang LQ, Guo LP, Van Staden J** (2012) Effects of temperature,
569 light, nutrients and smoke-water on seed germination and seedling growth of *Astragalus*
570 *membranaceus*, *Panax notoginseng* and *Magnolia officinalis* – highly traded Chinese
571 medicinal plants. *South African Journal of Botany* **79**, 62–70.
- 572

Figure 1. Mean ($\pm$ SE, SE shown if $\geq 5\%$) germination of three fire ephemerals on water, KAR₁, and smoke water during 4 weeks of incubation in light at 18/7°C following 0 (fresh)-5 months *in situ* soil storage under *Banksia* woodland conditions (either buried in soil or placed on the soil surface). Seeds were buried on 9 December 2008. Means with dissimilar letters within each panel are significantly different among solutions for the May 2009 test (PLSD, $P \leq 0.05$); those without letters are not. Data based upon three replicates of 25 seeds each.

Figure 2. Mean ($\pm$ SE, SE shown if $\geq 5\%$) germination of four fire ephemerals on water, KAR₁, and smoke water during 4 weeks in two light regimes at 18/7°C following 0 (fresh)-5 months storage under standardized laboratory conditions (polyethylene box with 50% relative humidity in an incubator at 30°C). Seeds were placed into the after-ripening box on 9 December 2008. Means with dissimilar letters within each panel are significantly different among solutions for the May 2009 test (PLSD, $P \leq 0.05$); those without letters are not. Data based on three replicates of 25 seeds each.

Figure 3. Box plots and mean ($\pm$ SE, SE shown if $\geq 5\%$) germination for after-ripened seeds of two fire ephemerals with small underdeveloped embryos incubated for up to 28 days on water, KAR₁, and smoke water. Embryos in 25 seeds were measured for each treatment at each time, and germination is based upon three replicates of 25 seeds each.

Figure 4. Mean ($\pm$ SE) time to onset (1%) of emergence, to median (50%) emergence, and spread (time interval) between 10 and 90% emergence of fire ephemerals exposed to aerosol smoke during the first and second emergence seasons for the 2007 cohort and during the first

season for the 2008 cohort (for additional details see Supplementary Data Figs S1, S2). Means with dissimilar letters within each graph are significantly different (PLSD, $P \leq 0.05$); those with similar letters are not.

Figure 5. Conceptual diagram showing seeds with different types of embryos (fully developed and under-developed) in relation to key life stage transitions from seed dispersal to germination according to seasons. Fire (indicated by flames) could occur at any time primarily in summer. Seeds with under-developed embryos germinate later than those with fully-developed embryos due to a period of embryo growth. Under-developed embryos must develop and elongate (dashed arrow) inside the seed before radicle emergence (blocked arrow).

Figure 1

Figure 2

1

Figure 3

2

3

Figure 4

Figure 5

Figure S1

Figure S2

SUPPLEMENTARY MATERIAL

Temporal dynamics of seedling emergence among fire ephemerals: importance of initial embryo size

Hidayati S.N., Merritt D.J., Turner S.R., Dixon K.W. and Walck J.L.

Supplementary data consist of the following:

Figure S1. Maximum and minimum daily temperatures from a regional weather station (Perth, Western Australia) and from a tunnel-house, regional total daily precipitation, and emergence (mean $\pm$ SE, SE shown if $\geq 5\%$) of seedlings from two fire ephemerals. Seeds were collected at maturity and sown on soil in November 2007 in a tunnel-house with ambient temperatures. Treatment pots were exposed to aerosol smoke (S) in April 2008 and 2009; control pots were not (NS). Soil was watered daily or twice weekly from April to August (hatched areas) or it was watered weekly at all other times of the year (non-hatched areas). Data based on four replicates of 200 seeds each.

Figure S2. Maximum and minimum daily temperatures from a regional weather station (Perth, Western Australia) and from the tunnel-house, regional total daily precipitation, and emergence (mean $\pm$ SE, SE shown if $\geq 5\%$) of seedlings from four fire ephemerals. Seeds were collected at maturity and sown on soil in December 2008 in a tunnel-house with ambient temperatures. Treatment pots were exposed to aerosol smoke (S) in April 2009; control pots were not (NS).

24 Soil was watered daily or twice weekly from April to June (hatched area) or it was watered
25 weekly at all other times of the year (non-hatched areas). Data based on four replicates of 200
26 seeds each.

27