Author Manuscript

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/pai.13063](https://doi.org/10.1111/pai.13063)

This article is protected by copyright. All rights reserved

DR JONATAN LEFFLER (Orcid ID : 0000-0001-5674-8462)

Article type : Original

TITLE PAGE

Progressive increase of FcεRI expression across several PBMC subsets is associated with atopy and atopic asthma within school-aged children

Jonatan Leffler PhD^{1,*}, James F. Read BSc (Hons)^{1,2,*}, Anya C. Jones PhD^{1,2}, Danny Mok PhD¹, Elysia M. Hollams PhD¹, Ingrid A. Laing PhD^{1,2}, Peter N. Le Souef MD^{1,2}, Peter D. Sly MD³, Merci M.H. Kusel MBBS¹, Nicholas H. de Klerk PhD¹, Anthony Bosco PhD¹, Patrick G. Holt PhD^{1,3}, Deborah H. Strickland PhD^{1,#}

¹Telethon Kids Institute, University of Western Australia, WA, Australia

²School of Medicine, University of Western Australia, WA, Australia

³Child Health Research Centre, University of Queensland, QLD, Australia

*Jonatan Leffler and James Read should be considered joint first author.

Running title: FcεRI expression is associated with atopic asthma

#Corresponding author

A/Prof. Deborah H. Strickland

Telethon Kids Institute

Northern Entrance, Perth Children's Hospital

15 Hospital Avenue

Nedlands, WA 6009, Australia

Email: Deb.Strickland@telethonkids.org.au

Phone: +61 8 6319 1528

Fax +61 8 6319 1777

Word count: 2543 **Number of tables:**1 **Number of figures:** 4

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

JL, JR, ACJ, DM, EMH performed the experiments and analysed data. JL, JR and NHK performed statistical analysis. JL and JR drafted the manuscript. IAL, PNLS, PDS, MMHK, AB, PGH, DHS assisted in conceiving the project, data interpretation and manuscript editing. All authors read and approved the final version of the manuscript.

FINANCIAL SUPPORT

Authors would like to acknowledge the following funding bodies; Australian National Health and Medical Research Council, University of Western Australia and Telethon Kids Institute. JL is supported by a Fellowship from the Swedish Society for Medical Research, AB is supported by a Fellowship from the Simon Lee Foundation and DHS is supported by a Telethon Kids Institute Fellowship.

ABSTRACT

Background Antigen specific IgE binds the Fcε receptor I (FcεRI) expressed on several types of immune cells, including dendritic cells (DC). Activation of FcεRI on DCs in atopy has been shown to modulate immune responses that potentially contribute to asthma development. However, the extent to which DC subsets differ in

FcεRI expression between atopic children with or without asthma is currently not clear.

This study aimed to analyse the expression of FcεRI on peripheral blood mononuclear cells (PBMC) from atopic children with and without asthma, and non-atopic/non-asthmatic age-matched healthy controls.

Methods We performed multiparameter flow cytometry on PBMC from 391 children across three community cohorts and one clinical cohort based in Western Australia.

Results We confirmed expression of FcεRI on basophils, monocytes, plasmacytoid and conventional DCs, with higher proportions of all cell populations expressing FcεRI in atopic compared to non-atopic children. Further, we observed that levels of FcεRI expression was elevated across plasmacytoid and conventional DC as well as basophils in atopic asthmatic compared to atopic non-asthmatic children also after adjusting for serum IgE levels.

Conclusion Our data suggest that the expression pattern of FcεRI on DC and basophils differentiates asthmatic and non-asthmatic atopic children. Given the significant immune modulatory effects observed as a consequence of FcεRI expression, this altered expression pattern is likely to contribute to asthma pathology in children.

Key words FcεRI expression, dendritic cells, basophils, monocytes, atopic asthma, IgE, childhood asthma, PBMC, flow cytometry

BACKGROUND

Allergic IgE sensitisation and early life respiratory infections are major risk factors for development of atopic asthma during childhood¹. Asthma affects 300 million

people, consists of several phenotypes and atopic asthma is the most common in children. The disease manifests initially as wheeze which is triggered at the time of respiratory viral infection, but in the majority of children the wheezy phenotype is only transient and resolves during the preschool years, whereas in a smaller subset dominated by sensitized children it persists and intensifies and eventually develops into full-blown atopic asthma. At the time of puberty, asthma transitions from being most common in boys to becoming more common in females².

IgE sensitization is characterised by induction of allergen-specific T helper 2 (Th2) immune memory and the production of corresponding specific IgE. Allergen specific IgE binds Fcε receptors that are expressed on several cell types within the airways and circulation including mast cells, basophils, monocytes and dendritic cells (DC)³. Fcε receptor I (FcεRI) binds IgE with high affinity and consists of three subunits; the IgE binding α-subunit and the signalling/regulatory β and γ subunits. Mast cells and basophils express a αβγ₂ tetrameric complex, whereas DC expresses a αγ₂ trimeric complex⁴. Crosslinking of IgE-bound FcεRI complexes by allergen leads to release of inflammatory mediators by mast cells/basophils and this is largely dependent on signal amplification by the β-subunit⁵. Thus, the function of trimeric FcεRI on DCs and monocytes is more complex. On the one hand conventional DC (cDC) utilize surface FcεRI expression to enhance their allergen capture functions⁶, which can result in multi-logfold increased capacity to re-activate Th2-memory cells⁷ but have also been reported to be involved in IgE clearance through internalisation of serum IgE⁸ and induction of immune modulatory mechanisms such as tolerance⁹. Conversely, cross-linking of FcεRI on plasmacytoid DC (pDC) results in reduced type I-interferon signalling and impaired anti-viral responses^{10,11}. Expression of FcεRI on

pDCs and cDC has been reported to be dependent on serum IgE levels³, and in sensitised children with elevated levels of specific IgE (above 0.35 IU/mL), respiratory viral infections are often more prevalent/severe compared to non-atopic children as reviewed¹². The proportion of FcεRI⁺ DCs, and FcεRI expression intensity, is generally higher in atopic and/or asthmatic individuals compared to healthy controls but variation exists between studies^{8,11}. The association of FcεRI expression and total serum IgE levels appears strongest for basophils and pDCs^{8,13}.

Although FcεRI expression has been reported to be elevated in atopic and/or asthmatic individuals compared to healthy controls¹³⁻¹⁵, there are currently few studies that have investigated whether FcεRI expression across cell populations within the PBMC fraction can differentiate atopic asthmatics from atopic but non-asthmatic individuals. In the current study we aimed to identify and characterise FcεRI expressing cell populations in atopic children with or without asthma and healthy controls across four different allergy/asthma cohorts based in Western Australia. Our findings suggest that FcεRI expression may have clinical relevance to the pathology of childhood asthma, and has potential to be used as a biomarker related to disease susceptibility and phenotype.

METHODS

Study cohorts

A subset of school-age individuals (age range 7-15 years) from four independent cohorts based in Perth, Western Australia, were analysed. The participant numbers and clinical parameters are summarised in Table I. The Childhood Asthma Study (CAS) cohort is a birth cohort which investigated respiratory infections in children

from families with a history of allergy or asthma and thus with a high risk for developing atopic disease at the age of 10 years¹⁶. The Western Australian Twin study of Child Health (WATCH) was established as a population-based twin register to examine the genetics of asthma and attention deficit hyperactivity disorder. This was extended to include a follow-up subsample of participant families enriched for (but not exclusively) physician-diagnosed asthmatics¹⁷ at the median age of 10 years. The Western Australian Pregnancy Cohort (Raine) is a large (>2000 children) unselected general birth cohort where approximately 10% of participants developed asthma for a period during their first 14 years of life^{18,19}. From this cohort, 79 atopic individuals were included in the analysis and selected so that 40 individuals also had an asthma diagnosis at the age of 14 years. Peripheral blood samples from the CAS, WATCH, and Raine cohorts were collected when the children did not display any clinical symptoms of airway disease. The Mechanism of Acute Viral Respiratory Infection in Children (MAVRIC) cohort enrolls children with median age of 9 years, who seek emergency medical attention with an acute respiratory wheeze at Princess Margaret Hospital in Perth, Australia²⁰. MAVRIC cohort samples were collected during an acute wheezy respiratory episode requiring hospitalization, with follow-up samples collected ≥ 4 weeks later at convalescence. All work on human samples was carried out after informed consent was given in accordance with the declaration of Helsinki.

Clinical parameters and measurements

Atopy was defined as a positive skin prick test (SPT, wheal size ≥ 3 mm) or levels of specific IgE above 0.35 IU/mL as measured using ImmunoCap (Phadia, Sweden) for any of the most common aeroallergens in the Perth region including house dust mite (HDM, *Dermatophagoides pteronyssinus*), cat dander, dog dander, *Alternaria*

alternata, Aspergillus spp., cockroach, Rye grass (*Lolium perenne*), mixed grass and mixed mould. SPT, total IgE and specific IgE were measured in the CAS, Raine and MAVRIC cohorts; SPT and total IgE were measured in the WATCH study. Current asthma was defined as (i) ever having physician diagnosed asthma, (ii) a wheezy episode within the last 12 months and (iii) current use of asthma medication, such as short acting adrenergic β 2 agonists. This definition of asthma excluded children who only wheeze at a very young age.

PBMC collection, storage and thawing

Peripheral blood was collected and processed within 2 hours (CAS and Raine), or 18 hours (WATCH). For the acute cohort (MAVRIC), samples were collected within 72 hours of presentation to Princess Margaret Hospital (Perth, Western Australia) and processed within 18 hours²⁰. PBMCs were obtained using LymphoprepTM (Axis-Shield, Norway) and cryopreserved in liquid nitrogen. Frozen PBMC were thawed as previously²¹. Identical PBMC isolation, cryopreservation, storage and thaw protocols were applied to all cohorts. Viability was assessed using trypan blue exclusion and was above 80%.

Flow cytometry analysis

Approximately 1×10^6 thawed PBMCs were stained as previously described²¹. The same FACS staining panel and protocol was used for all cohorts. Antibodies were titrated and a single batch was used across all samples within a cohort. Additionally, unstained and Fc ϵ R1 isotype stained samples (pooled per run) were included as controls for each staining and acquisition run. Samples were acquired using the LSR-Fortessa platform with FACSDiva software (BD Bioscience, Franklin Lakes, NJ).

Quality control was performed prior to each cytometry run for all cohorts; CS&T beads (BD) to ensure standardised setup and technical performance, and Rainbow calibration particles (BD) to determine alignment and track performance across detection channels and lasers. Following acquisition, data were analysed using FlowJo 10.3 (FlowJo, Ashland, Ore) and FlowSOM in R (v3.2.4)²².

Statistical Analysis

The statistical significance of difference between two groups was estimated using Student's t-test or the Mann-Whitney U test depending on the data distribution; for comparison of three groups the Kruskal-Wallis test followed by Dunn's post test was used; for comparison of paired samples Wilcoxon's Signed rank test was used and for comparison of categorical data between groups, Pearson's χ^2 test was used. Adjustment for IgE levels was calculated using Nominal logistic regression for current asthma within atopic individuals using log(serum IgE levels) and proportion of FcεRIα⁺ subset or log(FcεRIα MFI). Experimental and technical variation was minimised within and between cohorts, however the presence of latent variables between cohorts is a common consideration inherent to cross-cohort studies. To account for this when combining cohorts to compare FcεRIα expression, each cohort was normalised with respect to the 10% of subjects with the highest MFI for each cohort and cell type. The 10% threshold was considered reasonable as it captures the characteristic plateau observed in all cell types for the highest FcεRIα expressing individuals (Fig S3). Additionally, complete randomisation of subjects with respect to asthmatic status within the Raine cohort was not possible due to the experimental design. For analysis of FcεRIα MFI in the Raine cohort, expression was normalised to the run-specific isotype control. This consideration was not required for the CAS,

WATCH and MAVRIC cohorts determined by the consistency of the isotype controls (data not shown). All statistical analysis was performed using JMP 13 (SAS, NSW, Australia) and data was visualised using Prism 7.0a (GraphPad, La Jolla, Calif).

RESULTS

To identify FcεRI expressing cell types in the PBMC fraction, we used a 12-colour flow cytometry panel²¹ (Fig S1) and an unsupervised learning approach (FlowSOM)²² overlaid with manual gating to determine key cell populations (Fig 1A-B) and expression of FcεRIα across these populations (Fig 1C). Using this approach, we identified FcεRIα expression on pDCs, monocytes, cDCs and basophils (Fig 1D), as previously demonstrated²³.

To investigate the association between atopy and asthma with FcεRIα expression, we compared proportion and level of FcεRIα expression in above identified cell populations from (i) non-atopic/non-asthmatic, (ii) atopic/non-asthmatic or (iii) atopic/asthmatic individuals across the CAS, WATCH and Raine cohorts. Compared to non-atopic/non-asthmatic individuals, there was a significantly higher proportion of FcεRIα⁺ pDCs in both atopic groups from the WATCH and CAS cohorts, as well as increased proportion of FcεRIα⁺ monocytes in atopic/non-asthmatic compared to non-atopic in the WATCH or compared to atopic/asthmatic in the Raine cohort (Fig 2A). FcεRIα expression amongst FcεRIα⁺ pDC, cDC, monocyte and basophil populations displayed a progressive gradation of FcεRIα expression from non-atopics to atopic/non-asthmatics, and ultimately to atopic/asthmatics across all cohorts (Fig 2B). As the cohorts consisted of different proportions of individuals with the clinical phenotypes of interest, the power of each individual cohort to detect differences

between some groups differed, and therefore we additionally combined the cohorts as part of the analyses performed as a way of interrogating trends observed within each individual cohort. Once combined, we observed an increased proportion of FcεRIα⁺ pDCs in atopic/asthmatics compared to atopic/non-asthmatics (Fig 2C) and we also observed significantly greater FcεRIα expression levels on FcεRIα⁺ pDCs, cDCs and basophils in atopic individuals (non-asthmatic and asthmatic) compared to non-atopic/non-asthmatics (Fig 2D). Amongst the cohorts the atopic/asthmatics stand out most prominently by this criterion in Raine cohort (Fig 2A-B), and this may be a reflection of relative atopic disease chronicity given that these subjects represent the oldest group tested. We further investigated whether FcεRIα expression was influenced by biological sex (Fig S2A-C) or use of corticosteroid medication within asthmatic individuals (Fig S3A-D) observing that asthmatic individuals on corticosteroids expressed higher levels of FcεRIα compared to individuals on other asthma medication.

Given the well established correlations of FcεRIα expression and total IgE serum levels, which we also observed (Fig S4A-D), we compared total serum IgE between our groups after combining cohorts (Fig 3A). Atopics (non-asthmatic and asthmatic) displayed significantly higher serum IgE levels compared to non-atopic/non-asthmatics. In addition, atopic/asthmatics also displayed higher levels compared to atopic/non-asthmatics. Therefore, to determine if the observed differences in proportion of FcεRIα⁺ pDCs between atopic/non-asthmatics and atopic/asthmatics as well as the increased levels of FcεRIα expression were dependent of total IgE levels, we adjusted for IgE levels using logistic regression on atopic individuals with asthma as the outcome. We observed that both the proportion of FcεRIα⁺ pDCs (Fig 3B) as

well as increased level of expression on FcεRIα⁺ pDCs, cDCs and basophils (Fig 3C) were associated with asthma independently of serum IgE levels, suggesting the operation of a mechanism by which FcεRIα expression is associated with asthma which goes beyond the control of IgE production.

We finally sought to assess the impact of asthma exacerbations on circulating FcεRIα⁺ cell subsets in atopic children. Through the MAVRIC cohort, we recruited subjects presenting at hospital during an acute wheezing illness. For each subject, an acute sample was assessed against a paired convalescent sample obtained once symptoms had resolved several weeks later (Fig 4). Plasmacytoid DC, cDC, and basophil cell numbers were significantly decreased during respiratory exacerbation compared to convalescence (Fig 4A-B), consistent with their likely exit from circulation and migration to sites of inflammation in the airways and their subsequent participation in the local host immunoinflammatory response. No correlation between systemic steroid administration and cell numbers were observed (Fig S5A-D). Together this suggests that these subsets are systemically influenced by wheezing airway disease in atopic individuals.

DISCUSSION

In the present study, we have systematically characterised the expression of FcεRI on cells in the PBMC fraction in non-atopic and atopic children with and without asthma. Our findings are broadly consistent with upregulated expression of FcεRI on both effector (basophil) and regulatory (myeloid, especially pDC/cDC) populations and this upregulation is associated with current atopic asthma. Interestingly, the correlation of FcεRI expression and total IgE levels is not as strong as previously reported²⁴,

potentially due to the young age of the participants. There are also indications that these subsets migrate to the airways during an acute exacerbation²⁵. However given the small sample size and potential interference from administration of systemic steroids, this requires further investigation.

Regarding underlying mechanism(s), enhanced FcεRI expression on basophils equates to reducing the allergen-exposure threshold required for triggering IgE-mediated degranulation and attendant acute inflammation, but the implications of comparable upregulation on myeloid cells are more complex. Circulating cDC are the immediate source of precursors required to renew the airway mucosal DC network which turns over rapidly during infections/exacerbations²⁶, and their migration to this tissue pre-armed with FcεRI would increase the intensity of acute aeroallergen-specific Th2-effector responses and promote expansion of corresponding Th2-memory populations. A similar mechanism applies to circulating monocytes which traffic to airways following allergen challenge²⁷ and further give rise to monocyte derived DC²⁸. However, expression of FcεRI on monocytes does not appear to separate atopic/non-asthmatic and asthmatic individuals in this data set.

Plasmacytoid DCs are recognised as the major producers of type I interferons, the most important mediators of host viral defence²⁹. Consequently, pDCs are a likely candidate to bridge the gap between IgE sensitisation and respiratory viral infection as major asthmatic risk factors and may have a central role in regulating airway inflammation during asthma exacerbations triggered by viruses²⁹. In a subsample from the COAST cohort, an increased proportion of pDCs expressing FcεRI in atopic asthmatic children was observed compared to non-atopic non-asthmatics children¹¹.

However, no significant difference was observed when compared to allergic non-asthmatics, possibly a result of sample size. Notably within the COAST study, allergic asthmatic children demonstrated significantly impaired ability to produce type I interferons following FcεRI cross-linking with IgE in the presence of human rhinovirus, indicating that FcεRI expressing pDCs may have a reduced ability to initiate viral defence¹¹. Omalizumab treatment decreases circulating IgE levels and associated FcεRI cell surface expression. Importantly, omalizumab treatment reduces asthma exacerbation frequency and results in restored type I interferon response to rhinovirus³⁰.

In conclusion, our findings support the previous reports outlining an important role for FcεRI in atopic disease. Our study also suggest that FcεRI expression may be used to identify children with maximal susceptibility to allergic asthma and thus identify kids that may benefit the most from intervention therapies targeting atopy or allergic/Th2 mediated inflammation.

ACKNOWLEDGMENTS

The authors would like to acknowledge all study participants and families involved in the study, their treating physicians and technicians involved in sample processing.

REFERENCES

1. Mohammad HR, Belgrave D, Kopec Harding K, Murray CS, Simpson A, Custovic A. Age, sex and the association between skin test responses and IgE titres with asthma. *Pediatric allergy and immunology : official publication of the European Society of Pediatric Allergy and Immunology*. 2016;27(3):313-319.
2. Holgate ST, Wenzel S, Postma DS, Weiss ST, Renz H, Sly PD. Asthma. *Nature Reviews Disease Primers*. 2015;10.1038/nrdp.2015.25.

3. Dehlink E, Baker AH, Yen E, Nurko S, Fiebiger E. Relationships between levels of serum IgE, cell-bound IgE, and IgE-receptors on peripheral blood cells in a pediatric population. *PLoS One*. 2010;5(8):e12204.
4. Gould HJ, Sutton BJ. IgE in allergy and asthma today. *Nature reviews Immunology*. 2008;8(3):205-217.
5. Lin S, Cicala C, Scharenberg AM, Kinet J-P. The Fc(epsilon)RIbeta Subunit Functions as an Amplifier of Fc(epsilon)RIgamma-Mediated Cell Activation Signals. *Cell*. 1996;85(7):985-995.
6. Maurer D, Ebner C, Reininger B, et al. The high affinity IgE receptor (Fc epsilon RI) mediates IgE-dependent allergen presentation. *The Journal of Immunology*. 1995;154(12):6285-6290.
7. Novak N, Gros E, Bieber T, Allam JP. Human skin and oral mucosal dendritic cells as 'good guys' and 'bad guys' in allergic immune responses. *Clin Exp Immunol*. 2010;10.1111/j.1365-2249.2010.04162.x:28-33.
8. Greer AM, Wu N, Putnam AL, et al. Serum IgE clearance is facilitated by human FcepsilonRI internalization. *The Journal of clinical investigation*. 2014;124(3):1187-1198.
9. Bieber T. The pro- and anti-inflammatory properties of human antigen-presenting cells expressing the high affinity receptor for IgE (FcεRI). *Immunobiology*. 2007;212(6):499-503.
10. Gill MA, Bajwa G, George TA, et al. Counterregulation between the FcepsilonRI pathway and antiviral responses in human plasmacytoid dendritic cells. *J Immunol*. 2010;184(11):5999-6006.
11. Durrani SR, Montville DJ, Pratt AS, et al. Innate immune responses to rhinovirus are reduced by the high-affinity IgE receptor in allergic asthmatic children. *J Allergy Clin Immunol*. 2012;130(2):489-495.
12. Holt PG, Sly PD. Viral infections and atopy in asthma pathogenesis: new rationales for asthma prevention and treatment. *Nat Med*. 2012;18(5):726-735.
13. Vasudev M, Cheung DS, Pincsak H, et al. Expression of high-affinity IgE receptor on human peripheral blood dendritic cells in children. *PLoS One*. 2012;7(2):e32556.
14. Sihra BS, Kon OM, Grant JA, Kay AB. Expression of high-affinity IgE receptors (Fc epsilon RI) on peripheral blood basophils, monocytes, and eosinophils in atopic and nonatopic subjects: Relationship to total serum IgE concentrations. *Journal of Allergy and Clinical Immunology*. 1997;99(5):699-706.
15. Tversky JR, Le TV, Bieneman AP, Chichester KL, Hamilton RG, Schroeder JT. Human blood dendritic cells from allergic subjects have impaired capacity to produce interferon-α via toll-like receptor 9. *Clinical & Experimental Allergy*. 2008;38(5):781-788.
16. Kusel MM, de Klerk NH, Kebadze T, et al. Early-life respiratory viral infections, atopic sensitization, and risk of subsequent development of persistent asthma. *J Allergy Clin Immunol*. 2007;119(5):1105-1110.
17. McClenaghan J, Warrington NM, Jamrozik EF, et al. The PHF11 gene is not associated with asthma or asthma phenotypes in two independent populations. *Thorax*. 2009;64(7):620-625.
18. Newnham PJ, Evans FS, Michael AC, Stanley JF, Landau IL. Effects of frequent ultrasound during pregnancy: a randomised controlled trial. *Lancet*. 1993;342:887-891.

19. Oddy WH, de Klerk NH, Sly PD, Holt PG. The effects of respiratory infections, atopy, and breastfeeding on childhood asthma. *Eur Respir J*. 2002;19(5):899-905.
20. Cox DW, Bizzintino J, Ferrari G, et al. Human rhinovirus species C infection in young children with acute wheeze is associated with increased acute respiratory hospital admissions. *Am J Respir Crit Care Med*. 2013;188(11):1358-1364.
21. Leffler J, Jones AC, Hollams EM, et al. Basophil counts in PBMC populations during childhood acute wheeze/asthma are associated with future exacerbations. *J Allergy Clin Immunol*. 2018;142(5):1639-1641 e1635.
22. Van Gassen S, Callebaut B, Van Helden MJ, et al. FlowSOM: Using self-organizing maps for visualization and interpretation of cytometry data. *Cytometry A*. 2015;87(7):636-645.
23. Berings M, Gevaert P, De Ruyck N, et al. FcεpsilonRI expression and IgE binding by dendritic cells and basophils in allergic rhinitis and upon allergen immunotherapy. *Clin Exp Allergy*. 2018;48(8):970-980.
24. Foster B, Metcalfe DD, Prussin C. Human dendritic cell 1 and dendritic cell 2 subsets express FcεpsilonRI: correlation with serum IgE and allergic asthma. *J Allergy Clin Immunol*. 2003;112(6):1132-1138.
25. Dua B, Watson RM, Gauvreau GM, O'Byrne PM. Myeloid and plasmacytoid dendritic cells in induced sputum after allergen inhalation in subjects with asthma. *Journal of Allergy and Clinical Immunology*. 2010;126(1):133-139.
26. Holt PG, Strickland DH, Wikstrom ME, Jahnsen FL. Regulation of immunological homeostasis in the respiratory tract. *Nat Rev Immunol*. 2008;8(2):142-152.
27. Eguíluz-Gracia I, Malmstrom K, Dheyaaldeen SA, et al. Monocytes accumulate in the airways of children with fatal asthma. *Clinical & Experimental Allergy*. 2018;10.1111/cea.13265.
28. Plantinga M, Guillems M, Vanheerswynghe M, et al. Conventional and monocyte-derived CD11b(+) dendritic cells initiate and maintain T helper 2 cell-mediated immunity to house dust mite allergen. *Immunity*. 2013;38(2):322-335.
29. Lynch JP, Mazzone SB, Rogers MJ, et al. The plasmacytoid dendritic cell: at the cross-roads in asthma. *Eur Respir J*. 2014;43(1):264-275.
30. Teach SJ, Gill MA, Togias A, et al. Preseasonal treatment with either omalizumab or an inhaled corticosteroid boost to prevent fall asthma exacerbations. *J Allergy Clin Immunol*. 2015;136(6):1476-1485.

TABLES

Table 1 – Clinical and demographic features for the respective cohorts.

	CAS	WATCH	Raine	MAVRIC	p [†]
Number of individuals, n (% of total)	119 (32.0%)	174 (46.8%)	79 (21.2%)	19 n/a	n/a

individuals)					
Males, n (%)	71 (59.7%)	100 (57.5%)	44 (55.7%)	12 (63.2%)	0.91
Age, median years (range)	10.1 (10.0-10.7)	10.6 (7.2-13.0)	14.1 (13.5-14.9)	8.8 (6.9-13.2)	<0.001
Total IgE, median IU/mL (range)	166.8 (0-9892.1)	61.5 (1.0-747.0)	378.9 (38.3- 5206.6)	585.3 (123.7- 2342.1)	<0.001
SPT ⁺ , n (%)	74 (62.2%)	132/172 (76.7%) [‡]	65/75 (86.7%) [‡]	11/13 (84.6%) [‡]	<0.001
Atopy (SPT ⁺ or sIgE > 0.35 IU/mL), n (%)	98 (82.4%)	132 (75.9%)	79 (100%)	19 (100%)	<0.001
Current Asthma, n (%)	7 (5.9%)	28 (16.1%)	40 (50.6%)	19 (100%)	<0.001
Additional atopic disease [§] within atopic individuals, n (%)	63 (64.3%)	73/90 (81.1%) [‡]	51 (64.6%)	n/a	0.02
Admitted to hospital with acute wheeze, n (%)	n/a	n/a	n/a	15/17 (88.2%) [‡]	n/a
Hours from last systemic corticosteroid administration (1 mg/kg) to blood collection, median (range)	n/a	n/a	n/a	4.1 (0.4-14.8) [‡] , n = 13	n/a
Respiratory viral infection positive, n (%)	n/a	n/a	n/a	11/14 (78.6%) [‡]	n/a
Total PBMC fraction cell	1.9 (0.8-4.2)	1.1 (0.6-5.1)	1.5 (0.8-4.1)	0.5 (0.3-2.3)	<0.001

count, x 10 ⁶ /mL median (range)					
--	--	--	--	--	--

[†]p-values of statistical difference between cohorts using either Kruskal-Wallis or Pearson's χ^2 test depending on characteristics of the data.

[‡]data missing from some individuals.

[§]additional atopic diseases were allergic rhinitis or eczema within last 12 months.

FIGURE LEGENDS

Figure 1

A Algorithm (FlowSOM) based clustering of cells found within the PBMC fraction followed by manual classification using surface marker expression. **B** Cell numbers per mL of blood of the major cell populations of individuals included in the WATCH study. **C-D** Mapping of FcεRIα expression across the cell populations identified by FlowSOM (**C**) and representative gates of FcεRIα⁺ cell subsets (**D**) Representative figures from one individual are shown in A, C and D. In B, data is displayed as median, 25-75% percentile and range, n=174.

Figure 2

A-D Proportion (A,C) and expression of FcεRIα (B,D) on FcεRIα⁺ cells within each cell type found in PBMC fraction in non-atopic/non-asthmatic, atopic/non-asthmatic or atopic/asthmatic individuals from each cohort (**A-B**) or combined (**C-D**). Significance of difference was calculated using Kruskal-Wallis test followed by Dunn's post test or Mann-Whitney's for comparisons of two groups in the Raine cohort and indicated as; *** p<0.001, ** p<0.01, * p<0.05. Data are displayed as median, 25-75% percentile and range, CAS n_(non-atopic/non-asthmatic)=21, CAS n_{(atopic/non-}

asthmatic)=91 and CAS $n_{(\text{atopic/asthmatic})}=7$. WATCH $n_{(\text{non-atopic/non-asthmatic})}=42$, WATCH
 $n_{(\text{atopic/non-asthmatic})}=104$ and WATCH $n_{(\text{atopic/asthmatic})}=28$. Raine $n_{(\text{atopic/non-asthmatic})}=39$ and
 Raine $n_{(\text{atopic/asthmatic})}=40$. combined $n_{(\text{non-atopic/non-asthmatic})}=63$, combined $n_{(\text{atopic/non-asthmatic})}=235$ and combined $n_{(\text{atopic/asthmatic})}=75$.

Figure 3

A Total IgE serum levels in non-atopic/non-asthmatic, atopic/non-asthmatic or atopic/asthmatic individuals in the combined cohorts. **B-C** Association of $\text{Fc}\epsilon\text{RI}\alpha^+$ cells proportion in PBMC fraction (log odds ratio $\pm 95\%$ CI; **B**) or normalised expression levels of $\text{Fc}\epsilon\text{RI}\alpha$ on $\text{Fc}\epsilon\text{RI}\alpha^+$ subsets (log expression level $\pm 95\%$ CI; **C**) to current asthma within atopics of the combined cohorts, adjusted for $\log(\text{serum IgE})$. Data in **A** are displayed as median, 25-75% percentile and range, combined $n_{(\text{non-atopic/non-asthmatic})}=63$, combined $n_{(\text{atopic/non-asthmatic})}=235$ and combined $n_{(\text{atopic/asthmatic})}=75$. In **B-C** the estimated effect size (log odds ratio) for asthma among atopics $\pm 95\%$ CI per unit increase in % positive cells (**B**) or in $\log(\text{expression level})$ (**C**) is displayed, $n_{(\text{atopic})}=309$.

Figure 4

A-B Concentrations of cells in the PBMC fraction at exacerbation and convalescence (**A**) or fold change of cell concentration in the same individual during exacerbation compared to convalescence (**B**). Significance of difference on paired samples was calculated using Wilcoxon's Signed ranks test and indicated as; *** $p < 0.001$. Data are displayed as individual values, median, 25-75% percentile and range, $n=19$.

Figure 2

pai_13063_f2bw.tiff

Figure 3

pai_13063_f3bw.tiff

Figure 4

D

CD25 - BV421

pDC

Fc γ R1

Figure 2

pai_13063_f2.tiff

Figure 3

pai_13063_f3.tiff

Figure 4

pai_13063_f4.tiff