EFFECTS OF MUSCLE STRENGTH AND ENDURANCE ON BLOOD PRESSURE AND RELATED CARDIO-METABOLIC RISK FACTORS FROM CHILDHOOD TO ADOLESCENCE

Denise L. DEMMER¹, Lawrence J. BEILIN¹, Beth HANDS², Sally BURROWS¹, Kay L. COX¹, Leon STRAKER³, Trevor A. MORI¹

¹*The University of Western Australia (UWA), School of Medicine and Pharmacology, Royal Perth Hospital, Perth, Australia* ²*Notre Dame University, Perth, Australia,* ³*School of Physiotherapy and Exercise Science, Curtin University, Perth, Australia,*

Sources of funding:

Funding for Core Management of the Raine Study was provided by the: University of Western Australia (UWA); Telethon Kids Institute; Raine Medical Research Foundation; UWA Faculty of Medicine, Dentistry and Health Sciences; Women's and Infant's Research Foundation; and Curtin University. This work was supported by grants from the Raine Medical Research Foundation for the 10 year cohort review; National Health and Medical Research Council (NH&MRC) Program Grants (ID 211912; ID 003209) and the Raine Medical Research Foundation for the 14 year cohort review; and NH&MRC Program Grants (ID 35351417, ID 403981) for the 17 year follow up.

Conflict(s) of interest/Disclosure(s) statement: None.

Address all correspondence and requests for reprints to:

Denise Demmer, School of Medicine and Pharmacology, Royal Perth Hospital Unit,
University of Western Australia, GPO Box X2213, Perth, Western Australia 6847, Australia.
Telephone: +61 8 92240343 Fax: +61 8 92240243 E-mail: denise.demmer@uwa.edu.au

ABSTRACT

Background: Muscle function as assessed by strength and endurance is an important aspect of physical fitness. Low levels of muscle strength have been associated; insulin resistance and dyslipidaemia in adults but the effects on blood pressure are less clear. Moreover, there is a lack of data examining the evolution of relationships between muscle function and cardio-metabolic risk factors in childhood and adolescence. This study aimed to examine the evolution of relationships between measures of muscle strength and endurance with individual cardio-metabolic risk factors from childhood to late adolescence in a prospective population based cohort.

Methods: Participants from the Western Australian Pregnancy Cohort (Raine) Study at ages 10, 14 and 17 were analysed, using longitudinal linear mixed model analyses.

Results: Handgrip strength after adjusting for the confounding effects of BMI was positively associated with systolic BP, but not diastolic BP. The association between handgrip strength and systolic BP was stronger in males than females at all time points (coefficient (females): 0.18, $p<0.001$; sex*handgrip strength coefficient: 0.09, $p=0.002$). The association was strongest at 10 years and significantly attenuated over time (year*handgrip coefficient from 10 to 14 years: -0.11, $p=0.003$; year*handgrip coefficient from 10 to 17 years: -0.19, $p<0.001$). After the inclusion of BMI as a confounder, handgrip strength was significantly negatively associated with HOMA-IR and hs-CRP over time in both sexes. Back muscle endurance was positively associated with systolic BP, but not diastolic BP, after adjustment for the confounding effects of BMI (coefficient: 0.01, $p=0.002$). There were small, albeit

significant inverse associations between back muscle endurance and log HOMA-IR and log hs-CRP.

Conclusion: The positive association between handgrip strength and back muscle endurance with systolic BP throughout childhood and adolescence contrasts with beneficial effects on other related traditional cardio-metabolic risk factors. Mechanisms underlying these paradoxical effects on systolic BP warrant further investigation.

INTRODUCTION

Moderate to high levels of cardiorespiratory fitness have been associated with lower mortality rates (1) and lower levels of hypertension, dyslipidaemia and insulin resistance (2, 3). Muscle strength and endurance are key contributors to measures of physical fitness. Muscle strength is defined as the ability to generate a maximal force during a single contraction, whereas muscle endurance is the ability to maintain a repeated exertion of force over an extended period of time (4). The handgrip test acts as a proxy for general muscle strength and is associated with total body strength measures in adults (5) and children (6).

In middle-aged adults, low muscle strength has been associated with high morbidity and mortality rates (7). At least part of this effect is suggested to be mediated by beneficial influences on standard cardio-metabolic risk factors. For example, increased muscular mass due to resistance training has been shown to be protective against the onset of insulin resistance (8). Furthermore, large epidemiological studies have shown high levels of handgrip strength are associated with lower levels of undiagnosed hypertension (9) and to be a moderately strong predictor of incident cardiovascular disease in adults (10). However, there are limited data in children and adolescents.

Adolescence is a critical period of substantial physiological, psychological and lifestyle changes which can impact on future adult behaviours and health outcomes (11). Muscle strength and endurance increase as children age, due to growth in muscle mass and muscle fibre size (12). In addition, BP has been shown to linearly increase throughout adolescence (13). With a view to understanding the ontogeny of effects of muscle strength and endurance on adolescent risk factors for cardio-metabolic disease, we have examined two modes of

muscle function, specifically handgrip strength and back muscle endurance. We then analysed their relationships with a range of individual cardio-metabolic risk factors in participants from the population based Western Australian Pregnancy Cohort (Raine) Study. We hypothesised that higher levels of handgrip strength and back muscle endurance would be associated with lower levels of cardio-metabolic risk factors.

METHODS

Participants

The Raine Study is a prospective population study that recruited 2900 pregnant women between 16 to 20 weeks' gestation from King Edward Memorial Hospital and closely located practices, in Perth, Western Australia, Australia. The mothers gave birth to 2868 live infants. The offspring have been followed at approximately three-year intervals with a comprehensive range of anthropometry, BP, lifestyle, psychosocial and biochemical measures recorded. Details on methods of the Raine Study have previously been reported (14). Ethics approvals for the 10, 14 and 17 year assessments were obtained from the Human Research Ethics Committees at the University of Western Australia, Curtin University and Princess Margaret Hospital. Written informed consent was obtained from the participant's parent and the participant at 17 years age. Handgrip strength measures, BMI and BP recordings were recorded at 10, 14 and 17 years. Back muscle endurance, cardiorespiratory fitness, puberty and biochemistry measures were recorded at 14 and 17 years.

Muscle strength

Muscle strength was assessed using a handgrip dynamometer (Dynamometer, Therapeutic Instruments Clifton N.J.) following the protocol developed by McCarron (15). Participants

were asked to extend and hold the hand dynamometer at shoulder length and pull hard on the device to record grip force (kg). Measurements were assessed with the right hand and the left hand with the process being repeated. The best score of the two trials for each hand was recorded. These were combined to create a total handgrip strength score (kg).

Muscle Endurance

The sustained back muscle endurance test was employed to evaluate dynamic back muscle endurance by testing the postural trunk muscles using a procedure described by Biering-Sorensen (16). Participants were instructed to lie in the prone position with their lower body strapped to a bed and the upper body extended beyond the bed. Subjects were required to maintain the horizontal position isometrically for as long as possible. The endurance time was recorded (seconds).

Cardiorespiratory fitness

Cardiorespiratory fitness was estimated from heart rate recordings during sub-maximal cycle ergometry using the PWC₁₇₀ protocol (Monark cycle ergometer). Heart rate at three different workloads (watts) was used to extrapolate the load required to reach a heart rate of 170 beats per minute. PWC₁₇₀ is the maximal steady state power attained for a heart rate of 170 beats per minute on a cycle ergometer (17).

Anthropometry

Height was measured using a mounted stadiometer (to the nearest 0.1 cm) and weight was measured (to the nearest 100g) with participants dressed in light clothes. BMI was calculated as weight (kg) divided by the square of height (m²).

Puberty

Puberty was assessed using the Tanner Stages of physical development. This scale asks participants to rate their development based on a number of physical sexual characteristics, such as development of breasts, genitalia and pubic hair. The five stages range from 1 (prepubertal) to 5 (fully developed) (18). At 14 years of age, misreporting occurred when approximately 330 individuals were shown the wrong developmental stage chart. To account for this, these individuals were omitted from the analyses when using puberty as a confounder.

Biochemistry

Venous blood samples were taken after an overnight fast and analysed in the PathWest Laboratory at Royal Perth Hospital for serum glucose, insulin, total cholesterol, HDL-C, triglycerides and hs-CRP (19). LDL-C was calculated using the Friedewald formula (20). HOMA-IR was calculated using the formula: fasting insulin ($\mu\text{U/ml}$) x fasting glucose (mmol/L) / 22.5 (21).

Blood pressure

Systolic and diastolic BP were measured by trained personnel using an oscillometric sphygmomanometer (DINAMAP vital signs monitor 8100, DINAMAP XL vital signs monitor or DINAMAP ProCare 100; GE Healthcare, Soma Technology, Bloomfield, Connecticut, USA) after resting for 5 minutes and using the appropriate cuff size on the right arm. Three cuff sizes were available. At 10 years, two BP readings were obtained every 2 minutes in the seated position and the average calculated. At 14 years, six BP readings were obtained every 2 minutes within a 10-minute period in the seated position, whereas at 17

years, the same protocol was employed in the supine position. The average BP was calculated using the last five readings at 14 and 17 years. BP was recorded on a separate day to handgrip strength and back muscle endurance assessments.

Statistical analysis

Descriptive data were summarised by year and sex using means, geometric means and their 95% confidence intervals. There is no consensus in the literature regarding the need to adjust handgrip strength for body size or how this might be achieved. To explore this issue, handgrip strength was calculated at each year by regressing handgrip strength on height or weight respectively, we then added the residuals from these analyses to the mean handgrip strength, to obtain a height adjusted and weight adjusted handgrip strength. Potential non-linearity was accommodated by using fractional polynomials. Hierarchical linear mixed models of a selection of cardio-metabolic outcomes (systolic BP, diastolic BP, log HOMA-IR, log triglycerides, HDL-C, hs-CRP) were then used to compare the performance of raw handgrip strength, height adjusted handgrip strength and weight adjusted handgrip strength. A simple model was adopted for this purpose, containing time, sex and subsequently an adjustment for adiposity, using BMI. Models were assessed using Akaike Information Criterion (AIC) where the lowest value indicates the better model.

Longitudinal patterns of raw handgrip strength and back muscle endurance over time as well as individuals cardio-metabolic risk factors were investigated using hierarchical linear mixed models, to account for nesting of time within an individual and siblings within a family. In these models, maximum likelihood estimation was utilised to retain those with outcome data from at least one-time point in the analysis, resulting in unbiased estimates when missing data

is missing at random. Random effects Tobit regression (for censored data) was utilised for hs-CRP due to the lower boundary of the test (but does not make any adjustment for siblings) (22). HOMA-IR, triglycerides and hs-CRP were not normally distributed and hence were log transformed. Initial associations between cardio-metabolic risk factors and handgrip strength and back muscle endurance were investigated after adjusting for sex and year. A further adjustment for adiposity was investigated in a subsequent model using BMI. Interactions between handgrip strength or back muscle endurance with time or sex were tested to determine whether the relationship varied over time or sex. In addition, a time and sex interaction was included to determine whether an outcome varied over time and sex. Interactions were excluded from the model if $p \geq 0.05$. Cardiorespiratory fitness was included as a confounder in both handgrip strength and back muscle endurance models with either systolic BP or diastolic BP as the outcome. Data were analysed using STATA (StataCorp, 2011. *Stata Statistical Software: release 13*. College Station, Texas, USA). All reported P values are 2-tailed and significance was set at $\alpha=0.05$.

RESULTS

Participant characteristics

At 10, 14 and 17 years, 1632, 1560 and 1234 participants, respectively had complete data for handgrip strength and systolic BP (Figure 1). Handgrip strength and biochemistry at 14 and 17 years was available for 1346 and 1048 participants, respectively. At 14 and 17 years, 1554 and 1158 participants, respectively, had complete data for back muscle endurance and systolic BP, whereas 1341 and 983 participants, respectively, had complete data for back muscle endurance and biochemistry.

Figure 1. Flow diagram of Raine Study participants attending the 10, 14 and 17 year follow-up with complete handgrip strength and back muscle endurance data

* Multiple assessments on n= 1522

Multiple assessments on n= 1025

+ Multiple assessments on n= 888

^ Multiple assessments on n= 833

Interpretation of handgrip strength: Comparison of unadjusted and adjusted measures

For the determination of the ‘better’ handgrip strength measure, identical models of raw handgrip strength, height adjusted handgrip strength and weight adjusted handgrip strength were compared (Supplementary Table 1a, 1b and 1c.). Raw handgrip strength in unadjusted and BMI confounder adjusted models exhibited the lowest AIC compared to height and weight adjusted handgrip strength. Therefore, raw handgrip strength was deemed a better fit for the outcome of systolic BP and hence was employed as the preferred handgrip strength measure in the final analyses. The same pattern was observed for other cardio-metabolic risk factors investigated (only systolic BP data shown as supplementary material).

Longitudinal linear mixed models of handgrip strength, back muscle endurance, blood pressure and biochemical risk factors

Handgrip strength increased between 10 to 17 years in both sexes ($p<0.001$), but to a greater extent in males ($p<0.001$) (Table 1). Back muscle endurance was higher in females than in males at 14 years ($p=0.021$) but by 17 years there was no difference between the sexes ($p=0.200$) (Table 1). Back muscle endurance increased between 14 and 17 years in females ($p<0.001$) and also in males but to a greater extent ($p=0.001$) (Table 1). Handgrip strength and back muscle endurance were weakly correlated ($r=0.07$, $p=0.008$ at 14 years; $r=0.08$, $p=0.004$ at 17 years). Systolic BP increased between 10, 14 and 17 years in both sexes but to a greater extent in males ($p<0.001$) (Table 1). Diastolic BP increased between years 10 and 14 in both sexes but to a greater extent in females ($p<0.001$). There was no significant change in diastolic BP between 14 and 17 years for either sex. Log HOMA-IR decreased between 14 and 17 years in both sexes to a similar extent ($p<0.001$) (Table 1). Cholesterol decreased in males ($p<0.001$) with no change in females over time ($p=0.281$). Log triglycerides increased

in males ($p<0.001$) with no change in females over time ($p=0.547$). LDL-C increased in females ($p=0.004$) with no change in males over time ($p=0.818$). HDL-C decreased in both sexes but to a greater extent in males ($p<0.001$). Log hs-CRP increased from 14 to 17 years in females ($p<0.001$), but decreased in males ($p<0.001$).

Longitudinal linear mixed models examining associations between handgrip strength and cardio-metabolic risk factors over time

Handgrip strength was positively associated with systolic BP at all time points in both sexes. The association was strongest at 10 years and attenuated over time (Table 2, Supplementary Table 2 and Supplementary Figure 1).

After adjustment for the confounding effects of BMI, similar associations were detected (Table 2, Supplementary Table 2 and Figure 2). The association between handgrip strength and systolic BP was stronger in males than females at all time points (sex*handgrip strength coefficient: 0.09, $p=0.002$). In both sexes, handgrip strength was positively associated with systolic BP at 10 years of age and attenuated over time, (year*handgrip for 10 to 14 years coefficient: -0.11, $p=0.003$; year*handgrip for 10 to 17 years coefficient: -0.19, $p<0.001$) resulting in a non-significant association at 17 years of age in females. The change in systolic BP resulting from a 1 SD change (noted in brackets) in females was 1.07 mm Hg (5.94 kg) and 0.72 mm Hg (9.02 kg) for years 10 and 14 respectively. The corresponding changes in males were 1.78 mm Hg (6.37 kg), 2.50 mm Hg (14.70 kg) and 1.43 mm Hg (15.86 kg) for years 10, 14 and 17 respectively.

To investigate whether these associations may be attributable to cardiorespiratory fitness (only available at 14 and 17 years); PWC₁₇₀ was included as a further confounder with BMI

(Supplementary Table 2). The same pattern was observed after the additional adjustment for PWC₁₇₀ between 14 and 17 years when BMI was the sole confounder in adjusted models.

There were no associations between handgrip strength and diastolic BP at any time point in the unadjusted, BMI-adjusted (Table 2) and the PWC₁₇₀ and BMI-adjusted models (data not shown). Handgrip strength was positively associated with HOMA-IR in both sexes at all time points (coefficient (log HOMA-IR): 0.003, $p=0.013$) (Table 3 and Supplementary Table 3). After the inclusion of BMI as a confounder in the model, handgrip strength was negatively associated with HOMA-IR in both sexes at all time points, (coefficient (log HOMA-IR): -0.003, $p<0.001$). A 1 SD change in handgrip strength at 14 years (13.40 kg) and 17 years (20.58 kg) equated to a decrease in HOMA-IR of 3.9% and 6 %, respectively.

Handgrip strength was positively associated with triglycerides at all time point in both sexes in the unadjusted model (coefficient (log triglycerides): 0.002, $p=0.006$) (Table 3). After the inclusion of BMI as a confounder, the aforementioned association was no longer significant (coefficient (log triglycerides): -0.0007, $p=0.302$). Handgrip strength was negatively associated with HDL-C and this association attenuated from 14 to 17 years in both sexes, (coefficient at 14 years: -0.004, $p<0.001$; year*handgrip coefficient: 0.003, $p<0.001$) (Table 4 and Supplementary Table 4). After, the inclusion of BMI as a confounder, handgrip strength remained negatively associated with HDL-C at 14 years (coefficient: -0.003, $P<0.001$) but the association was no longer significant at 17 years (year*handgrip coefficient: 0.003, $p<0.001$). A 1 SD change in handgrip strength at 14 years (13.40 kg) equated to a 0.04 mmol decrease in HDL-C. There was a positive association between handgrip strength and hs-CRP (coefficient (log hs-CRP): 0.005, $p=0.013$) (Table 3 and Supplementary Table 5). After the inclusion of BMI as a confounder, there was a small negative, albeit significant, association

between handgrip strength and hs-CRP (coefficient (log hs-CRP): -0.006, $p=0.007$). These associations did not vary with sex or time. A 1 SD change in handgrip strength at 14 years (13.40 kg) and 17 years (20.58 kg) equated to a decrease in hs-CRP of 7.7% and 11.3%, respectively.

Longitudinal linear mixed models examining associations between back muscle endurance and cardio-metabolic risk factors over time

There were no interactions for back muscle endurance and time or back muscle endurance and sex for any of the cardio-metabolic risk factors investigated. The associations reported therefore apply to both time points in both sexes. Back muscle endurance was not associated with systolic BP in unadjusted analysis (coefficient: -0.001, $p=0.723$) (Table 5, Supplementary Table 6 and Supplementary Figure 2). After the inclusion of BMI as a confounder in the model, back muscle endurance was positively associated with systolic BP (coefficient: 0.01, $p=0.002$) (Table 5, Supplementary Table 6 and Figure 3). A 1 SD (14 years: 62.83 seconds; 17 years: 56.67 seconds) increase in back muscle endurance equated to a 0.06 mm Hg increase in systolic BP. After the inclusion of cardiorespiratory fitness as a further confounder, these associations were not altered (Supplementary Table 6). Back muscle endurance was not associated with diastolic BP in unadjusted, BMI adjusted (Table 5) and the PWC₁₇₀ and BMI-adjusted models (data not shown).

Back muscle endurance was inversely associated with HOMA-IR (coefficient (log HOMA-IR): -0.002, $p<0.001$) (Table 5 and Supplementary Table 7). With BMI as a confounder in the model, back muscle endurance remained negatively associated with HOMA-IR (coefficient (log HOMA-IR): -0.001, $P<0.001$). A 1 SD (14 years: 62.83 seconds; 17 years: 56.67

seconds) increase in back muscle endurance equated to a 6% and 5.5% decrease in HOMA-IR at 14 and 17 years, respectively. Back muscle endurance was inversely associated with triglycerides (coefficient (log triglycerides): -0.0009, $p<0.001$) and remained significant after the inclusion of BMI as a confounder (coefficient (log triglycerides): -0.0004, $p=0.007$) (Table 5 and Supplementary Table 8). A 1 SD (14 years: 62.83 seconds; 17 years: 56.67 seconds) increase in back muscle endurance equated to a 2.5% and 2.2% decrease in triglycerides at 14 and 17 years, respectively. Back muscle endurance was positively associated with HDL-C (coefficient: 0.0005, $p<0.001$), but this was no longer significant after the inclusion of BMI as a confounder (coefficient: 0.0001, $p=0.209$) (Table 5). Back muscle endurance was negatively associated with hs-CRP (coefficient (log hs-CRP): -0.003, $p<0.001$) and the relationship remained significant after inclusion of BMI in the model (coefficient (log hs-CRP): -0.001, $p=0.018$) (Table 5 and Supplementary Table 9). A 1 SD (14 years: 62.83 seconds; 17 years: 56.67 seconds) increase in back muscle endurance equated to a 6% and 5.5% decrease in hs-CRP at 14 and 17 years, respectively.

Table 1. Descriptive characteristics of participants from the Raine Study by age and sex

	Males			Females		
	10 years n=849	14 years n=815	17 years n=628	10 years n=783	14 years n=774	17 years n=616
Height (cm)	143.69 (143.25, 144.14)	166.27 (165.65, 166.88)	178.3 (177.74, 178.88)	143.64 (143.20, 144.09)	162.02 (161.58, 162.46)	165.90 (165.39, 166.41)
Weight (kg)	38.57 (38.00, 39.14)	58.62 (57.66, 59.59)	72.29 (71.13, 73.41)	38.71 (38.13, 39.31)	56.67 (55.82, 57.52)	63.66 (62.61, 64.71)
Body mass index (kg/m ²)	20.97 (19.86, 21.18)	21.06 (20.78, 21.35)	22.69 (22.36, 23.02)		21.53 (21.24, 21.83)	23.12 (22.76, 23.48)
Waist circumference (cm)		76.31 (75.52, 77.09)	80.53 (79.67, 81.04)		74.60 (73.88, 75.31)	77.55 (76.63, 78.47)
Handgrip strength (kg)	32.02 (31.60, 32.45)	57.00 (55.98, 58.02)	81.18 (79.93, 82.43)	29.01 (28.59, 29.43)	46.29 (45.64, 46.93)	49.74 (48.95, 50.53)
Back muscle endurance (sec)		78.07 (73.86, 82.28)	102.82 (98.55, 107.10)		85.82 (81.26, 90.39)	98.99 (94.03, 103.95)
Cardiorespiratory fitness PWC ₁₇₀ (watts)		124.35 (122.14, 126.56)	154.39 (151.08, 157.69)		97.20 (95.79, 98.63)	99.62 (97.60, 101.34)
Systolic BP (mm Hg)	106.9 (106.2, 107.5)	113.84 (113.1, 114.6)	117.82 (117.07, 118.57)	106.22 (105.54, 106.89)	108.80 (108.15, 109.46)	108.66 (107.94, 109.38)
Diastolic BP (mm Hg)	56.7 (56.2, 57.1)	58.6 (58.0, 59.0)	58.2 (57.7, 58.7)	56.8 (56.3, 57.2)	59.3 (58.8, 59.7)	59.5 (58.9, 60.0)
Glucose (mmol)		4.72 (4.64, 4.92)	4.65 (4.82, 4.92)		4.88 (4.68, 4.76)	4.87 (4.61, 4.70)
Insulin (mU/L)		12.29 (11.29, 13.30)	8.96 (8.36, 9.57)		12.71 (12.06, 13.36)	9.91 (8.93, 10.89)
HOMA-IR*		0.65 (0.61, 0.70)	0.50 (0.46, 0.55)		0.72 (0.68, 0.77)	0.52 (0.47, 0.56)

Triglycerides (mmol)*	0.22 (0.20, 0.26)	0.27 (0.24, 0.30)	0.19 (0.17, 0.22)	0.21 (0.18, 0.24)
Total cholesterol (mmol)	4.06 (4.01, 4.11)	3.96 (3.90, 4.01)	4.28 (4.23, 4.34)	3.96 (4.24, 4.36)
HDL-C (mmol)	1.35 (1.33, 1.37)	1.20 (1.18, 1.22)	1.43 (1.41, 1.45)	1.38 (1.36, 1.41)
LDL-C (mmol)	2.25 (2.21, 2.30)	2.25 (2.20, 2.31)	2.38 (2.36, 2.43)	2.44 (2.38, 2.49)
hs-CRP (mg/L)*	0.61 (0.52, 0.71)	0.65 (0.55, 0.76)	0.54 (0.42, 0.68)	0.74 (0.64, 0.84)

Descriptive characteristics are presented as means, geometric means and 95% CI. CI, confidence interval; PWC₁₇₀, physical work capacity protocol; BP, blood pressure; HOMA-IR, homeostatic model assessment of insulin resistance; HDL-C, high density lipoprotein-cholesterol; LDL-C, low density lipoprotein-cholesterol; hs-CRP, high-sensitivity C-reactive protein. BP was measured in the seated position at ages 10 and 14 years, and in the supine position at 17 years

*Variables were log transformed

Table 2. Handgrip strength associations with blood pressure generated from a longitudinal linear mixed model

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Systolic BP				
Females				
10 years	0.29 (0.21, 0.37)	<0.001	0.18 (0.11, 0.26)	<0.001
14 years	0.15 (0.09, 0.20)	<0.001	0.08 (0.02, 0.13)	0.007
17 years	0.07 (0.01, 0.12)	0.016	-0.001 (-0.06, 0.05)	0.960
Males				
10 years	0.35 (0.28, 0.42)	<0.001	0.28 (0.20, 0.35)	<0.001
14 years	0.21 (0.17, 0.25)	<0.001	0.17 (0.13, 0.20)	<0.001
17 years	0.13 (0.09, 0.17)	<0.001	0.09 (0.05, 0.13)	<0.001
Diastolic BP				
All years	-0.006 (-0.02, 0.01)	0.558	-0.13 (-0.03, 0.01)	0.196

N= 4425 observations from 1916 participants

Coefficients reported were generated from longitudinal linear mixed models (shown in Supplementary tables). CI, confidence interval; BP, blood pressure

Figure 2. Handgrip strength and systolic BP at 10, 14 and 17 years after adjustment for BMI generated from longitudinal linear mixed models (mean and 95% CI's represented)

Figure 3. Back muscle endurance and systolic BP at 14 and 17 years after adjustment for BMI generated from longitudinal linear mixed models (mean and 95% CI's represented)

Table 3. Handgrip strength associations with HOMA-IR, triglycerides and hs-CRP generated from a longitudinal linear mixed model or random effects tobit model

	Unadjusted model	BMI adjusted model
Log HOMA-IR		
Coefficient	0.003	-0.003
(95% CI)	(0.0006, 0.005)	(-0.005, -0.002)
<i>P</i> value	0.013	0.001
Log Triglycerides		
Coefficient	0.002	-0.0007
(95% CI)	(0.0006, 0.003)	(-0.002, 0.0006)
<i>P</i> value	0.006	0.302
Log hs-CRP*		
Coefficient	0.005	-0.006
(95% CI)	(0.001, 0.01)	(-0.01, -0.002)
<i>P</i> value	0.013	0.007

N= 4425 observations from 1916 participants

*N= 1967 observations from 1363 participants as participants with values >10 were excluded

Coefficients reported were generated from longitudinal linear mixed models or random effects tobit model (shown in Supplementary tables). CI, confidence interval; HOMA-IR, homeostatic model assessment insulin resistance; hs-CRP, high-sensitivity C-reactive protein

Table 4. Handgrip strength associations with HDL-C generated from a longitudinal linear mixed model

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
14 years	-0.004 (-0.006, -0.003)	<0.001	-0.003 (-0.004, -0.002)	<0.001
17 years	-0.002 (-0.003, -0.0007)	0.002	-0.0002 (-0.001, 0.001)	0.796

N= 2396 observations from 1506 participants

Coefficients reported were generated from longitudinal linear mixed models (shown in Supplementary tables). CI, confidence intervals; HDL-C, high density lipoprotein-cholesterol

Table 5. Longitudinal linear models of back endurance with cardio-metabolic outcomes generated from a longitudinal linear mixed model or random effects tobit model

	Unadjusted model	BMI adjusted model
Systolic BP[^]		
Coefficient	-0.001	0.01
(95% CI)	(-0.007, 0.005)	(0.004, 0.02)
<i>P</i> value	0.723	0.002
Diastolic BP[^]		
Coefficient	0.002	0.002
(95% CI)	(-0.002, 0.006)	(-0.002, 0.007)
<i>P</i> value	0.384	0.285
Log HOMA-IR		
Coefficient	-0.002	-0.001
(95% CI)	(-0.002, -0.001)	(-0.001, -0.0006)
<i>P</i> value	<0.001	<0.001
Log Triglycerides		
Coefficient	-0.0009	-0.0004
(95% CI)	(-0.001, -0.0006)	(-0.0007, -0.0001)
<i>P</i> value	<0.001	0.007
HDL-C		
Coefficient	0.0005	0.0001
(95% CI)	(0.0003, 0.0007)	(-0.00007, 0.0003)
<i>P</i> value	<0.001	0.209
Log hs-CRP*		
Coefficient	-0.003	-0.001
(95% CI)	(-0.004, -0.002)	(-0.002, -0.0002)
<i>P</i> value	<0.001	0.018

N= 2325 observations from 1491 participants

[^]N= 2712 observations from 1687 participants

*N= 1895 observations from 1337 participants as participants with values >10 were excluded

Coefficients reported were generated from longitudinal linear mixed models or random effects tobit model (shown in Supplementary tables).

DISCUSSION

To our knowledge, this the first study examining the relationship between measures of muscle performance i.e. muscle strength and endurance, with BP and other cardio-metabolic risk factors from childhood to late adolescence. The study has shown unexpectedly, handgrip strength was positively associated with systolic BP and attenuated over time. The association between handgrip strength and systolic BP was stronger in males than females, with a non-significant association at 17 years of age in females. More predictably, handgrip strength and back muscle endurance were inversely associated with HOMA-IR and hs-CRP after the inclusion of BMI as a confounder. Additionally, back muscle endurance was inversely associated with plasma triglycerides.

Previous paediatric studies investigating the effects of muscle strength on cardiovascular risk factors have examined cardio-metabolic risk factors as components of a clustered metabolic risk score. These studies have shown an inverse relationship between muscle strength and clustered metabolic risk (23-25). Artero et al. (23) used a muscular fitness score computed using the standardised values of handgrip strength and standing long jump (separately for males and females) and showed a negative association with a clustered metabolic risk score which included systolic BP, HOMA-IR, total cholesterol/HDL-cholesterol and triglycerides in 12 to 17 year old participants of the HELENA study. In addition, the ACFIES study observed that poorer handgrip strength were associated with a worse cardio-metabolic risk in Columbian schoolchildren, aged 8-14 years (26). However, the aforementioned studies did not examine BP and related cardio-metabolic risk factors individually. More recently, Zaquout et al. (27) examined the longitudinal associations between individual components of physical fitness (cardiorespiratory fitness, handgrip strength and lower limb strength and speed/agility)

and the combined sum of these individual physical fitness components (using z -scores) with metabolic risk markers, in 6-11 year old European participants in the IDEFICS study. No longitudinal associations were observed between handgrip strength or the sum of physical fitness with clustered metabolic risk (sum of age and sex specific z -scores of waist circumference, systolic BP, diastolic BP, triglycerides, HDL-C and HOMA-IR) or with systolic BP, HOMA-IR and blood lipids. Lower limb strength (measured by standing long jump) showed longitudinal and inverse associations with clustered metabolic risk, blood lipids and HOMA-IR but there were no associations with systolic BP. No studies to date have examined BP or related cardio-metabolic risk factors individually or longitudinally from childhood through to adolescence where significant maturational changes occur. Additionally, no studies have examined the association between isometric endurance of the trunk extensor muscles and the individual risk factors.

Adult population studies present conflicting results with muscle fitness and BP. Sayer et al. (28) have shown a lower handgrip strength was associated with higher systolic BP in a population based sample of 2677 men and women aged 59 – 73 years from the Hertfordshire study. In that study, a 1 SD decrease in handgrip strength associated with high BP (odds ratio 1.13 $p=0.004$). Vaara et al. (29) found that muscle endurance was inversely associated with both systolic and diastolic BP in young men ($25\pm$ years). The younger age of our participants and our use of dynamic assessment of muscle endurance may explain the observational differences seen in our study. In contrast, Viitasalo et al. (30) found positive correlations between muscle strength of the trunk extensor muscles with systolic BP in young healthy men (mean age 19.7 years). Takaema et al. (31) reported higher levels of handgrip strength

associated with higher resting BP in a large population based sample of Japanese men and women (aged 66 years). In contrast to our study of muscle strength and muscle endurance per se, a study of self-reported aerobic exercise in young adults (<30 years), showed an association with lower diastolic BP and higher pulse pressure (32). Aerobic physical activity was self-reported via questionnaire (International Physical Activity Questionnaire, 2002) and individuals who reported regular resistance training were excluded. There were no differences in central or brachial systolic BP between exercising and non-active individuals. This study contrasts with our results, as we found a positive association between handgrip strength and back muscle endurance with systolic BP, even after adjustment for cardiorespiratory fitness. Additionally, we observed no associations between muscle strength or muscle endurance with diastolic BP in our young population. Discrepancies between the two studies are likely due to the different measurement methods, types of physical fitness assessed, and different ages of the populations.

Population data are in contrast to resistance training studies which have consistently shown improvements in handgrip strength are associated with reductions in systolic BP (33-35). Owen et al. (36) suggested handgrip exercise might increase forearm blood flow and/or improve local flow-mediated dilation (FMD). Enhanced brachial FMD response as well as endothelium dependent vasodilatation were observed after isometric handgrip training in older men, elderly hypertensives and chronic heart failure subjects (37). Similarly, forearm blood flow was increased with regular handgrip exercise training in young men and in middle-aged men (38). However, it is important to note various factors, such as exercise intensity, different populations and exercise modality can have differing influences on BP.

The lack of effect of muscle strength and muscle endurance on diastolic BP in our study is of interest given that systolic BP and more predominantly diastolic BP in young adults are predictive of subsequent cardiovascular events and mortality (39). Our findings of a positive relationship between handgrip strength and back muscle endurance with systolic BP across periods of the survey may be related to the fact that both muscle strength and BP increase in magnitude during adolescence. Nevertheless, we saw similar associations cross-sectionally at each time point in subjects that were predominantly pre-pubertal at 10 years to post-puberty at 17 years. Thus it is possible that changes in androgen levels related to puberty influenced muscle mass and strength (40, 41) and thereby the positive association with systolic BP (42). Increases in testosterone could increase muscle mass and may have an additive effect on BP, although the associations in our study were seen in females as well as males making this explanation unlikely. This critical period of growth in the Raine cohort is associated with other hormonal changes such as growth hormone, which might also contribute to parallel changes in muscle strength (12) and left ventricular mass and contractility (43, 44) and hence BP. The association between handgrip strength and back muscle endurance with systolic BP in our study could be explained by an increase in arterial stiffness and/or vascular hypertrophy with a higher degree of muscle fitness in childhood and adolescents.

In accordance with our findings with HOMA-IR and triglycerides, adult studies have consistently shown inverse associations between different measures of muscle strength and measures of insulin resistance and lipids (45, 46). Vaara et al. (29) showed that muscular endurance (sit-ups, push-ups and repeated squats) was inversely associated with blood triglycerides and glucose in young men. Increased muscle mass is the most likely explanation for the effects on glucose homeostasis and lipids which appear to persist through childhood

and adolescence into adult life. Skeletal muscle is one of the primary tissue sites for glucose and lipid metabolism and is considered a determinant of resting metabolic rate. Thus changes in muscle mass may influence multiple cardio-metabolic risk factors (47).

The results of our study are likely to be generalisable as Raine study participants were derived from a cohort broadly representative of the general population and were not a selected group of athletes. Strengths of our study include serial measures of detailed phenotypes in a large cohort of population-based children and adolescents, allowing the study of the evolution of associations between different aspects of muscle function and a panel of cardio-metabolic risk factors. Limitations include the use of BMI in the analyses which does not distinguish between fat and lean mass. In addition, not all measures were repeated at every time point. BP recordings varied across the three time points in terms of posture and number of measurements obtained. This could have influenced the associations observed. Relationships with home BP or 24-hour ambulatory monitoring would be of further interest. In conclusion, we have demonstrated that paradoxically handgrip strength and back muscle endurance were positively associated with higher systolic BP in children and adolescents despite inverse associations with HOMA-IR, lipids and hs-CRP. These BP findings warrant replication in other populations and further evaluation to the possible underlying mechanisms. In view of the fact grip strength increases from childhood through to 20-30 years of age (48, 49), it will be of particular interest to ascertain whether the association between handgrip strength and back muscle endurance, with systolic BP, are maintained or reversed in later adulthood. Equally important will be to determine the optimal period to gain a substantial benefit from resistance strength training for the early prevention of type II diabetes, dyslipidaemia and hypertension.

ACKNOWLEDGEMENTS

The authors thank the Raine Study participants and their families, the Raine Study Team for cohort co-ordination and data collection and the Telethon Institute for Child Health Research for long-term support of the Study. The authors also acknowledge the long-term contribution of the National Health and Medical Research Council of Australia. Funding for Core Management of the Raine Study was provided by the University of Western Australia (UWA); Telethon Kids Institute; Raine Medical Research Foundation; UWA Faculty of Medicine, Dentistry and Health Sciences; Women's and Infant's Research Foundation; and Curtin University. This work was supported by grants from the Raine Medical Research Foundation for the 10 year cohort review; National Health and Medical Research Council (NH&MRC) Programme Grants (ID 211912; ID 003209) and the Raine Medical Research Foundation for the 14 year cohort review; and NH&MRC Programme Grants (ID 35351417, ID 403981) for the 17 year follow-up.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

1. Johannsen NM, Swift DL, Lavie CJ, Earnest CP, Blair SN, Church TS. Categorical Analysis of the Impact of Aerobic and Resistance Exercise Training, Alone and in Combination, on Cardiorespiratory Fitness Levels in Patients With Type 2 Diabetes: Results from the HART-D study. *Diabetes Care*. 2013;36(10):3305-12.
2. Grøntved A, Ried-Larsen M, Ekelund U, Froberg K, Brage S, Andersen LB. Independent and Combined Association of Muscle Strength and Cardiorespiratory Fitness in Youth With Insulin Resistance and β -Cell Function in Young Adulthood: The European Youth Heart Study. *Diabetes Care*. 2013;36(9):2575-81.
3. Park Y-MM, Sui X, Liu J, Zhou H, Kokkinos PF, Lavie CJ, et al. The Effect of Cardiorespiratory Fitness on Age-Related Lipids and Lipoproteins. *Journal of the American College of Cardiology*. 2015;65(19):2091-100.
4. Clarkson HM, Gilewich GB. *Musculoskeletal Assessment: Joint Range of Motion and Manual Muscle Strength*. Baltimore, MD: Williams & Wilkins; 1989.
5. Bohannon RW, Magasi SR, Bubela DJ, Wang Y-C, Gershon RC. Grip and Knee Extension Muscle Strength Reflect a Common Construct among Adults. *Muscle & nerve*. 2012;46(4):555-8.
6. Wind A, Takken T, Helders P, Engelbert R. Is grip strength a predictor for total muscle strength in healthy children, adolescents, and young adults? *European Journal of Pediatrics*. 2010;169(3):281-7.
7. Ruiz JR, Sui X, Lobelo F, Morrow JR, Jackson AW, Sjöström M, et al. Association between muscular strength and mortality in men: prospective cohort study. *British Medical Journal*. 2008;337:a439.
8. Srikanthan P, Karlamangla AS. Relative Muscle Mass Is Inversely Associated with Insulin Resistance and Prediabetes. Findings from The Third National Health and Nutrition Examination Survey. *The Journal of Clinical Endocrinology & Metabolism*. 2011;96(9):2898-903.
9. Mainous Iii AG, Tanner RJ, Anton SD, Jo A. Grip Strength as a Marker of Hypertension and Diabetes in Healthy Weight Adults. *American Journal of Preventive Medicine*.
10. Leong DP, Teo KK, Rangarajan S, Lopez-Jaramillo P, Avezum A, Jr., Orlandini A, et al. Prognostic value of grip strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. *The Lancet*. 386(9990):266-73.
11. Sawyer SM, Afifi RA, Bearinger LH, Blakemore S-J, Dick B, Ezech AC, et al. Adolescence: a foundation for future health. *The Lancet*. 379(9826):1630-40.
12. Hulthén L, Bengtsson B-Å, Sunnerhagen KS, Hallberg L, Grimby G, Johannsson G. GH Is Needed for the Maturation of Muscle Mass and Strength in Adolescents. *The Journal of Clinical Endocrinology & Metabolism*. 2001;86(10):4765-70.
13. Huang R-C, Burrows S, Mori TA, Oddy WH, Beilin LJ. Lifecourse Adiposity and Blood Pressure Between Birth and 17 Years Old. *American Journal of Hypertension*. 2015.
14. Hangartner TN, Warner S, Braillon P, Jankowski L, Shepherd J. The Official Positions of the International Society for Clinical Densitometry: Acquisition of Dual-Energy X-Ray Absorptiometry Body Composition and Considerations Regarding Analysis and Repeatability of Measures. *Journal of Clinical Densitometry*. 2013;16(4):520-36.
15. McCarron LT. *McCarron Assessment of Neuromuscular Development*. Dallas, TX: McCarron Dial Systems; 1997.

16. Biering-Sørensen F. Physical measurements as risk indicators for low-back trouble over a one-year period. *Spine*. 1984;9(2):106-19.
17. Hands B, Larkin D, Parker H, Straker L, Perry M. The relationship among physical activity, motor competence and health-related fitness in 14-year-old adolescents. *Scandinavian Journal of Medicine & Science in Sports*. 2009;19(5):655-63.
18. Tanner JM. *Growth at Adolescence*. 2nd edn Blackwell Scientific Publications: Oxford 1962.
19. Huang R-C, Beilin LJ, Ayonrinde O, Mori TA, Olynyk JK, Burrows S, et al. Importance of cardiometabolic risk factors in the association between nonalcoholic fatty liver disease and arterial stiffness in adolescents. *Hepatology*. 2013;58(4):1306-14.
20. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the Concentration of Low-Density Lipoprotein Cholesterol in Plasma, Without Use of the Preparative Ultracentrifuge. *Clinical Chemistry*. 1972;18(6):499-502.
21. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;28(7):412-9.
22. Tobin J. Estimation of Relationships for Limited Dependent Variables. *Econometrica*. 1958;26(1):24-36.
23. Artero EG, Ruiz JR, Ortega FB, España-Romero V, Vicente-Rodríguez G, Molnar D, et al. Muscular and cardiorespiratory fitness are independently associated with metabolic risk in adolescents: the HELENA study. *Pediatric Diabetes*. 2011;12(8):704-12.
24. Peterson MD, Saltarelli WA, Visich PS, Gordon PM. Strength Capacity and Cardiometabolic Risk Clustering in Adolescents. *Pediatrics*. 2014;133(4):e896-e903.
25. Jiménez-Pavón D, Ortega FB, Valtueña J, Castro-Piñero J, Gómez-Martínez S, Zaccaria M, et al. Muscular strength and markers of insulin resistance in European adolescents: the HELENA Study. *European Journal of Applied Physiology*. 2012;112(7):2455-65.
26. Cohen DD, Gómez-Arbeláez D, Camacho PA, Pinzon S, Hormiga C, Trejos-Suarez J, et al. Low Muscle Strength Is Associated with Metabolic Risk Factors in Colombian Children: The ACFIES Study. *PLoS ONE*. 2014;9(4):e93150.
27. Zaqout M, Michels N, Bammann K, Ahrens W, Sprengeler O, Molnar D, et al. Influence of physical fitness on cardio-metabolic risk factors in European children. The IDEFICS study. *International Journal of Obesity*. 2016.
28. Sayer AA, Syddall HE, Dennison EM, Martin HJ, Phillips DIW, Cooper C, et al. The relationship between grip strength and features of the metabolic syndrome: findings from the Hertfordshire Cohort Study. *QJM : Monthly Journal of the Association of Physicians*. 2007;100(11):707-13.
29. Vaara JP, Fogelholm M, Vasankari T, Santtila M, Häkkinen K, Kyröläinen H. Associations of maximal strength and muscular endurance with cardiovascular risk factors. *International Journal of Sports Medicine*. 2014;35(4):356-60.
30. Viitasalo JT, Komi PV, Karvonen MJ. Muscle strength and body composition as determinants of blood pressure in young men. *European Journal of Applied Physiology and Occupational Physiology*. 1979;42(3):165-73.
31. Taekema DG, Maier AB, Westendorp RGJ, Craen AJMd. Higher Blood Pressure Is Associated With Higher Handgrip Strength in the Oldest Old. *American Journal of Hypertension*. 2011;24(1):83-9.
32. Straker LM, Hall GL, Mountain J, Howie EK, White E, McArdle N, et al. Rationale, design and methods for the 22 year follow-up of the Western Australian Pregnancy Cohort (Raine) Study. *BMC Public Health*. 2015;15:663.

33. Kelley GA, Kelley KS. Isometric handgrip exercise and resting blood pressure: a meta-analysis of randomized controlled trials. *Journal of Hypertension*. 2010;28(3):411-8.
34. Cornelissen VA, Fagard RH, Coeckelberghs E, Vanhees L. Impact of Resistance Training on Blood Pressure and Other Cardiovascular Risk Factors: A Meta-Analysis of Randomized, Controlled Trials. *Hypertension*. 2011;58(5):950-8.
35. Cornelissen VA, Smart NA. Exercise Training for Blood Pressure: A Systematic Review and Meta-analysis. *Journal of the American Heart Association*. 2013;2(1).
36. Owen A, Wiles J, Swaine I. Effect of isometric exercise on resting blood pressure: a meta analysis. *Journal of Human Hypertension*. 2010;24(12):796-800.
37. Horiuchi M, Okita K. Blood Flow Restricted Exercise and Vascular Function. *International Journal of Vascular Medicine*. 2012;2012:17.
38. Maslow AL, Sui X, Colabianchi N, Hussey J, Blair SN. Muscular Strength and Incident Hypertension in Normotensive and Prehypertensive Men. *Medicine and Science in Sports and Exercise*. 2010;42(2):288-95.
39. Sundström J, Neovius M, Tynelius P, Rasmussen F. Association of blood pressure in late adolescence with subsequent mortality: cohort study of Swedish male conscripts. *British Medical Journal*. 2011;342.
40. Alberti KGMM, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JJ, Donato KA, et al. Harmonizing the Metabolic Syndrome. *Circulation*. 2009;120(16):1640-5.
41. Notelovitz M. Androgen effects on bone and muscle. *Fertility and Sterility*. 2002;77, Supplement 4:34-41.
42. Reckelhoff JF, Roman RJ. Androgens and Hypertension: Role in Both Males and Females? *Hypertension*. 2011;57(4):681-2.
43. Adamopoulos S, Parissis JT, Paraskevaidis I, Karatzas D, Livanis E, Georgiadis M, et al. Effects of growth hormone on circulating cytokine network, and left ventricular contractile performance and geometry in patients with idiopathic dilated cardiomyopathy. *European Heart Journal*. 2003;24(24):2186-96.
44. Colao A, Marzullo P, Di Somma C, Lombardi G. Growth hormone and the heart. *Clinical Endocrinology*. 2001;54(2):137-54.
45. Steene-Johannessen J, Anderssen SA, Kolle E, Andersen LB. Low Muscle Fitness Is Associated with Metabolic Risk in Youth. *Medicine & Science in Sports & Exercise*. 2009;41(7):1361-7.
46. Grøntved A, Ried-Larsen M, Møller NC, Kristensen PL, Froberg K, Brage S, et al. Muscle strength in youth and cardiovascular risk in young adulthood (the European Youth Heart Study). *British Journal of Sports Medicine*. 2015;49(2):90-4.
47. Jung UJ, Choi M-S. Obesity and Its Metabolic Complications: The Role of Adipokines and the Relationship between Obesity, Inflammation, Insulin Resistance, Dyslipidemia and Nonalcoholic Fatty Liver Disease. *International Journal of Molecular Sciences*. 2014;15(4):6184-223.
48. Granacher U, Muehlbauer T, Gruber M. A Qualitative Review of Balance and Strength Performance in Healthy Older Adults: Impact for Testing and Training. *Journal of Aging Research*. 2012;2012:1-16.
49. Dodds RM, Syddall HE, Cooper R, Benzeval M, Deary IJ, Dennison EM, et al. Grip Strength across the Life Course: Normative Data from Twelve British Studies. *PLoS ONE*. 2014;9(12):e113637.

SUPPORTING INFORMATION

Supplementary Table 1a. Raw handgrip strength with systolic BP

Systolic BP	Unadjusted for adiposity model					Adjusted for adiposity model			
		Coefficient	P Value	95% CI		Coefficient	P value	95% CI	
Year (ref 10)									
	14	1.80	0.154	-0.67	4.27	0.04	0.977	-2.42	2.50
	17	2.71	0.031	0.25	5.17	-0.44	0.726	-2.93	2.04
Handgrip strength		0.30	<0.001	0.25	5.17	0.22	<0.001	0.15	0.29
Year*handgrip strength									
	14	-0.08	0.026	-0.15	-0.01	-0.04	0.282	-0.12	0.03
	17	-0.11	0.001	-0.18	-0.04	-0.05	0.124	-0.12	0.01
Sex (ref females)		1.58	<0.001	0.84	2.33	2.22	<0.001	1.49	2.95
BMI		N/A	N/A	N/A	N/A	0.52	<0.001	0.44	0.60
Constant		97.12	<0.001	95.02	99.22	89.55	<0.001	87.17	91.93
AIC		31919.38				31768.82			

N= 4425 observations from 1916 participants. CI, confidence interval; BP, blood pressure; BMI, body mass index; AIC, Akaike Information Criterion

Supplementary Table 1b. Height adjusted handgrip strength with systolic BP

Systolic BP	Unadjusted for adiposity model					Adjusted for adiposity model			
		Coefficient	P Value	95% CI		Coefficient	P value	95% CI	
Year (ref 10)									
	14	-0.22	0.884	-3.25	2.80	-1.37	0.371	-4.36	1.63
	17	0.15	0.92	-2.88	3.18	-2.12	0.168	-5.13	0.89
Handgrip strength		0.14	0.001	0.05	0.22	0.06	0.172	-0.02	0.14
Year*handgrip strength									
	14	0.03	0.471	-0.05	0.12	0.05	0.198	-0.03	0.14
	17	0.02	0.619	-0.06	0.10	0.06	0.161	-0.02	0.14
Sex (ref females)		3.34	<0.001	2.60	4.07	3.90	<0.001	3.20	4.61
BMI		N/A	N/A	N/A	N/A	0.60	<0.001	0.52	0.68
Constant		101.34	<0.001	98.86	103.82	92.29	<0.001	0.52	0.68
AIC		32130.51				31939.96			

N= 4425 observations from 1916 participants. CI, confidence interval; BP, blood pressure; BMI, body mass index; AIC, Akaike Information Criteria

Supplementary Table 1c. Weight adjusted handgrip strength with systolic BP

Systolic BP	Unadjusted for adiposity model					Adjusted for adiposity model			
		Coefficient	P Value	95% CI		Coefficient	P value	95% CI	
Year (ref 10)									
	14	-3.15	0.036	-6.10	-0.20	-6.03	<0.001	-8.93	-3.13
	17	-3.12	0.035	-6.03	-0.21	-7.56	<0.001	-10.44	-4.68
Handgrip strength		0.01	0.806	-0.07	0.09	0.06	0.132	-0.02	0.14
Year*handgrip strength									
	14	0.14	0.001	0.06	0.22	0.13	0.001	0.05	0.22
	17	0.14	0.001	0.06	0.23	0.13	0.002	0.05	0.21
Sex (ref females)		2.94	<0.001	2.16	3.71	2.72	<0.001	2.00	3.45
BMI		N/A	N/A	N/A	N/A	0.77	<0.001	0.69	0.85
Constant		105.38	<0.001	2.16	3.71	89.62	<0.001	86.72	92.52
AIC		32137.14				31809.22			

N= 4425 observations from 1916 participants. CI, confidence interval; BP, blood pressure; BMI, body mass index; AIC, Akaike Information Criterion

Supplementary Table 2: Longitudinal linear mixed model for the association between handgrip strength and systolic BP over time

	Unadjusted model		BMI adjusted model		BMI and PWC ₁₇₀ adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year						
14 years	3.35 (0.70, 6.00)	0.013	1.99 (-0.62, 4.61)	0.135	Reference group	Reference group
17 years	6.69 (3.87, 9.52)	<0.001	4.52 (1.72, 7.32)	0.002	3.70 (1.25, 6.14)	0.003
Sex Male	-2.23 (-4.22, -0.22)	0.030	-2.86 (-4.82, -0.90)	0.004	-0.34 (-3.67, 2.99)	0.842
Year*sex						
14 years	2.24 (0.65, 3.83)	0.006	2.19 (0.63, 3.75)	0.006	Reference group	Reference group
17 years	4.40 (2.08, 6.72)	<0.001	4.95 (2.68, 7.22)	<0.001	3.25 (1.49, 5.00)	<0.001
Handgrip strength	0.29 (0.21, 0.37)	<0.001	0.18 (0.11, 0.26)	<0.001	0.12 (0.06, 0.19)	<0.001
Year*handgrip strength						
14 years	-0.14 (-0.21, -0.07)	<0.001	-0.11 (-0.18, -0.04)	0.003	Reference group	Reference group
17 years	-0.22 (-0.29, -0.15)	<0.001	-0.19 (-0.26, -0.11)	<0.001	-0.10 (-0.15, -0.06)	<0.001
Sex*handgrip strength						
Males	0.06 (-0.0004, 0.12)	0.051	0.09 (0.03, 0.15)	0.002	0.09 (0.02, 0.15)	0.008

BMI	N/A	N/A	0.58 (0.50, 0.67)	<0.001	0.50 (0.40, 0.59)	<0.001
PWC ₁₇₀	N/A	N/A	N/A	N/A	-0.02 (-0.03, -0.007)	0.003
Constant	90.61 (88.08, 93.13)	<0.001	90.61 (88.08, 93.13)	<0.001	94.09 (90.79, 97.40)	97.40

N (unadjusted and BMI adjusted model) = 4425 observations from 1916 participants; (BMI and PWC₁₇₀ adjusted model) = 2617 observations from 1649 participants. CI, confidence interval; BP, blood pressure; BMI, body mass index; PWC₁₇₀, physical work capacity protocol

Supplementary Table 3: Longitudinal linear mixed model for the association between handgrip strength and log HOMA-IR over time

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year				
17 years	-0.38 (-0.44, -0.31)	<0.001	-0.46 (-0.52, -0.40)	<0.001
Sex Male	-0.12 (-0.19, -0.05)	0.001	-0.01 (0.08, -0.05)	0.663
Year*sex				
17 years	-0.06 (-0.16, 0.04)	0.221	0.06 (-0.03, 0.15)	0.210
Handgrip strength	0.003 (0.0006, 0.005)	0.013	-0.003 (-0.005, -0.002)	0.001
BMI	N/A	N/A	0.06 (0.05, 0.07)	<0.001
Constant	0.72 (0.61, 0.83)	<0.001	-0.39 (-0.53, -0.24)	<0.001

N= 2392 observations from 1505 participants. CI, confidence interval; HOMA-IR, homeostatic model assessment of insulin resistance; BMI, body mass index

Supplementary Table 4: Longitudinal linear mixed model for the association between handgrip strength and HDL-C over time

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year				
17 years	-0.15 (-0.22, -0.08)	<0.001	-0.12 (-0.18, -0.05)	0.001
Sex Male	-0.02 (-0.06, 0.01)	0.181	-0.05 (-0.09, -0.02)	0.001
Year*sex				
17 years	-0.11 (-0.15, -0.06)	<0.001	-0.14 (-0.18, -0.02)	<0.001
Handgrip strength	-0.004 (-0.006, -0.003)	<0.001	-0.003 (-0.004, -0.002)	<0.001
Year*handgrip strength				
17 years	0.003 (0.001, 0.004)	<0.001	0.003 (0.001, 0.004)	<0.001
BMI	N/A	N/A	-0.02 (-0.02, -0.01)	<0.001
Constant	1.64 (1.58, 1.70)	<0.001	2.02 (1.94, 2.09)	<0.001

N= 2396 observations from 1506 participants. CI, confidence interval; HDL-C, high density lipoprotein-cholesterol; BMI, body mass index

Supplementary Table 5: Random effects tobit model for the association between handgrip strength and log hs-CRP over time

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year				
17 years	0.43 (0.30, 0.56)	<0.001	0.35 (0.23, 0.48)	<0.001
Sex Male	0.06 (-0.09, 0.22)	0.439	0.22 (0.08, 0.37)	0.003
Year*sex				
17 years	-0.73 (-0.94, -0.53)	<0.001	-0.50 (-0.70, -0.31)	<0.001
Handgrip strength	0.005 (0.001, 0.01)	0.013	-0.006 (-0.01, -0.002)	0.007
BMI	N/A	N/A	0.11 (0.10, 0.13)	<0.001
Constant	-0.84 (-1.06, -0.62)	<0.001	-2.89 (-3.18, -2.59)	<0.001

N= 1967 observations from 1363 participants. CI, confidence interval; hs-CRP, high-sensitivity C-reactive protein; BMI, body mass index

Supplementary Table 6: Longitudinal linear mixed model for the association between back muscle endurance and systolic BP over time

	Unadjusted model		BMI adjusted model		BMI and PWC ₁₇₀ adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year*						
17 years	-0.34 (-1.17, 0.50)	0.429	-1.34 (-2.18, -0.51)	0.002	-1.40 (-2.27, -0.52)	0.002
Sex Male	4.99 (4.05, 5.95)	<0.001	5.39 (4.47, 6.31)	<0.001	5.55 (4.54, 6.57)	<0.001
Year*sex						
17 years	4.33 (3.17, 5.49)	<0.001	4.07 (2.93, 5.21)	<0.001	3.90 (2.67, 5.14)	<0.001
Back muscle endurance	-0.001 (-0.007, 0.005)	0.723	0.01 (0.004, 0.02)	0.002	0.01 (0.003, 0.02)	0.003
BMI	N/A	N/A	0.61 (0.52, 0.71)	<0.001	0.62 (0.52, 0.72)	<0.001
PWC ₁₇₀	N/A	N/A	N/A	N/A	0.003 (-0.01, 0.02)	0.654
Constant	108.86 (107.99, 109.72)	<0.001	94.69 (92.32, 97.07)	<0.001	94.21 (91.66, 96.76)	<0.001

N (unadjusted and BMI adjusted model) = 2712 observations from 1687 participants; (BMI and PWC₁₇₀ adjusted model) = 2557 observations from 1638 participants. CI, confidence interval; BP, blood pressure; BMI, body mass index; PWC₁₇₀, physical work capacity protocol

Supplementary Table 7: Longitudinal linear mixed model for the association between back muscle endurance and log HOMA-IR over time

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year				
17 years	-0.36 (-0.43, -0.30)	<0.001	-0.46 (-0.52, -0.39)	<0.001
Sex Male	-0.10 (-0.17, -0.04)	0.002	-0.06 (-0.12, -0.004)	0.036
Year*sex				
17 years	0.04 (-0.05, 0.12)	0.403	0.01 (-0.07, 0.10)	0.768
Back muscle endurance	-0.002 (-0.002, -0.001)	<0.001	-0.001 (-0.001, -0.0006)	<0.001
BMI	N/A	N/A	0.06 (0.05, 0.10)	<0.001
Constant	1.02 (0.97, 1.08)	<0.001	-0.29 (-0.44, -0.14)	<0.001

N= 2325 observations from 1491 participants. CI, confidence interval; HOMA-IR, homeostatic model assessment of insulin resistance; BMI, body mass index

Supplementary Table 8: Longitudinal linear mixed model for the association between back muscle endurance and log triglycerides over time

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year				
17 years	-0.0001 (-0.04, 0.04)	0.996	-0.04 (-0.08, -0.004)	0.033
Sex Male	-0.09 (-0.13, -0.04)	<0.001	-0.07 (-0.11, -0.03)	0.001
Year*sex				
17 years	0.10 (0.04, 0.15)	<0.002	0.09 (0.03, 0.14)	0.002
Back muscle endurance	-0.0009 (-0.001, -0.0006)	<0.001	-0.0004 (-0.0007, -0.0001)	0.007
BMI	N/A	N/A	0.03 (0.02, 0.03)	<0.001
Constant	0.03 (-0.007, 0.07)	0.112	-0.58 (-0.69, -0.47)	<0.001

N= 2325 observations from 1491 participants. CI, confidence interval; BMI, body mass index

Supplementary Table 9: Random effects tobit model for the association between back muscle endurance and log hs-CRP over time

	Unadjusted model		BMI adjusted model	
	Coefficient (95% CI)	<i>P</i> value	Coefficient (95% CI)	<i>P</i> value
Year				
17 years	0.45 (0.32, 0.59)	<0.001	0.34 (0.21, 0.46)	<0.001
Sex Male	0.10 (-0.04, 0.25)	0.169	0.15 (0.02, 0.29)	0.029
Year*sex				
17 years	-0.58 (-0.76, -0.39)	<0.001	-0.60 (-0.78, -0.43)	<0.001
Back muscle endurance	-0.003 (-0.004, -0.002)	<0.001	-0.001 (-0.002, -0.0002)	0.018
BMI	N/A	N/A	0.10 (0.09, 0.12)	<0.001
Constant	-0.31 (-0.44, -0.18)	<0.001	-2.81 (-3.14, -2.49)	<0.001

N= 1895 observations from 1337 participants. CI, confidence interval; hs-CRP, high-sensitivity C-reactive protein; BMI, body mass index

Supplementary Figure 1. Handgrip strength and systolic BP at 10, 14 and 17 years generated from longitudinal linear mixed models (mean and 95% CI's represented)

Supplementary Figure 2. Back muscle endurance and systolic BP at 14 and 17 years generated from longitudinal linear mixed models (mean and 95% CI's represented)