The Structure of CCl_5^- in the Gas Phase

Timothy R. Corkish, Christian T. Haakansson, Allan J. McKinley, and Duncan A.
Wild*

*School of Molecular Sciences, The University of Western Australia, Crawley, Western
Australia, 6009*

E-mail: duncan.wild@uwa.edu.au

Abstract

The first experimental evidence of the structure of the CCl_5^- gas phase anion complex is presented in conjunction with results from high level theoretical calculations. The photoelectron spectrum of the system shows a single peak with a maximum at 4.22 eV. CCSD(T) detachment energies of two stable C_{3v} ion-molecule complexes of the form $\text{Cl}^- \cdots \text{CCl}_4$ were also determined. The first complex found features the Cl^- bound linearly in a $\text{Cl}^- \cdots \text{Cl}-\text{C}$ bonding arrangement, while the second, less stable minimum has the Cl^- positioned at the face of the CCl_4 molecule, midway between three chlorine atoms. The calculated detachment energy for the first complex was found to be in excellent agreement with experiment, allowing the structure of CCl_5^- in the gas phase to be postulated as a non-covalent $\text{Cl}^- \cdots \text{CCl}_4$ anion complex, with the Cl^- anion tethered by a typical halogen bond.

Graphical TOC Entry

The structure of a chloride ion bound to a CCl_4 molecule in the gas phase has not yet been the subject of experimental scrutiny. The existence of a stable form of the CCl_5^- anion was first proposed in 1966 by McDaniel and Deiters, who suggested either a $\text{Cl}_3\text{C}-\text{Cl}-\text{Cl}^-$ bonding arrangement or a penta-coordinated carbon structure.¹ Later X-ray data of crystalline carbon tetrahalide complexes showed that one halide ion is bound to each apex of the core molecule,² while Raman studies found that in solution, the halide is positioned at the ‘face’ of the carbon tetrahalide, equally bound to three halogen atoms.^{3,4} Despite this, no structural information has been gained relative to the gas phase species; although several mass spectrometry experiments have confirmed its presence,⁵ with estimations for the heat of association as approximately -60 kJ mol^{-1} .⁶⁻⁸

The nomenclature ‘ CCl_5^- ’ implies a stable, hypervalent, and pentahalogenated carbon species of D_{3h} symmetry, although theoretical studies have since confirmed that this purported structure is a saddle point linking two degenerate $\text{C}_{3v} \text{Cl}^- \cdots \text{CCl}_4$ complexes along an $\text{S}_{\text{N}}2$ style pathway.⁹⁻¹² The stable complex in this form involves the halide at the face of the CCl_4 molecule. In addition, a second, more stable minimum has been predicted by theory, with the chloride bound linearly to a Cl atom of CCl_4 in what resembles a three-centred, four-electron bonding motif.^{8,11-13} However, no experimental results to date have been able to elucidate the structure of CCl_5^- , henceforth referred to as $\text{Cl}^- \cdots \text{CCl}_4$, in the gas phase.

The aim of this paper is to present a combination of theoretical and experimental results that support the existence of one stable isomer of $\text{Cl}^- \cdots \text{CCl}_4$ over another, and ascertain the gas phase structure of the anion complex for the first time. It will be shown through *ab initio* calculations that each isomer can be experimentally distinguished via their individual electron binding energies (*eBE*). Subsequent experimental data, in the form of mass spectrometry and photoelectron spectroscopy, will be used to present evidence as to the structure of the $\text{Cl}^- \cdots \text{CCl}_4$ anion complex in the gas phase.

Figure 1 displays the optimised geometries of the $\text{Cl}^- \cdots \text{CCl}_4$ isomers at the MP2/aug-cc-pVQZ level of theory, using the aug-cc-pV(Q+d)Z basis set for Cl.^{14,15} Vibrational analyses

found all real frequencies for both conformers, while further details on structures and energetics can be found in the Supporting Information attachment, in addition to the optimised geometry of the D_{3h} saddle point mentioned previously. CCSD(T) energies were computed on these stable complexes with all values extrapolated to the complete basis set limit (CBS) using the W2w method unless stated otherwise.¹⁶

Figure 1: Optimised MP2/aug-cc-pVQZ (with the aug-cc-pV(Q+d)Z basis set for Cl) geometries of the C_{3v} $Cl^- \cdots CCl_4$ gas phase anion complexes.

The first C_{3v} chlorine-bound complex, to be referred to as Complex 1 as per Figure 1, features the Cl^- anion tethered linearly to one Cl of CCl_4 at a distance of 2.780 Å, and has a zero-point corrected dissociation energy (D_0) of 46.7 kJ mol⁻¹. Carrion and Dewar¹³ first reported this structure with an accompanying $Cl^- \cdots Cl$ distance of 2.13 Å and a heat of formation of 134 kJ mol⁻¹ using the MNDO method, prompting a call to label the ion as being covalently bound. Subsequent HF investigations by Kobychiev et al.¹¹ as well as Vetter and Zülicke¹² found this distance to be 2.93 Å and 2.92 Å respectively, with dissociation energies of 43.1 kJ mol⁻¹ and 37.7 kJ mol⁻¹. Hiraoka et al. converged on this complex as well, reporting a B3LYP/6-31+G* distance of 2.82 Å and a QCISD(T) dissociation energy of 42.7 kJ mol⁻¹.⁸ With the exception of the MNDO calculations, we present the shortest $Cl^- \cdots Cl$ distance reported for the anion complex, while all dissociation energies are in relative agreement.

In contrast, in the second C_{3v} structure of this work, labelled Complex 2, the Cl^- anion is equidistant to three Cl atoms of CCl_4 , 3.563 Å apart, and has a significantly lower D_0 value of 20.2 kJ mol $^{-1}$. The $\text{Cl}^- \cdots \text{C}$ distance is 3.708 Å for our reported structure. This distance has been presented in the range from 3.78 Å to 4.15 Å in previous theoretical studies, with dissociation energies from 7.5 kJ mol $^{-1}$ to 19.7 kJ mol $^{-1}$.⁸⁻¹² Additionally, Marchaj et al. most recently argued, upon taking into consideration thermal and entropic corrections, that the $\text{Cl}^- \cdots \text{CCl}_4$ geometry of Complex 2 is unlikely to be observed experimentally, especially at elevated temperatures.⁹ Indeed, a simple Boltzmann distribution calculated here shows that the relative proportion of Complex 2 would approach 0.002% at 1000 K, and would be negligible at the typical experimental beam temperature of 30 K.

Based on the D_0 values of each complex presented in this work, it would be expected that Complex 1 is the more likely to be observed of the two isomers. Table 1 summarises the structural parameters of these systems, in addition to presenting the CCSD(T)/CBS vertical detachment energies (VDE) for each complex. These energies correspond to a simulation of the detachment of an electron associated with the Cl^- anion of each structure, with the calculated geometries suggesting a non-covalent van der Waals interaction, as opposed to the formation of a molecular anion. Raw photodetachment energies were split into transitions to the $^2\text{P}_{3/2}$ and $^2\text{P}_{1/2}$ neutral states of Cl using the experimental spin-orbit coupling constant.

Table 1: Summary of structural parameters of the $\text{Cl}^- \cdots \text{CCl}_4$ complexes. Geometrical information is from the relevant MP2 geometry optimised complex while the D_0 and VDE values were determined from CCSD(T)/CBS energies. See text for further information.

	$r_{\text{Cl}^- \cdots \text{Cl}_1}$	$r_{\text{Cl}^- \cdots \text{Cl}_2}$	D_0	VDE	
	(Å)	(Å)	(kJ mol $^{-1}$)	(eV)	
				$^2\text{P}_{3/2}$	$^2\text{P}_{1/2}$
Complex 1	2.780	5.426	46.7	4.23	4.34
Complex 2	5.500	3.563	20.2	3.72	3.83

Complex 1 has VDE values of 4.23 eV and 4.34 eV for transitions to the $^2P_{3/2}$ and $^2P_{1/2}$ states of Cl respectively, while these values are 3.72 eV and 3.83 eV for Complex 2. For the case of Complex 2, the data is in good agreement with the 3.83 eV value presented by Marchaj et al.⁹ That the electron would be more tightly bound to the Cl^- of Complex 1 by more than 0.5 eV can be rationalised by the strength of anion D_0 values in relation to one another.

Needless to say, the large difference in energy between the two detachment energies suggests that the presence of each conformer can be determined through experimental photoelectron spectroscopy. It is also noted at this point, that VDEs produced from theory to predict or rationalise experimental photoelectron spectra have been used previously with great success, including those at the CCSD(T) level of theory.¹⁷⁻¹⁹

Figure 2: A mass spectrum over the range of 140-200 m/z from a gas mixture of CCl_4 , $\text{CH}_3\text{CH}_2\text{OH}$, and Ar. The cluster of peaks from 187 m/z to 193 m/z corresponds to the presence of the $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion complex.

Shown in Figure 2 is a mass spectrum resulting from a gas mixture consisting of CCl_4 , $\text{CH}_3\text{CH}_2\text{OH}$, and made up to 400 kPa with Ar. The ethanol added to the mixture was in an attempt to characterise the $\text{Cl}^- \cdots \text{CH}_3\text{CH}_2\text{OH}$ system, although has no relevance to the investigation at hand, with the presence of the $\text{Cl}^- \cdots \text{CCl}_4$ anion complex detected in unrelated Cl^- based experiments. Cl^- anions were produced via a process of dissociative attachment to

CCl_4 , with electrons provided by a hot rhenium filament. The major characterising feature from the m/z region between 140 and 200 is a series of four peaks at m/z values of 187, 189, 191, and 193, highly indicative of the touted $\text{Cl}^- \cdots \text{CCl}_4$ system. The relative abundance of each mass peak is in good agreement with those that have been previously reported,⁷ and follows from the natural abundance of the chlorine isotopes. Also present in the mass spectrum are complexes resulting from a residual amount of CH_3I of a previous experiment; notably $\text{I}^- \cdots \text{H}_2\text{O}$ at 145 m/z , $\text{I}^- \cdots \text{Ar}$ at 167 m/z , and $\text{Cl}^- \cdots \text{CH}_3\text{I}$ at 177 and 179 m/z .

Photoelectron spectra were collected for the bare Cl^- and I^- ions for calibration, while the spectrum of the $\text{Cl}^- \cdots \text{CCl}_4$ system was recorded using the largest mass peak occurring at 189 m/z . This photoelectron spectrum is reported in Figure 3, characterised by a single peak with a maximum at 4.22 eV, the experimentally determined eBE . The 0.18 eV spread in photoelectron energy of this peak, resulting from the mass of the complex and a beam energy of 1500 eV,²⁰ in turn leading to a full width half maximum of 0.07 eV, obscures any possible distinction of a perturbed $^2\text{P}_{1/2}$ Cl spin-orbit state, as would be expected for a Cl^- anion core interacting non-covalently with an adduct molecule. However, there is a small shoulder occurring at slightly higher eBE which could amount to a small protrusion of the $^2\text{P}_{1/2}$ neutral state, or even a vibrational progression, as has been observed for similar systems.^{21,22}

The general shape of the novel peak is consistent with that of the bare Cl^- photoelectron spectrum shifted approximately 0.58 eV to higher eBE , as the maximum of the bare halide photoelectron peak was measured at 3.64 eV. The magnitude of this electron stabilisation energy (E_{stab}) is usually indicative of the strength of the van der Waals interaction between the halide anion and complexing molecule. For example, the $\text{Cl}^- \cdots \text{N}_2$ ion-induced dipole complex experiences an E_{stab} of 0.11 eV,²³ whereas this figure is 0.76 eV for the ion-dipole $\text{Cl}^- \cdots \text{H}_2\text{O}$ gas phase complex.²⁴ The E_{stab} of 0.58 eV observed in the current experiment is therefore indicative that the $\text{Cl}^- \cdots \text{CCl}_4$ complex is bound rather tightly, likely through the halogen bond motif of Complex 1, which is debated to comprise of purely coulombic

Figure 3: Photoelectron spectrum of the $\text{Cl}^- \cdots \text{CCl}_4$ gas-phase anion complex from 0 to 4.661 eV. The maximum of the photoelectron peak occurs at 4.22 eV. The theoretical VDEs to a neutral $^2\text{P}_{3/2}$ state of Complex 1 (dashed line) and Complex 2 (dotted line) are overlayed for comparison.

terms, being electrostatic and polarisation,^{25,26} in addition to some contribution resulting from charge transfer.^{27,28}

Figure 3 also shows the comparison between the experimental photoelectron spectrum (black line) and the $^2\text{P}_{3/2}$ VDE predictions of Complex 1 (dashed line) and Complex 2 (dotted line). It is readily apparent that the high level predictions for Complex 1 are in excellent agreement with the experimentally determined eBE . Therefore, the results we present here are indicative that the structure of $\text{Cl}^- \cdots \text{CCl}_4$ in the gas phase is that of a chloride ion forming a van der Waals complex with the CCl_4 molecule, bound linearly to a Cl atom as per the depiction of Complex 1 in Figure 1.

Additionally, there appears to be no indication that there is any photodetachment peak resulting from the presence of Complex 2. There is a small portion of noise between 3.50-4.00 eV in Figure 3, however this is likely due to background subtraction becoming more pronounced in a region where the applied Jacobi transform is beginning to show an effect. This

transform involves multiplying the photoelectron intensities by their time-of-flight cubed, t^3 , in order to move from time-dependent bins to energy-dependent. Therefore, we can say with confidence that little to none of the experimental $\text{Cl}^- \cdots \text{CCl}_4$ mass peak results from the existence of Complex 2, under the present experimental conditions.

The only caveat to the experimental results is that the $^2\text{P}_{1/2}$ neutral state of Cl is obscured; a feature that would be observable at higher resolution. The distinguishing of both neutral states of Cl would allow the conclusion to be made, beyond reasonable doubt, that the proposed CCl_5^- system is best described by a chloride ion adducting non-covalently to CCl_4 , although the combination of theory and experiment presented here does support the existence of a $\text{Cl}^- \cdots \text{CCl}_4$ van der Waals complex.

To complement the experimental results, we have undertaken two theoretical approaches to determine the bond index of the $\text{Cl}^- \cdots \text{Cl}$ bond in Complex 1. The first involves the methodology set out by Oliveira et al., using calculated local stretching force constants to determine a bond strength order (BSO).²⁹ This method yields a BSO of 0.219 for the $\text{Cl}^- \cdots \text{Cl}$ bond, and 1.006 for the $\text{Cl}-\text{C}$ bond along the axis of the Cl^- adduct. Interestingly, this would correspond to a percentage of three-centred four-electron bonding of 22%. The BSO of the $\text{Cl}^- \cdots \text{Cl}$ bond would place it in the category of ‘normal halogen bond’ in the bond strength hierarchy put forward by Oliveira et al.²⁹ As a note, the force constants used in this calculation can be found along with vibrational modes in the Supporting Information.

The second method utilised the Roby-Gould bond index program using the Tonto computational chemistry package.³⁰ Further information about this method has been reported.^{31–33} The overall $\text{Cl}^- \cdots \text{Cl}$ bond index was found to be 0.269 for the geometry of Complex 1, in what is reasonable agreement with the index predicted through stretching force constants. These results are consistent with the conclusion that the $\text{Cl}^- \cdots \text{CCl}_4$ complex studied experimentally in this work is likely to consist of a chloride ion bound non-covalently to a CCl_4 molecule in a $\text{Cl}^- \cdots \text{Cl}-\text{C}$ bonding arrangement.

In summary, the first experimental evidence as to the structure of the CCl_5^- , or rather,

the $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion species has been presented. Mass spectrometry experiments were used to confirm the production of the system, before characterisation with photoelectron spectroscopy. The photoelectron spectrum features a single peak with a maximum at 4.22 eV, found to be in excellent agreement with the CCSD(T)/CBS predicted VDE of Complex 1 of Figure 1, which was determined to be 4.23 eV and 4.34 eV for detachments to the neutral $^2\text{P}_{3/2}$ and $^2\text{P}_{1/2}$ states of Cl respectively.

The structure of this C_{3v} complex has the halide appended non-covalently to a singular Cl of CCl_4 in what resembles a typical halogen bond, and is bound by 46.7 kJ mol $^{-1}$. Based on a combination of the theoretical and experimental results, we assign this complex as being the form of $\text{Cl}^- \cdots \text{CCl}_4$ in the gas phase.

A second stable C_{3v} complex was also found by *ab initio* calculations, with the Cl^- anion positioned at the face of the CCl_4 molecule, and this is thought to be a more purely electrostatically driven interaction. Complex 2 was found to have a dissociation energy of 20.2 kJ mol $^{-1}$, less than half that of Complex 1. Rationalising the theoretical VDE predictions in relation to the experiment, there is no indication that Complex 2 was observed experimentally.

The experimental data outlined in this paper was recorded using a home built time of flight mass spectrometer coupled to a photoelectron spectrometer. The design of the mass spectrometer is owed to Wiley and McLaren,³⁴ while the magnetic-bottleneck feature of the photoelectron spectrometer is derived from Cheshnovsky.²⁰ Further information on the experimental design has been reported.³⁵

The fourth harmonic of a Nd:YAG laser (Spectra Physics Quanta Ray Pro) was used to generate the 4.661 eV photons used for this experiment. Upon photodetachment, electron counts were recorded as per their time of flight, before conversion to kinetic energy and subsequently *eBE* based on the energy of the input photon. Photoelectron spectra were recorded over 10 000 shots, before averaging, as well as the application of a Jacobi transform.

All calculations were carried out using the Gaussian 09 program,³⁶ unless stated otherwise. As mentioned, geometries and frequencies were determined at the MP2/aug-cc-PVQZ level of theory, with the aug-cc-pV(Q+d)Z basis set used for Cl.^{14,15} Very tight convergence criteria ($1 \times 10^{-8} E_h a_0^{-1}$) were applied during optimisations. Single point CCSD(T) energies on both the stable anion minima and the neutral vertically-detached counterparts were determined up to the quality of aug-cc-pV5Z/aug-cc-pV(5+d)Z before CBS extrapolation. Further details on the results of this process are outlined in the Supporting Information.

Acknowledgements

This research was undertaken with the assistance of computational resources from the Pople high-performance computing cluster of the Faculty of Science at the University of Western Australia. The Australian Research Council is acknowledged for funding the laser installation under the LIEF scheme (LE110100093). The School of Molecular Sciences and the Faculty of Science are acknowledged for financial support. T.R.C. thanks the support of an Australian Government Research Training Program (RTP) scholarship, while C.T.H. acknowledges the UWA Dean’s Excellence in Science PhD Scholarship. The authors would also like to thank Prof. Dylan Jayatilaka for his help with the Roby-Gould bond index program.

Supporting Information Available

- SuppInfo.pdf: Energetics, vibrational modes, and cartesian coordinates of the stable C_{3v} complexes outlined in this work, in addition to the D_{3h} saddle point.

References

- (1) McDaniel, D. H.; Deiters, R. M. The CCl_5^- Species. *J. Am. Chem. Soc.* **1966**, *88*, 2607-2608.

- (2) Creighton, J. A.; Thomas, K. M. Preparation and Structures of Halide Ion Complexes of Carbon Tetrahalides and Tetrahalogenoethylenes. *J. Chem. Soc., Dalton Trans.* **1972**, 403-410.
- (3) Creighton, J. A.; Thomas, K. M. Raman Spectroscopic Studies of Halide-Ion Complexes of Carbon Tetrahalides in Solution. *J. Chem. Soc., Dalton Trans.* **1972**, 2254-2257.
- (4) Creighton, J. A.; Thomas, K. M. The Structure and Bonding in Halide Ion Complexes of Carbon Tetrahalides. *J. Mol. Structure.* **1971**, 7, 173-178.
- (5) Kebabian, P.; Arshadi, M.; Scarborough, J. Hydration of Negative Ions in the Gas Phase. *J. Chem. Phys.* **1968**, 49, 817-822.
- (6) Dougherty, R. C.; Dalton, J.; Roberts, J. S_N2 Reactions in the Gas Phase. Structure of the Transition State. *Organic Mass Spectrometry*, **1973**, 8, 77-79.
- (7) Staneke, P. O.; Groothuis, G.; Ingemann, S.; Nibbering, N. M. M. Formation, Stability and Structure of Radical Anions of Chloroform, Tetrachloromethane and Fluorotrichloromethane in the Gas Phase. *Int. J. Mass Spectrom. Ion Process.* **1995**, 142, 83-93.
- (8) Hiraoka, K.; Mizuno, T.; Iino, T.; Eguchi, D.; Yamabe, S. Characteristic Changes of Bond Energies for Gas-Phase Cluster Ions of Halide Ions with Methane and Chloromethanes. *J. Phys. Chem. A* **2001**, 105, 4887-4893.
- (9) Marchaj, M.; Freza, S.; Skurski, P. Why Are SiX₅⁻ and GeX₅⁻ (X = F, Cl) Stable but Not CF₅⁻ and CCl₅⁻? *J. Phys. Chem. A* **2012**, 116, 1966-1973.
- (10) Gutsev, G. L. A Theoretical Investigation on the Structure of the Hypervalent Carbon and Silicon Pentahalogenides as Well as Their Singly Charged Anions. *Chem. Phys.* **1992**, 166, 57-68.

- (11) Kobychiev, V. B.; Vitkovskaya, N. M.; Abramoc, A. V.; Timokhin, B. V. Quantum-Chemical Study of Halophilic Interactions: III. *Ab initio* Study of the Attack of Tetra-chloromethane by Halide Ions. *Russ. J. Gen. Chem.*, **1999**, *69*, 788-793.
- (12) Vetter, R.; Zülicke, L. Non-empirical Study of the Structure and Stability of Penta-halocarbonate Anions CX_5^- ($X = F, Cl, Br$). All-electron and Valence-electron SCF Calculations. *Chem. Phys.* **1986**, *101*, 201-209.
- (13) Carrion, F.; Dewar, M. J. S. Ground States of Molecules. 59. MNDO Study of S_N2 Reactions and Related Processes. *J. Am. Chem. Soc.* **1984**, *106*, 3531-3539.
- (14) Kendall, R. A.; Dunning, Jr., T. H.; Harrison, R. J. Electron Affinities of the First-Row Atoms Revisited. Systematic Basis Sets and Wave Functions. *J. Chem. Phys.* **1992**, *96*, 6796-6806.
- (15) Dunning, Jr., T. H.; Peterson, K. A.; Wilson, A. K. Gaussian Basis Sets For Use in Correlated Molecular Calculations. X. The Atoms Aluminum Through Argon Revisited *J. Chem. Phys.* **2001**, *114*, 9244-9253.
- (16) Karton, A.; Rabinovic, E.; Martin, J. M. L. W4 Theory for Computational Chemistry: In Pursuit of Confident sub-kJ/mol Predictions. *J. Chem. Phys.* **2006**, *125*, 144108.
- (17) Elliott, B. M.; Koyle, E.; Boldyrev, A. I.; Wang, X-B.; Wang, L-S. MX_3^- Superhalogens ($M = Be, Mg, Ca$; $X = Cl, Br$): A Photoelectron Spectroscopic and *ab Initio* Theoretical Study. *J. Phys. Chem. A* **2005**, *109*, 11560-11567.
- (18) Alexandrova, A. N.; Boldyrev, A. I.; Fu, Y-J.; Yang, X.; Wang, X-B.; Wang, L-S. Structure of the $Na_xCl_{x+1}^-$ ($x=1-4$) Clusters via *ab Initio* Genetic Algorithm and Photoelectron Spectroscopy. *J. Chem. Phys.* **2004**, *121*, 5709-5719.
- (19) Lapere, K. M. L.; Kettner, M.; Watson, P. D.; McKinley, A. J.; Wild, D. A. The

- Halide-Nitrogen Gas-Phase Clusters: Anion Photoelectron Spectroscopy and High Level *ab Initio* Calculations. *J. Phys. Chem. A*, **2015**, *119*, 9722-9728.
- (20) Cheshnovsky, O. H.; Yang, S. H.; Pettiette, C. L.; Craycraft, M. J.; Smalley, R. E. Magnetic Time-Of-Flight Photoelectron Spectrometer For Mass-Selected Negative Cluster Ions. *Rev. Sci. Instrum.* **1987**, *58*, 2131-2137.
- (21) Cyr, D. M.; Bisea, G. A.; Scarton, M. G.; Johnson, M. A. Observation of Charge-Transfer Excited States in the $\text{I}^- \cdot \text{CH}_3\text{I}$, $\text{I}^- \cdot \text{CH}_3\text{Br}$, and $\text{I}^- \cdot \text{CH}_2\text{Br}_2$ $\text{S}_\text{n}2$ Reaction Intermediates Using Photofragmentation and Photoelectron Spectroscopies. *J. Chem. Phys.* **1992**, *97*, 5911-5914.
- (22) Van Duzor, M.; Wei, J.; Mbaiwa, F.; Mabbs, R. $\text{I}^- \cdot \text{CH}_3\text{X}$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) Photodetachment: The Effect of Electron-Molecule Interactions in Cluster Anion Photodetachment Spectra and Angular Distributions. *J. Chem. Phys.* **2010**, *133*, 144303.
- (23) P. D. Watson, H. Yong, K. M. L. Lapere, M. Kettner, A. J. McKinley, D. A. Wild. Anion Photoelectron Spectroscopy and CCSD(T) Calculations of the $\text{Cl}^- \cdots \text{N}_2$ Complex. *Chem. Phys. Lett.* **2016**, *654*, 119.
- (24) Markovich, G.; Pollack, S.; Giniger, R.; Cheshnovsky, O. Photoelectron Spectroscopy of Cl^- , Br^- , and I^- Solvated in Water Clusters. *J. Chem. Phys.* **1994**, *101*, 9344-9353.
- (25) Clark, T.; Murray, J.; Politzer, P. A Perspective on Quantum Mechanics and Chemical Concepts in Describing Noncovalent Interactions. *Phys. Chem. Chem. Phys.* **2018**, *20*, 30076-30082.
- (26) Clark, T.; Heßelmann, A. The Coulombic σ -hole Model Describes Bonding in $\text{CH}_3\text{I} \cdots \text{Y}^-$ Complexes. *Phys. Chem. Chem. Phys.* **2018**, *20*, 22849-22855.
- (27) Řezáč, J.; de la Lande, A. On the Role of Charge Transfer in Halogen Bonding. *Phys. Chem. Chem. Phys.* **2017**, *19*, 791-803.

- (28) Thirman, J.; Engelage, E.; Huber, S. M.; Head-Gordon, M. Characterizing the Interplay of Pauli Repulsion, Electrostatics, Dispersion and Charge Transfer in Halogen Bonding with Energy Decomposition Analysis. *Phys. Chem. Chem. Phys.* **2018**, *20*, 905-915.
- (29) Oliveira, V.; Kraka, E.; Cremer, D. The Intrinsic Strength of the Halogen Bond: Electrostatic and Covalent Contributions Described by Coupled Cluster Theory. *Phys. Chem. Chem. Phys.* **2016**, *18*, 33031-33046.
- (30) Jayatilaka, D.; Grimwood, D. J. Tonto: A Fortran Based Object-Oriented System for Quantum Chemistry and Crystallography. In *Computational Science - ICCS 2003, Part 4*; Springer Verlag: New York, 2003; pp142-151. <https://github.com/dylan-jayatilaka/tonto> (accessed Aug 21, 2019)
- (31) Gould, M. D.; Taylor, C.; Wolff, S. K.; Chandler, G. S.; Jayatilaka, D. A Definition For the Covalent and Ionic Bond Index in a Molecule. *Theor. Chem. Acc.* **2008**, *119*, 275-290.
- (32) Alhameedi, K.; Bohman, B.; Karton, A.; Jayatilaka, D. Predicting the Primary Fragments in Mass Spectrometry Using *ab Initio* Roby-Gould Bond Indices. *Int. J. Quantum Chem.* **2018**, *118*, e25603.
- (33) Wońska, M.; Grabowsky, S.; Dominiak, P. M.; Wozniak, K.; Jayatilaka, D. Hydrogen Atoms Can Be Located Accurately and Precisely By X-ray Crystallography. *Sci. Adv.* **2016**, *2*, e1600192.
- (34) Wiley, W. C.; McLaren, I. H. Time-Of-Flight Mass Spectrometer With Improved Resolution. *Rev. Sci. Instrum.* **1955**, *26*, 1150-1157.
- (35) Lapere, K. M.; LaMacchia, R. L.; Quak, L. H.; McKinley, A. J.; Wild, D. A. Anion Photoelectron Spectroscopy and *ab Initio* Calculations of the Gas Phase Chloride-Carbon Monoxide Complex: $\text{Cl}^- \cdots \text{CO}$. *Chem. Phys. Lett.* **2011**, *504*, 13-19.

- (36) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian 09 Revision D.01, Gaussian Inc., Wallingford CT, 2013.

The Structure of CCl_5^- in the Gas Phase

Timothy R. Corkish, Christian T. Haakansson, Allan J. McKinley, Duncan A. Wild

School of Molecular Sciences, The University of Western Australia, Crawley, Western Australia, 6009, Australia

SUPPORTING INFORMATION

The supporting information presented here comprises results of *ab initio* calculations performed on $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion complexes, and the CCl_5^- transition state. Included are the energies, vibrational data, and cartesian coordinates predicted at the MP2 and CCSD(T) levels of theory, with aug-cc-pVTZ, QZ, and 5Z basis sets. The aug-cc-pV(X+d)Z basis sets were used for chlorine. Collectively, these basis sets will be referred to as AVXZ.

1 Energetics

Table S1: Energies of the bare chloride anion and chlorine radical determined from CCSD(T) calculations.

		$E_{\text{CCSD(T)}}$ [E_h]	VDE [eV]	Experimental SO* [eV]	Split [eV]	Lit. $^2P_{3/2}^\dagger$ [eV]	Shift ‡ [eV]
Cl ⁻ Cl	AVTZ	-459.8066265 -459.6778578	3.504		3.468 3.577		+0.145
	AVQZ	-459.6958913 -459.8283951	3.606		3.570 3.679		+0.043
	AV5Z	-459.7013126 -459.8348346	3.633		3.597 3.706		+0.016
	CBS	-459.7066880 -459.8412328	3.661	-0.036 +0.073	3.625 3.734	3.613	-0.012

* Values from <http://www.nist.gov/pml/data/handbook/index.cfm>

† Values from <http://webbook.nist.gov>

‡ Shift refers to the difference between the predicted and literature Electron Detachment Energy

Table S2: Energies of the C_{3v} halogen-appended Cl⁻...CCl₄ gas phase anion complex and neutral counterpart predicted from CCSD(T) calculations.

		Anion _{(CCSD(T))} [E_h]	Neutral _{(CCSD(T))} [E_h]	zpe [kJ mol ⁻¹]	De [kJ mol ⁻¹]	D ₀ [†] [kJ mol ⁻¹]	VDE [†] [eV]
Cl ⁻ ...CCl ₄	AVTZ	-2336.7998247	-2336.6479978				
	AVQZ	-2336.9106662	-2336.7549268	26.7*			
	AV5Z	-2336.9447580	-2336.7879148				
	CBS	-2336.9785121	-2336.8205701		47.5	46.7	4.23 4.34

* MP2/AVQZ value

† Determined using CCSD(T)/CBS energies and MP2/AVQZ zpe

Table S3: Energies of the C_{3v} Cl⁻...CCl₄ gas phase anion complex, featuring the halide at the face of the CCl₄ molecule, and neutral counterpart predicted from CCSD(T) calculations.

		Anion _{(CCSD(T))} [E_h]	Neutral _{(CCSD(T))} [E_h]	zpe [kJ mol ⁻¹]	De [kJ mol ⁻¹]	D ₀ [†] [kJ mol ⁻¹]	VDE [†] [eV]
Cl ⁻ ...CCl ₄	AVTZ	-2336.7899093	-2336.6539300				
	AVQZ	-2336.9006398	-2336.7607232	26.6*			
	AV5Z	-2336.9347036	-2336.7937173				
	CBS	-2336.9684218	-2336.8296220		21.0	20.2	3.72 3.83

* MP2/AVQZ value

† Determined using CCSD(T)/CBS energies and MP2/AVQZ zpe

2 Vibrational Data

Table S4: Vibrational frequencies for the halogen-appended $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion complex from MP2/AVQZ calculations. Frequencies in cm^{-1} , zero point energies (zpe) in kJ mol^{-1} , force constants in $\text{mDyne \AA}^{-1}$.

Mode	Symmetry	Frequency	Force constant
ω_1	A_1	786	4.749*
ω_2	E	775	4.609
ω_3	E	775	4.609
ω_4	A_1	448	4.128
ω_5	E	326	2.048
ω_6	E	326	2.048
ω_7	A_1	309	1.799
ω_8	E	226	1.053
ω_9	E	226	1.053
ω_{10}	A_1	125	0.317*
ω_{11}	E	68	0.093
ω_{12}	E	68	0.093
zpe	26.7		

*Respective $\text{Cl}-\text{C}$ and $\text{Cl}^- \cdots \text{Cl}$ local stretching force constants used in main text to determine an approximate bond strength order.

Table S5: Vibrational frequencies for the $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion complex, featuring the halide at the face of the CCl_4 molecule, from MP2/AVQZ calculations. Frequencies in cm^{-1} , zero point energies (zpe) in kJ mol^{-1} , force constants in $\text{mDyne \AA}^{-1}$.

Mode	Symmetry	Frequency	Force constant
ω_1	E	826	5.297
ω_2	E	826	5.297
ω_3	A_1	753	4.394
ω_4	A_1	475	4.584
ω_5	E	323	1.931
ω_6	E	323	1.931
ω_7	A_1	313	1.867
ω_8	E	221	1.004
ω_9	E	221	1.004
ω_{10}	A_1	77	0.117
ω_{11}	E	48	0.046
ω_{12}	E	48	0.046
zpe	26.6		

Table S6: Vibrational frequencies for the CCl_5^- transition state from MP2/AVQZ calculations. Frequencies in cm^{-1} , zero point energies (zpe) in kJ mol^{-1} , force constants in $\text{mDyne \AA}^{-1}$.

Mode	Symmetry	Frequency	Force constant
ω_1	E'	919	6.562
ω_2	E'	919	6.562
ω_3	A_1'	460	4.365
ω_4	A_2''	302	1.724
ω_5	E'	289	1.535
ω_6	E'	289	1.535
ω_7	E''	230	1.086
ω_8	E''	230	1.086
ω_9	E'	134	0.363
ω_{10}	E'	134	0.363
ω_{11}	A_1'	123	0.314
ω_{12}	A_2''	-557	2.367
zpe	24.1		

3 Cartesian Coordinates

Table S7: Cartesian coordinates of the geometry of the halogen-appended $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion complex optimised at MP2/AVQZ, in Å.

		x	y	z
C_{3v}	C	0.000000	0.000000	-0.817022
	Cl	0.000000	1.660488	-1.452849
	Cl	0.000000	0.000000	0.933545
	Cl	1.438025	-0.830244	-1.452849
	Cl	-1.438025	-0.830244	-1.452849
	Cl	0.000000	0.000000	3.713364

Table S8: Cartesian coordinates of the geometry of the $\text{Cl}^- \cdots \text{CCl}_4$ gas phase anion complex optimised at MP2/AVQZ, in Å, featuring the halide at the face of the CCl_4 molecule.

		x	y	z
C_{3v}	C	0.000000	0.000000	-0.668872
	Cl	0.000000	0.000000	-2.461208
	Cl	0.000000	1.659347	-0.113911
	Cl	1.437036	-0.829673	-0.113911
	Cl	-1.437036	-0.829673	-0.113911
	Cl	0.000000	0.000000	3.039013

Table S9: Cartesian coordinates of the geometry of the CCl_5^- transition state optimised at MP2/AVQZ, in Å.

		x	y	z
D_{3h}	C	0.000000	0.000000	0.000000
	Cl	0.000000	1.713893	0.000000
	Cl	-1.484275	-0.856946	0.000000
	Cl	0.000000	0.000000	2.430953
	Cl	1.484275	-0.856946	0.000000
	Cl	0.000000	0.000000	-2.430953