Short title: 2,4-D resistance cost

No Auxinic Herbicide Resistance Cost in Wild Radish (*Raphanus raphanistrum*)

Danica E. Goggin¹, Hugh J. Beckie², Chad Sayer³ and Stephen B. Powles⁴

¹Research Associate, Australian Herbicide Resistance Initiative, School of Agriculture and Environment, University of Western Australia, Perth, Australia; ²Professor, Australian Herbicide Resistance Initiative, School of Agriculture and Environment, University of Western Australia, Perth, Australia; ³Global Lead Technical Services, Nufarm Australia Limited, Laverton North, Victoria, Australia; ⁴Professor, Australian Herbicide Resistance Initiative, School of Agriculture and Environment, University of Western Australia, Perth, Australia

Author for correspondence: Danica E. Goggin, Research Associate (ORCID ID: 0000-0003-3856-9801), Australian Herbicide Resistance Initiative (M086), University of Western Australia, 35 Stirling Highway, Crawley 6009, Australia. (Email: danica.goggin@uwa.edu.au)

Abstract

Wild radish (*Raphanus raphanistrum* L.) is a problematic and economically damaging dicotyledonous weed infesting crops in many regions of the world. Resistance to the auxinic herbicides 2,4-D and dicamba is widespread in Western Australian *R. raphanistrum* populations, with the resistance mechanism appearing to involve alterations in the physiological response to synthetic auxins and in plant defence. This study aimed to determine if these alterations cause inhibition in plant growth or reproduction that could potentially be exploited to manage 2,4-D-resistant populations in cropping areas. Therefore, the morphology and seed production of resistant and susceptible populations were compared in an outdoor pot study, with plants grown in the presence and absence of competition by wheat. The susceptible and resistant *R. raphanistrum* populations were equally suppressed by wheat competition, with plant growth and seed production being decreased by approximately 50%. Although resistant populations produced less vegetative biomass than susceptible populations, there was no negative association between resistance and seed production. Therefore, it is unlikely that any non-herbicidal management practices will be more efficacious on 2,4-D-resistant than -susceptible *R. raphanistrum* populations.

Key words: 2,4-D, Brassicaceae, competition, dicamba, seed dormancy.

Introduction

Auxinic herbicides have been applied to crops such as wheat and barley for over 60 years, with usage levels in the major grain crops generally decreasing as newer herbicides were added to the market (Peterson et al. 2016). However, growers are once again becoming more reliant on auxinic herbicides to control weeds in cropping situations, due to the introduction of auxin-resistant transgenic crops and/or to the development of widespread weed resistance to other herbicide modes of action. Consequently, there is a risk that this high selection pressure could accelerate the evolution of auxinic herbicide resistance in weeds. This scenario has already been observed in Western Australian populations of wild radish (*Raphanus raphanistrum* L.), where almost universal resistance to the acetolactate synthase (ALS)-inhibiting herbicides has resulted in increased use of the synthetic auxins 2,4-dichlorophenoxyacetic acid (2,4-D) and 4-methyl-2-chlorophenoxyacetic acid (MCPA) and a concomitant increase in the number of 2,4-D-resistant populations (Owen et al. 2015). Research aimed at characterisation of the resistance mechanism(s) in a number of these populations has ruled out common non-target-site mechanisms such as metabolic detoxification, vacuolar sequestration and reduced herbicide uptake, and have pointed to altered synthetic auxin perception or signalling and/or enhanced defence at the plasma membrane (Goggin et al. 2018). Although many resistant populations exhibit restricted phloem movement of 2,4-D, the extent of this reduced translocation is not correlated with resistance level. Therefore, the precise molecular mechanism(s) of auxinic herbicide resistance in *R. raphanistrum* remains to be determined.

Arabidopsis [*Arabidopsis thaliana* (L.) Heynh.] plants with resistance to natural or synthetic auxins conferred by mutations in auxin transporters, receptors, signalling proteins or metabolic enzymes often display characteristic phenotypes such as dwarfism (e.g. Estelle and Somerville 1987; Woodward et al. 2007), lack of shoot apical dominance and inhibited

lateral root formation (e.g. Ruegger et al. 1997), twisting and epinasty (e.g. Geisler et al. 2003), altered leaf serration (e.g. Bilsborough et al. 2011; Kasprzewska et al. 2015) and delayed flowering (e.g. Tognetti et al. 2010), but this does not always lead to a reproductive cost (Roux and Reboud 2005). In general, the phenotypes of weeds with evolved auxin resistance are less differentiated from their susceptible counterparts, although a wild mustard (*Sinapis arvensis* L.) population with resistance to four auxinic herbicides (2,4-D, MCPA, dicamba and picloram) contained plants that were smaller, lacked apical dominance, and had darker green leaves that were more serrated than those of a susceptible population (Hall and Romano 1995; Mithila et al. 2012). Plants in a dicamba-resistant *Chenopodium album* L. population were also smaller in stature and less competitive, but in this case the leaves were lighter green and less serrated than those of the susceptible population (Rahman et al. 2014; Ghanizadeh and Harrington 2018).

Initial observations made during the generation of seed resources for the 2,4-D-resistant *R. raphanistrum* populations studied in Goggin et al. (2018) suggested possible differences between populations in terms of leaf size and serration, plant stature, and reproductive capacity. Therefore, the current study quantified parameters relating to seedling morphology, flowering and seed production (with and without competition from wheat) between two herbicide-susceptible and eleven 2,4-D-resistant populations that have been previously characterised at the physiological and biochemical level. The potential confounding effects of (1) the different genetic backgrounds of each population and (2) the presence of resistance to non-auxinic herbicides (Owen et al. 2015) are partially offset by the use of a large number of populations (Van Etten et al. 2016; Osipitan and Dille 2017). Although of different genetic backgrounds and (potentially) auxin resistance mechanisms, these populations represent the spectrum of auxin-resistant *R. raphanistrum* populations

present in Australian cropping fields. Thus, it is important to identify any weaknesses in growth or reproduction that could be used to develop non-herbicidal management strategies.

Materials and Methods

Plant Material. The two herbicide-susceptible (S1 and S2) and eleven resistant (R1 – R11) *R. raphanistrum* populations originally collected from across the Western Australian grain belt (Owen et al. 2015) and characterised in Goggin et al. (2018) were used for the current study. The field-collected populations had been screened for resistance to the recommended field rates of chlorsulfuron, imazamox+imazapyr (hereafter abbreviated as IMI), diflufenican, atrazine, glyphosate and bromoxynil+pyrasulfotole as well as to 2,4-D (Owen et al. 2015), and the observed levels of resistance to these herbicides were taken into account when examining potential effects of 2,4-D resistance on plant characteristics. None of the populations were resistant to atrazine, glyphosate or bromoxynil+pyrasulfotole (0% plant survival). Survival after chlorsulfuron treatment was >90% in all R populations, whereas resistance to diflufenican (0 – 30% survival) and IMI (0 – 50% survival) was more variable (Owen et al. 2015). In the study of Goggin et al. (2018), the original R populations were twice-selected at the recommended field rate (500 g a.e. ha⁻¹) of 2,4-D amine in an effort to increase the homogeneity of the 2,4-D resistance mechanism(s) in each population, and their level of resistance to 2,4-D and dicamba was quantified in dose-response experiments. The proportion of applied 2,4-D retained in the treated leaf was also quantified in Goggin et al. (2018) using ¹⁴C-labelled 2,4-D. A summary of the characteristics of each population is presented in Table 1.

All plants were grown in potting mix (composted pine bark:peat moss:washed river sand, 2:1:1) in pots of 250 mm diameter and 230 mm depth. To ensure uniformity, seeds

were pre-germinated on 0.6% (w/v) agar-water in the dark for 3 d at 20°C and then seedlings with radicles of similar length were transplanted into moist potting mix and grown outdoors at the University of Western Australia (31.99°S, 115.82°E) between May and October 2018 (average climatic conditions: maximum/minimum temperature, 20/11°C; day length, 11 h, 6 min; daily solar exposure 14 MJ m⁻²). Pots were arranged randomly on five adjacent 4 m² benches in full sun, watered as required, and fertilised weekly with 500 mL of 1.5 g L⁻¹ commercial soluble fertiliser (Diamond Red: N 27%, P 5.7%, K 10.9%, trace elements; Campbells Fertilisers, Laverton North, Australia). Aphids were controlled either by hand-picking or by spot spraying with commercial pyrethrum formulated for use in home gardens (Pyrethrum Long Life Insect Spray, 0.9 g L⁻¹ pyrethrum + 3 g L⁻¹ piperonyl butoxide; Multicrop, Scoresby, Australia).

Assessment of Seedling Growth and Morphology. For each population, there were five replicates of five seedlings per pot. When plants had reached the 3- to 4-leaf stage (11 days after transplant, DAT), the first leaf was removed from the biggest plant in each pot, pressed, and dried at 70°C for 48 h, at which time they were photographed, measured and weighed. Leaf area was measured by weighing cut-out photographs of the leaves and comparing to a standard curve of the masses of known areas of paper, allowing the area of the leaf photograph to be calculated from its mass. At 15 DAT, plant diameter and height were recorded, along with the number of leaves per plant. When floral buds were just visible in most pots (34 DAT), the above-ground plant parts were harvested and dried at 70°C for 72 h and then weighed.

Assessment of Plant Performance in the Presence and Absence of Competition. Five replicates of one plant per pot were set up for each population, either without competition,

or in competition with five wheat (*Triticum aestivum* L. ‘Mace’) seedlings per pot. This corresponds to a wheat density of 100 plants m⁻², which approximates commercial wheat crop densities. Wheat seeds were germinated on agar and transplanted into pots at the same time as the *R. raphanistrum* seeds. *Raphanus raphanistrum* plants were monitored daily during the experiment and the following parameters were recorded: date when floral buds first became visible; date of anthesis of first flower; height above soil of first flower; and flower colour. The latter was arbitrarily quantified by giving scores based on the range of colours observed in the study (Figure 1) and the fact that yellow (carotenoid production) is recessive (colour score: 0) whilst purple (anthocyanin production) is dominant (colour score: 7) in *R. sativus* (Irwin and Strauss 2005).

Wheat booting commenced at 68 DAT, heading at 74 DAT, and anthesis at 87 DAT. When the wheat had reached maturity, watering and fertilisation of all pots was ceased even if the *R. raphanistrum* plants were still actively growing and flowering. This protocol more closely reproduces conditions in an Australian cropping field where late-flowering plants might experience terminal drought. *Raphanus raphanistrum* siliques were collected as they matured to minimise losses through shedding (most plants started shedding siliques before the wheat ears had matured) and stored in a cool shed until harvest was completed. The date of final harvest for each *R. raphanistrum* plant was recorded, and the above-ground vegetative parts were collected and dried at 70°C for 48 h and then weighed. Once the harvest was complete, siliques were dried in a 30°C oven for 7 d, weighed, and then milled to obtain the seeds. Seeds were weighed and counted and their dormancy and viability assessed in a germination test (see below). Wheat ears were counted and weighed, threshed, and the seeds were counted and weighed.

Seed Germination Tests. Fifty seeds from each *R. raphanistrum* plant were placed on 0.6% (w/v) agar-water and incubated in the dark at 20°C. Seed germination was counted every 3 – 4 days for 14 d, and the germinated seedlings removed. At 14 d, ungerminated seeds were transferred to agar containing 50 µM each of gibberellin A₃ (GA₃) and fluridone, a combination which is known to break dormancy in *R. raphanistrum* seeds (Goggin and Powles 2014), and placed in a growth cabinet at 20/15°C day/night with a 12 h photoperiod of white incandescent and LED light at 200 µmol m⁻² s⁻¹. Germination was counted every 3 – 4 days for another 17 d, and the ungerminated seeds tested for viability by cutting through the embryo and assessing its firmness and colour.

Statistical Analysis. Whilst keeping in mind that the auxinic herbicide resistance mechanism(s) in the populations studied is not fully characterised, the fitness of each population was calculated using the equation cited in Vila-Aiub et al. (2015), where fitness (W) is the product of the proportion of plants surviving after seed dispersal (P) and the number of offspring produced (N). For this study, seed viability was used as a measure of P, and the total number of seeds produced as N. Differences between populations and treatments were assessed by one- and two-factor analysis of variance and the Fisher's least significant difference (LSD) test at a significance level of 5%, as well as by principal component analysis (PCA) using MetaboAnalyst 4.0 (Xia et al. 2009) with auto-scaling used for data normalisation. As the data were normally distributed according to a quantile-quantile plot (Wilk and Gnanadesikan 1968), but heteroscedastic according to a Bartlett's test (McDonald 2014), weighted least squares regression was used to identify possible links between population characteristics and their resistance to 2,4-D and dicamba or ability to translocate 2,4-D out of the treated leaf (all quantified in Goggin et al. 2018). Weighted least squares regression was performed using the Real Statistics Resource Pack software (release

6.2) in Excel (Zaiontz 2019), and correlation coefficients were considered to be significant if $P < 0.05$. Resistance was expressed as both “survival”, based on the herbicide dose required to kill 50% of the plants in the population (LD_{50}) and as “growth”, based on vegetative biomass of survivors following auxinic herbicide treatment. To account for possible confounding effects due to the resistance of the R populations to other herbicides, the correlation between the various parameters measured in this study and the level of resistance (in this case expressed as the percentage survival of a population sprayed with the recommended rate of each herbicide, from Owen et al. 2015) was also calculated. All significant correlations were plotted to check for outliers distorting the correlation coefficient.

Results and Discussion

Overall, there were no striking differences among populations in the parameters measured (Supplementary Table S1), despite their geographically distinct origins. As was observed for the biochemical characteristics assessed in Goggin et al. (2018), a distinguishing feature of these *R. raphanistrum* populations is their wide intra-population variability that often precludes identification of statistically significant inter-population differences. The PCA showed separation between the S and R populations when resistance parameters (LD_{50} values and post-spray biomass for 2,4-D and dicamba) were included in the analysis, but not when only plant growth and seed production parameters were used (Figure 2).

Growth and Morphology of Young Plants. The first leaf of population R8 was around 20% larger than that of most other populations, whilst the first leaf of R5 was 20% smaller (Supplementary Table S1). Populations R6 and R8 had a 12% larger seedling diameter than

most others, R1 had an average of 0.5 extra leaves at 15 DAT, and S2 had 20% higher biomass than most other populations at 34 DAT whilst that of R5 was around 25% lower (Supplementary Table S1).

Seedling diameter and the dry mass and area of the first leaf were positively correlated with vegetative growth following dicamba treatment, whilst there was a negative relationship between specific leaf area and resistance to 2,4-D (post-spray growth) and dicamba (survival and post-spray growth) (Table 2). Seedling height and dry mass were negatively correlated with IMI resistance, and the same relationship was observed between seedling height and growth after 2,4-D treatment. First-leaf serration was not quantified, but visual examination of leaf photographs shows a level of variability within populations that was similar to the inter-population variation, and no consistent differences between the S and R populations (Figure 3).

Although the dicamba-resistant seedlings tended to be of larger diameter and leaf area, the difference between the largest resistant populations and the smallest susceptible population was only around 1.3-fold, which may be difficult for a grower to visually identify before spraying a field with an auxinic herbicide. However, the almost 2-fold difference in height between the most IMI-resistant and the susceptible populations could represent an opportunity for growers to identify fields with high numbers of IMI-resistant seedlings and use an alternative weed control method with a higher likelihood of success.

Performance of *Raphanus raphanistrum* Populations in the Absence of Competition.

Bud formation and anthesis were 2 – 3 weeks later in population R2 than in most others, and occurred at least 2 weeks earlier in R1 (Supplementary Table S1). The first flower of population S2 was positioned at least 15 cm higher above the soil than the other populations, whilst those of R5 were around 14 cm lower (Supplementary Table S1). Vegetative shoot

dry mass at harvest was around 30% higher in S1 and S2 and 40% lower in R3, and silique dry mass and seed number were close to 40% higher in R9 and 65% lower in R2 (Supplementary Table S1). Mass of 100 seeds was 25% higher in R1 (Supplementary Table S1). Survival and growth after both 2,4-D and dicamba treatment were strongly negatively correlated with vegetative biomass at harvest (Table 2). There was also a positive correlation between the ratio of silique:vegetative biomass and growth following dicamba treatment (Table 2).

The consistent association between auxinic herbicide resistance and lower accumulation of above-ground vegetative biomass suggests that the resistance mechanism(s) in *R. raphanistrum* causes the plants to divert resources from growth into other processes (reviewed in Vila-Aiub et al. 2015); for example, constitutive activation of plant defence pathways (Goggin et al. 2018). However, this potential resistance cost did not extend to seed production, and indeed the populations showing the strongest vegetative growth following dicamba treatment also invested the most in silique production.

Performance of *Raphanus raphanistrum* Populations Under Competition with Wheat.

Overall, the presence or absence of wheat competition did not cause a separation of *R. raphanistrum* populations in the PCA (Figure 2). Competition did not significantly affect certain parameters (e.g. timing of bud formation and anthesis, relative partitioning between vegetative and reproductive mass; mass of 100 seeds), whilst others (e.g. plant biomass, silique mass, seed number) were decreased by approximately 50% (Table 3). Additionally, the average height of the first flowers was around 25% greater when plants were grown with wheat, and the final harvest occurred 6 d earlier (Table 3). There were no significant differences between pots in terms of number of wheat ears produced (overall mean: 15 ± 0.2

ears per pot), total ear dry mass (40 ± 1 g), total wheat seed mass (31 ± 1 g) or total wheat seed number (605 ± 15).

When *R. raphanistrum* populations were analysed individually, there were few differences among them in terms of time of bud formation, anthesis, date of final harvest or reproductive performance (Supplementary Table S1). The first flowers of population S2 were around 15 cm higher and S2 also had a 50% greater vegetative biomass at harvest, and R1 and R5 had first flowers that were at least 15 cm shorter than most other populations (Supplementary Table S1). Populations with earlier bud formation tended to show better growth following dicamba treatment, and vegetative biomass at harvest was negatively correlated with dicamba survival and post-treatment growth, reflecting the result observed with plants grown in the absence of competition (Table 2). The fact that there was not also a relationship between 2,4-D resistance and biomass at harvest in the presence of wheat suggests that the overall suppression of biomass production caused by competition masks the observed growth penalty inflicted by 2,4-D resistance in the absence of competition. As the relationship between dicamba resistance and plant biomass was expressed in both the presence and absence of competition, it is possible that dicamba resistance imposes a greater cost on vegetative growth than does 2,4-D resistance, but a firm conclusion cannot be drawn until the precise resistance mechanisms are known.

Previous work has shown that a combination of 2,4-D treatment and competition from wheat at ≥ 100 plants m^{-2} was successful in controlling one 2,4-D-resistant *R. raphanistrum* population, but not another (Walsh et al. 2009). The surviving population from that study corresponds to the population here designated as R1, which exhibits the lowest level of 2,4-D resistance of the 11 R populations tested (Goggin et al. 2018). Therefore, although useful in suppressing *R. raphanistrum* in general, as also observed by Eslami et al.

(2006), crop competition alone is not likely to be more effective on auxin-resistant vs. -susceptible populations.

Seed Dormancy and Viability. Overall, *R. raphanistrum* seed viability and responsiveness to germination stimulants was not affected by wheat competition during seed production, but the level of primary dormancy was slightly higher (i.e. lower germination at 14 d after the start of imbibition, just before the seeds were transferred to germination stimulants) under competitive conditions, and fitness was decreased by more than 50% (Table 3). Between populations, the viability of seed produced by S2 and R9 grown without competition was lower than that of the other populations, but seed viability was >90% in all populations (Supplementary Table S1). In spite of its slightly lower seed viability, the calculated fitness of population R9 in the absence of competition was at least 80% higher than most of the others due to its high seed production. Under competitive conditions, however, there were no differences between populations in terms of fitness (Supplementary Table S1). Primary dormancy was highest in populations R3, R6 and R9 under both competitive and non-competitive conditions, whilst R7 seeds were the least dormant (Supplementary Table S1). Population R7 was also the most responsive to germination stimulants, with 50% germination occurring at least 5 days earlier than in most other populations (Supplementary Table S1). Under competitive conditions, there was a moderate negative correlation between vegetative growth after 2,4-D treatment and germination at 14 d (Table 2).

Based on the slightly higher dormancy of seeds produced by the R populations, it could be beneficial to delay sowing of the next season's crop in a field with a high number of 2,4-D survivors so that the next generation of *R. raphanistrum* can be controlled with non-selective, pre-sowing herbicides. Alternatively, a highly competitive crop or pasture

could be sown early, to suppress the late-emerging auxin-resistant *R. raphanistrum* seedlings. A similar situation exists in glyphosate-resistant (Beckie et al. 2018) and auxin-resistant (Kumar and Jha 2016) *Kochia scoparia*, in which the slower germination of the resistant populations could potentially be exploited using the same strategies.

Other Considerations. Populations which retained a higher proportion of applied 2,4-D in the treated leaf tended to have a higher number of leaves at 15 DAT, a higher ratio of seed:silique biomass (no competition), a higher silique biomass (under competition), and, interestingly, flowers that were more likely to be purple (Table 2). High concentrations of auxin have been shown to initiate leaf formation in seedlings (Bar and Ori 2014), stimulate fruit production (Figueiredo and Köhler 2018) and inhibit biosynthesis of anthocyanins (Liu et al. 2014) and carotenoids (Liu et al. 2015). Therefore, the alteration in auxin transport that seems to have developed under repeated selection with 2,4-D may have minor effects on leaf formation, resource allocation and floral pigment production by affecting local auxin concentrations, although this does not have a significant impact on plant performance. These correlations are, however, based on population-level characteristics, and experiments in which translocation or resource allocation are assessed in individual seedlings with known flower colour have not yet been performed.

Management of auxin-resistant *Raphanus raphanistrum*. The lack of an obvious resistance cost in 2,4-D- and dicamba-resistant *R. raphanistrum* is similar to the situation observed in several other auxin-resistant weeds, such as *Conyza canadensis* (L.) Cronquist (Kruger et al. 2010), *Carduus nutans* L. (Bonner et al. 1998), *Centaurea solstitialis* L. (Sterling et al. 2001) and *Kochia scoparia* (L.) Schrad. (Menalled and Smith 2007) (although it must be noted that a different *K. scoparia* population, resistant to both dicamba and

fluroxypyr, did exhibit a significant resistance cost: Kumar and Jha (2016)). In *Ranunculus acris* L., the fitness penalty identified in the resistant population was too slight to make a practical difference (Bourdote et al. 1996).

There also seems to be little opportunity to exploit any resistance costs in 2,4-D-resistant *R. raphanistrum* populations, apart from the slightly higher seed dormancy described above. In spite of the lower biomass of resistant populations, there was no impairment in seed production, so it is expected that 2,4-D-resistant individuals of *R. raphanistrum* found across the Western Australian grain belt would not decline in frequency relative to susceptible individuals in populations if use of 2,4-D was discontinued. Therefore, effective practices already in place, such as rotation and mixing of still-effective herbicides (including synthetic auxins at the full recommended dose), harvest weed seed control and high crop sowing rates, need to be rigorously maintained to mitigate the further evolution and spread of auxin-resistant weeds. It is also worth exploring the observed reduction in seedling height that is associated with resistance to imidazolinones, by studying populations that are resistant to this mode of action but still susceptible to auxinic herbicides. If the relationship is found to be reliable, growers may be able to readily identify fields with potential imidazolinone resistance problems and avoid the use of these herbicides on plants that will not be adequately controlled.

Acknowledgements

This research was funded by the Australian Research Council (project no. LP150100161). Chad Sayer is an employee of Nufarm Australia, which manufactures and sells auxinic herbicide products. We thank Paula Reeve for threshing the wheat and wild radish seeds generated in the study, Mechelle Owen for providing the original wild radish populations

362 and accompanying information, and the two anonymous reviewers whose suggestions
363 helped to improve the manuscript.

364

365

References

- Bar M, Ori N (2014) Leaf development and morphogenesis. *Development* 141:4219-4230.
- Beckie HJ, Blackshaw RE, Leeson JY, Stahlman PW, Gaines TA, Johnson EN (2018) Seedbank persistence, germination and early growth of glyphosate-resistant *Kochia scoparia*. *Weed Res* 58:177-187.
- Bilsborough GD, Runions A, Barkoulas M, Jenkins HW, Hasson A, Galinha C, Laufs P, Hay A, Prusinkiewicz P, Tsiantis M (2011) Model for the regulation of *Arabidopsis thaliana* leaf margin development. *Proc Natl Acad Sci USA* 108:3424-3429.
- Bonner KI, Rahman A, James TK, Nicholson KS, Wardle DA (1998) Relative intra-species competitive ability of nodding thistle biotypes with varying resistance to the herbicide 2,4-D. *NZ J Agric Res* 41:291-297.
- Bourdote GW, Saville DJ, Hurrell GA (1996) Ecological fitness and the decline of resistance to the herbicide MCPA in a population of *Ranunculus acris*. *J Appl Ecol* 33:151-160.
- Eslami SV, Gill GS, Bellotti B, McDonald G (2006) Wild radish (*Raphanus raphanistrum*) interference in wheat. *Weed Sci* 54:749-756.
- Estelle MA, Somerville C (1987) Auxin-resistant mutants of *Arabidopsis thaliana* with an altered morphology. *Mol Gen Genet* 206:200-206.
- Figueiredo DD, Köhler C (2018) Auxin: a molecular trigger of seed development. *Genes Dev* 32:479-490.
- Geisler M, Kolukisaoglu HU, Bouchard R, Billion K, Berger J, Saal B, Frangne N, Koncz-Kálmán Z, Koncz, Dudler R, Blakeslee JJ, Murphy AS, Martinoia E, Schulz B (2003) TWISTED DWARF1, a unique plasma membrane-anchored immunophilin-

- 390 like protein, interacts with *Arabidopsis* multidrug resistance-like transporters
391 AtPGP1 and AtPGP19. Mol Biol Cell 14:4238-4249.
- 392 Ghanizadeh H, Harrington KC (2018) Fitness costs associated with multiple resistance to
393 dicamba and atrazine in *Chenopodium album*. Planta 249:787-797.
- 394 Goggin DE, Kaur P, Owen MJ, Powles SB (2018) 2,4-D and dicamba resistance
395 mechanisms in wild radish: subtle, complex and population-specific? Ann Bot
396 122:627-640.
- 397 Goggin DE, Powles SB (2014) Fluridone: a combination germination stimulant and
398 herbicide for problem fields? Pest Manag Sci 70:1418-1424.
- 399 Hall JC, Romano ML (1995) Morphological and physiological differences between the
400 auxinic herbicide-susceptible (S) and -resistant (R) wild mustard (*Sinapis arvensis*
401 L.) biotypes. Pestic Biochem Physiol 52:149-155.
- 402 Irwin RE, Strauss SY (2005) Flower color microevolution in wild radish: evolutionary
403 response to pollinator-mediated selection. Am Nat 165:225-237.
- 404 Kasprzewska A, Carter R, Swarup R, Bennett M, Monk N, Hobbs JK, Fleming A (2015)
405 Auxin influx importers modulate serration along the leaf margin. Plant J 83:705-718.
- 406 Kruger GR, Davis VM, Weller SC, Johnson WG (2010) Growth and seed production of
407 horseweed (*Conyza canadensis*) populations after exposure to postemergence 2,4-D.
408 Weed Sci 58:413-419.
- 409 Kumar V, Jha P (2016) Differences in germination, growth, and fecundity characteristics of
410 dicamba-fluroxypyr-resistant and susceptible *Kochia scoparia*. PLoS One
411 11:e0161533.
- 412 Liu L, Shao Z, Zhang M, Wang Q (2015) Regulation of carotenoid metabolism in tomato.
413 Mol Plant 8:28-39.

- 414 Liu Z, Shi M-Z, Xie D-Y (2014) Regulation of anthocyanin biosynthesis in *Arabidopsis*
 415 *thaliana* red *pap1-D* cells metabolically programmed by auxins. *Planta* 239:765-781.
- 416 McDonald JH (2014) Handbook of Biological Statistics. 3rd ed. Baltimore, MD: Sparky
 417 House Publishing. Pp 137-139.
- 418 Menalled FD, Smith RG (2007) Competitiveness of herbicide-resistant and herbicide-
 419 susceptible kochia (*Kochia scoparia* [L.] Schrad.) under contrasting management
 420 practises. *Weed Biol Manage* 7:115-119.
- 421 Mithila J, McLean MD, Chen S, Hall JC (2012) Development of near-isogenic lines and
 422 identification of markers linked to auxinic herbicide resistance in wild mustard
 423 (*Sinapis arvensis* L.). *Pest Manag Sci* 68:548-556.
- 424 Osipitan OA, Dille JA (2017) Fitness outcomes related to glyphosate resistance in Kochia
 425 (*Kochia scoparia*): what life history stage to examine? *Front Plant Sci* 8:1090.
- 426 Owen MJ, Martinez NJ, Powles SB (2015) Multiple herbicide-resistant wild radish
 427 (*Raphanus raphanistrum*) populations dominate Western Australian cropping fields.
 428 *Crop Pasture Sci* 66:1079-1085.
- 429 Peterson MA, McMasters SA, Riechers DE, Skelton J, Stahlmann PW (2016) 2,4-D past,
 430 present and future: a review. *Weed Technol* 30:303-345.
- 431 Rahman A, James T, Trollove M (2014) Characteristics and control of dicamba-resistant
 432 common lambsquarters (*Chenopodium album*). *Weed Biol Manage* 14:88-98.
- 433 Roux F, Reboud X (2005) Is the cost of herbicide resistance expressed in the breakdown of
 434 the relationships between characters? A case study using synthetic-auxin-resistant
 435 *Arabidopsis thaliana* mutants. *Genet Res* 85:101-110.
- 436 Ruegger M, Dewey E, Hobbie L, Brown D, Bernasconi P, Turner J, Muday G, Estelle M
 437 (1997) Reduced naphthylphthalamic acid binding in the *tir3* mutant of *Arabidopsis*

- 438 is associated with a reduction in polar auxin transport and diverse morphological
439 defects. *Plant Cell* 9:745-757.
- 440 Sterling TM, Lownds NK, Murray LW (2001) Picloram-resistant and -susceptible yellow
441 starthistle accessions have similar competitive ability. *Weed Sci* 49:42-47.
- 442 Tognetti VB, Van Aken O, Morreel K, Vandenbroucke K, van de Cotte B, De Clercq I,
443 Chiwocha S, Fenske R, Prinsen E, Boerjan W, Genty B, Stubbs KA, Inzé D, Van
444 Breusegem F (2010) Perturbation of indole-3-butyric acid homeostasis by the UDP-
445 glucosyltransferase UGT74E2 modulates *Arabidopsis* architecture and water stress
446 tolerance. *Plant Cell* 22:2660-2679.
- 447 Van Etten ML, Kuester A, Chang S-M, Baucom RS (2016) Fitness cost of herbicide
448 resistance across natural populations of the common morning glory, *Ipomoea*
449 *purpurea*. *Evolution* 70:2199-2210.
- 450 Vila-Aiub MM, Gundel PE, Preston C (2015) Experimental methods for estimation of plant
451 fitness costs associated with herbicide-resistance genes. *Weed Sci.* 63:203-216.
- 452 Walsh MJ, Maguire N, Powles SB (2009) Combined effects of wheat competition and 2,4-
453 D amine on phenoxy herbicide resistant *Raphanus raphanistrum* populations. *Weed*
454 *Res* 49:316-325.
- 455 Wilk MB, Gnanadesikan R (1968) Probability plotting methods for the analysis of data.
456 *Biometrika* 55:1-17.
- 457 Woodward AW, Ratzel SE, Woodward EE, Shamoo Y, Bartel B (2007) Mutation of *E1-*
458 *CONJUGATING ENZYME-RELATED1* decreases RELATED TO UBIQUITIN
459 conjugation and alters auxin response and development. *Plant Physiol* 144:976-987.
- 460 Xia J, Psychogios N, Young N, Wishart DS (2009) MetaboAnalyst: a web server for
461 metabolomic data analysis and interpretation. *Nucleic Acids Res* 37:W652-W660.

462 Zaiontz C (2019) Real Statistics Using Excel. <http://www.real-statistics.com>. Accessed:
463 June 11, 2019.

464

Table 1. Summary of herbicide resistance characteristics of the *Raphanus raphanistrum* populations. Rainfall zones (H, 450 – 750 mm; M, 325 – 450 mm; L, <325 mm) from which the populations were originally collected are given in parentheses after the population name. The resistance indices for 2,4-D and dicamba were expressed as the ratio of the LD₅₀ value (survival) or post-spray biomass (growth) of each R population and the mean of the S1 and S2 populations (see Goggin et al. 2018 for original data). Resistance to chlorsulfuron (Chlor), imazamox+imazapyr (IMI) and diflufenican (DFF) was classified as S (0% plant survival at the recommended field rate); r (1 – 19%); or R ($\geq 20\%$) (Owen et al. 2015).

Population	Resistance index (R:S)				Resistance classification		
	2,4-D	2,4-D	Dicamba	Dicamba	Chlor	IMI	DFF
	survival	growth	survival	growth			
S1 (M)	1	1	1	1	S	S	S
S2 (H)	1	1	1	1	S	S	S
R1 (M)	7	19	2	2	R	S	R
R2 (H)	42	37	4	10	R	r	S
R3 (M)	43	62	8	41	R	r	r
R4 (H)	43	18	4	36	R	r	r
R5 (M)	48	74	12	9	R	R	r
R6 (M)	27	65	27	23	R	S	S
R7 (M)	18	15	6	6	R	S	r
R8 (L)	27	41	11	16	R	r	S
R9 (L)	35	41	10	7	R	R	r
R10 (L)	21	20	9	7	R	R	S
R11 (L)	17	41	15	41	R	r	S

Table 2. Significant ($P<0.05$) associations between herbicide resistance parameters and plant characteristics as calculated by weighted least squares regression. The data used to generate the correlation coefficients are presented in Supplementary Table S1. The full correlation matrix is in Supplementary Table S2. DAT, days after transplant; IMI, imazamox+imazapyr.

Parameter 1	Parameter 2	Correlation coefficient
2,4-D survival	Plant dry mass (-wheat)	-0.747
2,4-D growth	Seedling height	-0.557
	Specific leaf area of first leaf	-0.631
	Plant dry mass (-wheat)	-0.891
	Germination at 14 d (+wheat)	-0.657
Dicamba survival	Specific leaf area of first leaf	-0.639
	Plant dry mass (-wheat)	-0.638
	Plant dry mass (+wheat)	-0.624
Dicamba growth	Seedling diameter	0.578
	Dry mass of first leaf	0.764
	Area of first leaf	0.827
	Specific leaf area of first leaf	-0.604
	Time of first flower buds (+wheat)	-0.572
	Plant dry mass (-wheat)	-0.782
	Plant dry mass (+wheat)	-0.727
	Ratio silique:vegetative dwt (-wheat)	0.618
2,4-D retained in treated leaf	No. leaves at 15 DAT	0.591

	Flower colour	0.597
	Silique dry mass (+wheat)	-0.554
	Ratio seed:silique dwt (-wheat)	0.658
IMI resistance	Seedling height	-0.689
	Seedling dry mass	-0.571

480

481

Table 3. Effect of wheat competition on *Raphanus raphanistrum* growth and reproduction.

The data for all 13 populations were pooled. Significant differences between competition treatments ($P < 0.05$) are denoted by an asterisk in the third column. DAT, days after transplant.

Parameter	<i>Raphanus raphanistrum</i> grown alone	<i>Raphanus raphanistrum</i> grown with wheat
Bud formation (DAT)	39.3 ± 2.2	39.8 ± 2.8
Height of first flower (cm)	27.8 ± 3.1	$38.1 \pm 2.5^*$
First flower anthesis (DAT)	58.1 ± 3.7	57.7 ± 2.8
Final plant dry mass (g)	21.8 ± 1.7	$9.0 \pm 1.0^*$
Time of final harvest (DAT)	153 ± 1	$147 \pm 1^*$
Silique dry mass (g)	21.1 ± 1.9	$9.1 \pm 0.8^*$
100-seed mass (g)	0.58 ± 0.03	0.59 ± 0.02
Total no. seeds	698 ± 70	$321 \pm 24^*$
Ratio silique:vegetative mass	1.1 ± 0.1	1.1 ± 0.1
Ratio seed:silique mass	0.20 ± 0.01	0.21 ± 0.01
Seed viability (%)	97.2 ± 0.7	97.6 ± 0.4
Germination at 14 d (%)	25.8 ± 2.8	$19.8 \pm 2.2^*$
Time to 50% germination (d)	19.0 ± 1.0	19.6 ± 0.7
Fitness	669 ± 70	$294 \pm 54^*$

Figure Legends

Figure 1. Variation in *Raphanus raphanistrum* flower colour, with arbitrary scores assigned as follows: (A) yellow: 0; (B) pale yellow/cream: 1; (C) white: 2; (D) yellow with purple edges: 3; (E) white with purple edges: 4; (F) purple with white centre: 5; (G) pale purple: 6; (H) purple: 7.

Figure 2. Principal component analysis of growth and seed production of 2,4-D-susceptible (S) and -resistant (R) *Raphanus raphanistrum* populations grown in the presence (+ wheat) and absence (- wheat) of competition, with herbicide resistance parameters included (A) or excluded (B) from the analysis. Shaded areas represent the 95% confidence intervals of each group. The first two of eight components in total are shown.

Figure 3. Morphology of first-leaves of five seedlings from each population at 11 days after transplant. Leaves were pressed and dried at 70°C before they were photographed.

505

506

507

508 Figure 1.

509

510

511

512

513 Figure 2.

514

515

516

517 Figure 3.