Belowground-mediated and phase-dependent processes drive nitrogen-evoked community dynamic changes in grasslands

Journal:	<i>Journal of Ecology</i>
Manuscript ID	Draft
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Tian, Qiuying; Institute of Botany Chinese Academy of Sciences Yang, Liuyi; Institute of Botany Chinese Academy of Sciences Ma, Pengfei; State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences Liu, Nana ; Peking University Bai, Wen Ming; the Chinese Academy of Sciences, Institute of Botany Wang, Hong; Hebei Agricultural University Ren, Lifei; Institute of Botany, CAS Lu, Peng; Institute of Botany Chinese Academy of Sciences Han, Wenwu; Institute of Botany, Photosynthesis Research Centre Schultz, Peggy; University of Kansas Bever, James; University of Kansas, Ecology & Evolutionary Biology Zhang, Fusuo ; China Agricultural University Lambers, Hans; Western Australia University, Faculty of Agriculture Zhang, Wen-Hao; Institute of Botany, Photosynthesis Research Centre;
Key-words:	nitrogen enrichment, plant biodiversity, grassland ecosystems, soil acidification, metal toxicity, belowground processes, functional groups, forbs and grasses, plant-soil interaction

SCHOLARONE™
Manuscripts

1 **Belowground-mediated and phase-dependent processes drive nitrogen-evoked**
2 **community dynamic changes in grasslands**
3
4 Qiuying Tian¹, Liuyi Yang^{1,2}, Pengfei Ma^{1,2}, Nana Liu³, Wenming Bai¹, Hong Wang⁴,
5 Lifei Ren¹, Peng Lu¹, Wenwu Han¹, Peggy A. Schultz⁵, James D. Bever⁵, Fu-Suo
6 Zhang⁶, Hans Lambers^{6,7}, Wen-Hao Zhang^{1*}
7 ¹State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,
8 Chinese Academy of Sciences, Beijing 100093, China.
9 ²University of Chinese Academy of Sciences, Beijing 100049, China.
10 ³Institute of Ecology and Key Laboratory for Earth Surface Processes of the Ministry
11 of Education, College of Urban and Environmental Sciences, Peking University,
12 Beijing 100871, China.
13 ⁴Mountain Area Research Institute, Agricultural University of Hebei, Baoding 071001,
14 China.
15 ⁵Department of Ecology and Evolutionary Biology, and the Kansas Biological Survey,
16 University of Kansas, Lawrence, KS 66047. USA
17 ⁶College of Resources and Environmental Sciences, Key Laboratory of Plant–Soil
18 Interactions of the Ministry of Education, China Agricultural University, Beijing
19 100193, China
20 ⁷School of Biological Sciences, University of Western Australia, Perth, WA 6009,
21 Australia
22 **Running title:** Nitrogen enrichment and grassland community

Summary

1. Enhanced deposition of atmospheric nitrogen (N) has greatly influenced terrestrial ecosystems across the globe. Loss of plant diversity by N enrichment has been widely reported in different types of grasslands. Although extensive studies have been conducted to evaluate the effects of N enrichments on plant biodiversity, few studies have linked time sequences of N enrichment to processes in soils and plants in grassland ecosystems worldwide.
2. We investigated the effects of N addition on plant diversity of temperate grasslands by monitoring dynamic changes in plant community, soil properties and physiological processes of dominant species in field N-addition experiments (up to 15 years of N addition) with multiple sites across northern China, and combined with greenhouse experiments.
3. We found that the N enrichment-induced declines in plant diversity of grasslands were phase-based, functional group-dependent, and driven by multiple belowground processes. The loss of plant biodiversity was broadly classified into three phases, and driven by different processes in belowground.
4. Nitrogen addition led to a rapid accumulation of inorganic N, particularly ammonium in soil, which rendered the ammonium-sensitive leguminous and broad-leaf-non-rhizomatous forbs toxicity, driving their loss in the community during early phase of N enrichment. The N-induced changes in this phase were independent of soil pH as evidenced by results from application lime to mitigate N-evoked soil acidification. With progression of N addition soil pH reduced, and the

soil acidification-induced mobilization metal ions in soil, particularly manganese (Mn^{2+}), caused excessive foliar accumulation of Mn in the narrow-leaf-non-rhizomatous forbs, contributing to their loss due to Mn toxicity in the second phase of N enrichment. When N addition proceeded longer than ~10 years, a greater allocation of photoassimilate to rhizomes stimulated rhizome growth and meristem differentiation of rhizomatous species in belowground, contributing to the dominance by rhizomatous sedges/grasses in the community at later phase of N enrichment.

5. Synthesis: The phase-based, functional group-specific and belowground-driven mechanism provides a comprehensive explanation for the decline in plant diversity by N enrichment in grassland ecosystems, and can be extended to explain plant diversity in other terrestrial ecosystems suffering from the enhanced N input.

Keywords: nitrogen enrichment, plant biodiversity, grassland ecosystems, soil acidification, metal toxicity, belowground processes, functional groups, forbs and grasses

Introduction

Enhanced atmospheric nitrogen (N) deposition due to human activities is altering N cycling, productivity, plant composition and biodiversity of terrestrial ecosystems globally (Vitousek *et al.* 1997; Clark *et al.* 2007; Galloway *et al.* 2008; Bobbink *et al.* 2010; Duprè *et al.* 2010; Kimball, Goulden, Suding, & Parker 2014). Nitrogen enrichment enhances plant productivity, but negatively affects species diversity in many N-limited terrestrial ecosystems (Stevens, Dise, Mountford, & Gowing 2004; Suding *et al.* 2005; Clark & Tilman 2008; LeBauer & Treseder 2008; Reich 2009), leading to reductions in ecosystem services (Cardinale *et al.* 2012; Zuppinger-Dingley *et al.* 2014; Isbell *et al.* 2015). Increased competition for aboveground resources has been suggested to be a key driver for the N-enrichment-induced reduction in species diversity (Grime 1973; Tilman 1987; Hautier, Niklaus, & Hector 2009; Borer *et al.* 2014; Farrer & Suding 2016). In addition, N deposition often results in limitation by other mineral nutrients, in particular phosphorus (P), for plant growth (Phoenix *et al.* 2004; Menge & Field 2007; Thayer, St Clair, Field, & Somerville 2008; Li, Niu, & Yu 2016). The environmental variables in soils have also been considered as the primary controls on plant diversity (Houdijk, Verbeek, Vandijk, & Roelofs 1993; Laliberté, Zemunik, & Turner 2014). For instance, N enrichment acidifies soils (Crawley *et al.* 2005; Horswill, O'sullivan, Phoenix, & Leake 2008; Tian & Niu 2015), thus threatens plant growth by altering availabilities of mineral nutrients in soils (Bowman, Cleveland, Halada, Hresko, & Baron 2008; Tian, Liu, Bai, Li, & Zhang 2015) and reduces niche dimensionality for species coexisting by decreasing spatial heterogeneity of resources

(Levin 1974; Tilman 1993; Stein, Gerstner, & Kreft 2014).

Although the above hypotheses provide potential explanations for the widely observed N-deposition-driven species loss, the relative importance and the temporal and spatial links among these mechanisms are generally unknown (Dickson & Foster 2011; Baer, Blair, & Collins 2015). According to the competition theory, elevated N input favors grass over forb growth, thus allowing grasses to outcompete forbs (Grime 1973; Tilman 1987; Hautier, Niklaus, & Hector 2009; Borer *et al.* 2014). However, these studies have rarely taken into account the temporal features of N effects on the community. The involvements of ammonium toxicity in the decline in N-induced plant diversity have been implicated in the literature (van den Berg *et al.* 2005; Stevens, Dise, Gowing, & Mountford 2006; Kleijn, Bekker, Bobbink, de Graaf, & Roelofs 2008). In contrast to this hypothesis, some of the lost plant species under N enrichment are insensitive to NH_4^+ concentrations (Bobbink, Ashmore, Braun, Flückiger, & van den Wyngaert 2003). Despite wide speculation on the ammonium toxicity in decline of plant biodiversity, the conclusion is mainly based on results from field survey and statistical analyses, while direct experimental evidence seems insufficient to verify this hypothesis. These results imply that multiple mechanisms may be involved in N enrichment-induced decline in plant diversity. The relative contributions of these mechanisms may depend on the duration and intensity of N deposition, and the differences in edaphic and community traits of grasslands. However, few studies have comprehensively evaluated the roles of these factors by the temporal approach yet.

106

107 The decline in plant diversity by N enrichment has mainly been observed in studies by
108 monitoring changes in plant community traits including aboveground biomass and
109 species richness from field surveys and *in situ* simulation experiments (Grime 1973;
110 Stevens, Dise, Mountford, & Gowing 2004; Suding *et al.* 2005; Clark *et al.* 2007; Clark
111 & Tilman 2008). However, the N-induced changes in these traits are relatively slow
112 (Larsen, Chase, Durance, & Ormerod 2018), and are consequences of altered
113 physiological processes in plants. Therefore, understanding physiological responses
114 underlying the changes in aboveground biomass and species richness can shed
115 important light on the N-driven decline in plant diversity.

116

117 Emerging evidence shows that N enrichment not only influences aboveground
118 processes, but it can also have a profound impact on underground processes, including
119 soil microbiomes, rhizosphere processes and root traits in grassland ecosystems (Bai *et*
120 *al.* 2015; Leff *et al.* 2015). However, few studies have focused on the underground
121 processes as well as the interactions between aboveground and underground processes
122 in N enrichment-induced changes in plant diversity. The spatial links and feedback
123 between the aboveground and underground processes can influence plant growth
124 (Mariotte *et al.* 2018). Therefore, research on interactions between plants and soils is
125 critical for our mechanistic understanding of vegetation changes driven by enhanced
126 atmospheric N deposition. Our previous studies have demonstrated that both soil- and
127 root-mediated processes are involved in N enrichment-induced loss of forbs in the

128 temperate grassland (Tian *et al.* 2016). Specifically, we found that N-induced
129 accumulation of manganese (Mn) inhibited photosynthetic rates of forbs, thus
130 contributing to their loss (Tian *et al.* 2016). However, this conclusion is based on the
131 results from a relatively long-term (9 years) N-addition experiment at a single site.
132 Whether this mechanism can also explain the short-term and long term effect of N
133 enrichment on plant diversity across different sites remains to be experimentally tested.
134
135 Plant species vary in their responsiveness to N enrichment (Inouye & Tilman 1995;
136 Suding *et al.* 2005; Payne, Dise, Stevens, & Gowing 2013; Tian *et al.* 2016), thus
137 leading to a temporal variation in the plant community (Duprè *et al.* 2010). Given the
138 differences in sensitivities of plant species to varying biotic and/or abiotic conditions,
139 we surmise that plant community dynamics may exhibit phase-based and functional
140 group-dependent characteristics in response to N enrichment. The phased changes in
141 plant community may be driven by N-induced processes in soil. To experimentally test
142 these hypotheses, we investigated dynamic changes in plant community of grasslands
143 in response to varying regimes of N enrichment, to uncover the relative importance of
144 these belowground-mediated processes in vegetation dynamics from long-term and
145 short-term N-addition studies across grassland ecosystems in China.
146

Materials and methods

Field experimental setup

Long-term N-addition experiment

To test whether loss of plant diversity in response to N deposition is phase-dependent, we conducted a long-term field N addition experiment to simulate N deposition in Duolun county (116°17'E, 42°02'N; 1350 m a.s.l.) in Inner Mongolia, China. The grassland in this region has its advantages for studying the phased variations in plant composition and diversity to N enrichment due to its relatively low rate of atmospheric N deposition, abundant forb species and neutrality or weak alkalinity of soil pH. The area is a semi-arid temperate steppe with mean annual temperature of 2.1°C and mean annual precipitation of 382 mm. The atmospheric bulk N deposition in this area was approximately 15.1 kg N ha⁻¹ yr⁻¹ (Xu *et al.* 2015). The detailed properties of soil in this area were described in our previous study (Tian *et al.* 2016). The vegetation is a typical grassland community, comprised by perennial and annual forbs, sedges and grasses (Table S1; Table S2).

The background survey was stated in 2003 after exclusion of livestock grazing and N addition treatment was conducted in 2004. A 8 × 8 Latin square experimental design was used with 8 levels of N addition (0, 1, 2, 4, 8, 16, 32, 64 g N·m⁻² ·yr⁻¹) and 8 replicates. The plot area was 150 m² (15 × 10 m) and each plot was separated by 4 m wide buffer strips. Nitrogen (urea) was applied by hand annually in July when maximal precipitation occurs. Plant species data including aboveground biomass, species

richness and species abundance were determined in mid-August using a 1 x 1 m quadrat each year. The quadrat was randomly placed in each plot that was not overlapped spatially among years by marking with a flag. The species richness was the number of plant species. Individual plants were harvested by clipping the entire aboveground part of each plant to determine their aboveground biomass.

The different functional groups (only including the perennial species) were classified based on plant traits (i.e., cotyledon number, N₂-fixing property, leaf and root characteristics). Based on the cotyledon numbers, the species were divided into forbs and grasses. The *Allium* species were classified as forbs. The N₂-fixing species were leguminous forbs. The non-N₂-fixing forbs with vegetative offspring produced by deep/long rhizomes underground were categorized as “rhizomatous forbs”. The forbs that produce vegetative offspring by stolons or shallow/short rhizomes and the forbs that do not naturally produce vegetative ramets were categorized as “non-rhizomatous forbs”. The non-rhizomatous forbs were classified into broad leaf forbs and arrow leaf forbs based on their leaf architecture. Grass and sedge species were assigned to one of two categories of rhizome-clonal plants included in published dataset (Gough *et al.* 2012): clumper or runner. Grass species with vegetative offspring produced adjacent to parents were categorized as “clumper/bunch”. Grass and sedge species with vegetative offspring produced farther away from parents were categorized as “runner/rhizomatous”. Therefore, each species was assigned to one of six growth forms: N₂-fixing forbs, broad-leaf-non-rhizomatous forbs (BLNR forbs), narrow-leaf-non-

rhizomatous forbs (NLNR forbs), narrow-leaf-rhizomatous forbs (NLR forbs), bunch
grasses, rhizomatous grasses/sedges.

N-addition experiment in Hulun Buir, Qinghai and Xinjiang

A N addition experiment was carried out in 2010 in Hulun Buir (119°35'-119°41' E,
48°27'-48°35' N, 760-770 m above sea level), Inner Mongolia Autonomous Region,
China. This area is characterized by annual precipitation and mean annual temperature
of 329 mm and -1.6°C, respectively. Soil in the site is dark chestnut type according to
China's soil classification system (Hou 1982). The grassland type belongs to the
temperate meadow steppe. The plant species in this grassland mainly includes
Melissitus ruthenica, *Thermopsis lanceolata*, *Viciamul ticaulis*, *Potentilla acaulis*,
Potentilla tanacetifolia, *Potentilla betonicifolia*, *Phlomis umbrosa*, *Sibbaldia sdpressa*,
Artemisia frigida, *Artemisia commutate*, *Thaictru petaloideum*, *Filifolium sibiricum*,
Rhaponticu uniflorum, *Serratula yamasutana*, *Carex pediformis*, *Vicia multicaulis*,
Stipa baicalensis, *Achnatherum sibiricum*, *Carex duriuscula*, *Cleistogenes squarrosa*
and *Leymus chinensis*. *Stipa baicalensis* and *Leymus chinensis* are main dominant
species of the community. The N-addition levels were 0, 1.5, 3, 5, 10, 15, 20, 30 g N
m⁻² yr⁻¹. The plot area was 64 m² (8 m x 8 m) with six replicates for each N treatment.
Nitrogen (NH₄NO₃) was supplied twice a year during the growing season (10-15 June
and 10-15 July) with half of the total amount of supplied on each occasion. Nitrogen
was dissolved in water (8 L) and sprayed into the plots. The control plots were treated
with same amount of water exclusively. For aboveground biomass measurement, the

213 aboveground shoots were clipped within a 1 m x 1 m quadrat in mid-August each year.

214 All shoots of each plant species were over-dried at 70°C for 48 h, and then weighted.

215

216 In Qinghai, the N addition was started in May 2013 on the Sanjiaocheng Sheep

217 Breeding Farm (37°18'N, 100°15'E, 3290 m a.s.l) in an alpine steppe. The mean annual

218 temperature is 0.08°C and the mean annual precipitation is 387 mm (1980-2012). The

219 soil pH is 9.5 and the inorganic nitrogen ($\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$) is about 11.6 mg kg⁻¹.

220 The other detailed properties of soil in this area were described by Peng *et al.* (2017).

221 Eight N levels (0, 1, 2, 4, 8, 16, 24, and 32 g m⁻² yr⁻¹) were established and arranged in

222 randomized complete blocks with five replications. The plot area was 36 m² (6 m x 6

223 m). The distance between any two adjacent plots within a block was 1 m and each block

224 was separated by 2 m buffer strips. Nitrogen fertilizer (ammonium nitrate) was

225 dissolved in 10 L of water and applied by a backpack sprayer at the beginning of each

226 month during the growing season from May to September every year. The same amount

227 of water without N was applied in the control plot. The aboveground biomass of plants

228 was measured by a 1 m x 1 m quadrat in mid-August each year.

229

230 In Xinjiang, The N addition experiment was conducted in May 2009 at the Bayinbuluk

231 Grassland Ecosystem Research Station, Chinese Academy of Sciences (42°53.1'N,

232 83°42.5'E, 2500 m a.s.l). Bayinbuluk alpine grassland is located in the southern

233 Tianshan mountains, Xinjiang Uygur Autonomous Region, China. The mean annual

234 precipitation is 282.3 mm and the mean annual temperature is -4.8 °C. The vegetation

of the grassland was dominated by perennial grasses (*Stipa purpurea*, *Festuca ovina*, *Agropyron cristatum* and *Koeleria cristata*) and perennial forbs (*Oxytropis glabra*, *Potentilla multifida* and *Potentilla bifurca*). Four N levels (0, 1, 3, 9 g m⁻² yr⁻¹) were established different blocks (each 4 m × 8 m) with five replications. Nitrogen (NH₄NO₃) dissolved in water (8 L) were applied to all plots in late May and June each year from 2009 to 2012. Aboveground biomass of plants was determined using a 1 × 1 m quadrat.

Lime and N-addition experiments

To demonstrate the contributions of soil acidification to N deposition-driven changes in plant community, we arranged another N and lime (CaCO₃) addition experiment on 2014 in Duolun county (116°17'E, 42°02'N; 1350 m a.s.l.) in Inner Mongolia. Sixty plots (each 8 m × 10 m) separated by 2 m wide buffer strips were established in a randomized block experimental design. The levels of N addition were 0, 2, 8, 16, 32 g N m⁻² yr⁻¹ and lime addition levels were 0 and 50 g m⁻² yr⁻¹. Each treatment included six replicates. NH₄NO₃ was sprayed into the plots with 12 L water three times a year (mid-May, mid-June and mid-July). Lime was supplied in the mid-October each year. In mid-August each year, aboveground biomass of plants was determined using a 1 × 1 m quadrat. The cover of different functional groups was determined with a square (1 m × 1 m) with one hundred a square decimeter (1 cm × 1 cm) (Yang *et al.* 2011).

Nitrogen-addition experiment with different N forms

To test the effect of different forms of N on different functional groups during the early

periods of N addition, we carried out another field N-addition experiment of different form in 2015 in Duolun county (116°17'E, 42°02'N; 1350 m a.s.l.) in Inner Mongolia of China. The plots (each 5 × 8 m) separated by 2 m wide buffer strips. The N forms were $\text{NH}_4^+\text{-N}$ (NH_4Cl) and $\text{NO}_3^-\text{-N}$ ($\text{Ca}(\text{NO}_3)_2$) and N-addition levels were 0, 2, 8 g N $\text{m}^{-2} \text{yr}^{-1}$. Each treatment included six replicates. Nitrogen (NH_4Cl and $\text{Ca}(\text{NO}_3)_2$) was supplied three times a year (mid-May, mid-June and mid-July). To rule out the effect of nitrification on the transformation of $\text{NH}_4^+\text{-N}$ into $\text{NO}_3^-\text{-N}$, we supplied nitrification inhibitor (0.3 g m^{-2}), 3,4-dimethylpyrazole phosphate (DMPP). The $\text{NH}_4\text{Cl}/\text{Ca}(\text{NO}_3)_2$ and DMPP in aqueous solution (12 L) was evenly sprayed into each plot and the control plots were only treaded with 12 L water.

Greenhouse experiments

Manganese addition to simulate soil Mn accumulation induced by N addition

To determine if the increases of metal concentrations in soils driven by N addition in the field experiment can affect forb and grass growth, we collected the soils from control plots. The air-dried soils were sieved (2 mm mesh size) and thoroughly mixed with different concentrations of MnCl_2 . We controlled exchangeable Mn^{2+} concentrations in soil to similar levels of the N-addition experiment in field. The seeds of dominant species of narrow leaf forbs and bunch grasses, *A. frigida* and *S. krylovii*, were planted in plastic pots (15 cm in diameter x 12 cm in height) filled with 1.5 kg soil. Eight *A. frigida* and *S. krylovii* plants were grown in each pot with six replicates for each treatment. Plants were irrigated with water to maintain 80%-85% of field

capacity. Shoot and root biomass of these plants were determined after their growth for about two months.

Manganese addition experiment under constant pH and nutrient sufficiency

To separate the direct effect of metal toxicity from soil acidification or deficiency in other nutrients, we further conducted Mn-addition experiment by controlling pH and supplying high availability of other nutrients. Seeds of three forb species (*Heteropappus altaicus*, *Artemisia frigida* and *Potentilla bifurica*) and three grass species (*Stipa krylovii*, *Leymus chinensis* and *Agropyron cristatum*) germinated in vermiculite. Three seedlings of each species were transplanted in pots (20-cm diameter) filled with vermiculite. Seedlings in each pot were treated with one-eighth strength Hoagland solution supplemented with varying concentrations of MnCl_2 (0, 50, 100, 250, 500, 1000 $\mu\text{mol L}^{-1}$) with five replicates for each treatment. The pH of nutrient solution was kept at 6.0 ± 0.1 by NaOH and HCl. The shoot and root biomass of these plants were measured after treated for 40 days.

Short-term N-addition experiment

To reveal which factor in soils, metal ion accumulation or increased N availability, determined N-addition-induced suppression of narrow leaf forbs in field, we applied different levels of nitrogen (NH_4NO_3) by controlling the same level of metal ions. Dominant forb species (*A. frigida*) and dominant grass species (*S. krylovii*) were pre-cultured for about one month in vermiculite watered with one-eighth strength Hoagland

solution (pH 6.0 ± 0.1). Six seedlings of each species were transplanted in pots (20 cm diameter) filled with vermiculite. The N (NH_4NO_3) levels was 0, 1, 2, 4, 8, 16, 32 g N m^{-2} . The replicates of each treatment were six. The shoot biomass, plant height, total root length and the number of ramets and tillers were measured after treated with different concentrations of nitrogen for two month.

Analyses of soil properties

Soil samples in each plot were randomly taken using a 3-cm diameter soil core in the plots at a depth of 0-10 cm in mid-August each year. Three-core soils were combined to one soil sample. Soil samples were thoroughly mixed, air-dried and sieved through a 2 mm mesh for measuring soil pH and the concentrations of exchangeable ions. Soil NH_4^+ -N and NO_3^- -N concentrations were measured using fresh soils. Soil NH_4^+ -N and NO_3^- -N concentrations were analyzed with a continuous flow analyzer (Seal XY-2, Australia) after extraction of 2 M KCl at the ratio of 1:5 (w/v). For determination of soil pH, 6 g of air-dried soil was shaken with 15 ml CO_2 -free deionized water for a minute, and equilibrated for an hour to determinate pH with a pH meter (HANNA, PH211, Italy). Exchangeable of Ca^{2+} , Mg^{2+} and K^+ in soil were extracted by 1M NH_4OAc (pH 7.0) at a 1:10 ratio (w/v) for 30 min. The exchangeable Mn^{2+} , Fe^{3+} , Cu^{2+} , Zn^{2+} in soil were extracted with a extracting agent (pH 7.3) consisted of 5 mM diethylenetriamine pentaacetic acid (DTPA), 10 mM CaCl_2 and 0.1 M triethanolamine (TEA) in 1:2 ratio (w/v) for 2 h. The extraction solution was filtered to determine the concentration of Ca^{2+} , Mg^{2+} , K^+ , Fe^{3+} , Mn^{2+} , Cu^{2+} , Zn^{2+} by ICP-OES (Thermo Electron

Corporation, USA).

Measurements of plant physiological traits

Photosynthetic rates of dominant species were measured on sunny days with a LI- 6400 XT portable photosynthesis system equipped with a LED leaf cuvette (Li-Cor, Lincoln, Nebraska, USA). Fully expanded leaves were selected for measure the photosynthetic rates. The parameters were according to those in our previous study (Tian *et al.* 2016). At least two individual plants of one species in each plot were measured for photosynthetic rates.

The negative effects on some plants of N addition were quantified by the inhibitory ratio of yellowing leaf to green leaf on a weight basis. The yellow parts of leaves were cut by scissors and weighed. The remaining green parts of the leaves were also weighed. The inhibitory ratio of N enrichment on functional groups are calculated as the log response ratio: $\ln[\text{mean}(\text{weight ratio of yellowing leaf to green leaf treatment})/\text{mean}(\text{weight ratio control})]$. Three plants of three grasses (*S. krylovii*, *C. squarrosa* and *A. cristatum*) and five forbs (*A. frigida*, *Potentilla acaulis*, *Potentilla tanacetifolia*, *Potentilla betonicifolia*, *Phlomis umbrosa*, and *Sibbaldia sdpressa*) were randomly sampled in each plot for the analysis of inhibitory ratio.

The plant height and primary root length were measured by rule. Roots were scanned with an Epson digital scanner (Expression 10000XL, Epson Inc., Hirooka, Japan) and

analyzed with the WinRHIZO/WinFOLIA software (Regent Instruments Inc., Quebec, Canada) to obtain the total root length. The bud numbers and the root and rhizome biomass were determined by soil cores (30 cm × 30 cm × 20 cm). Roots and rhizomes were randomly sampled in each plot digging a 30 × 30 × 20 cm soil core. The soil cores were transferred to root bags and cleaned with water. The rhizome internode length was measured by a rule. Shoots, roots and rhizomes were separated and dried at 75 °C for 48 h. The ratios of roots to shoots and the ratios of rhizomes to shoots were calculated by the dried biomass.

Data analyses and statistics

The species richness, species density and species frequency were respectively defined as: the number of plant species, plant number of each species and the proportion of species occurrences. We calculated the inhibitory ratios, response ratios and changes in inorganic N and metal ions in soil as the log response ratio: $\ln[\text{mean}(\text{treatment})/\text{mean}(\text{control})]$. The expansion potential of grasses was calculated by the total length of bud internodes (TLBI). We calculated the TLBI as: $\text{TLBI} = \text{LBI} \times \text{BN}$, where LBI and BN are the mean length of bud internodes and the number buds. The change of expansion potential (EP) was calculated as the log response ratio: $\ln[\text{mean}(\text{TLBI treatment})/\text{mean}(\text{TLBI control})]$.

Unpaired *t*-tests were used to evaluate the effect of N addition and metal ion addition on aboveground biomass and species richness. One-way analysis of variance

367 (ANOVAs) performed by the SPSS statistical software (SPSS 17.0 for windows, SPSS
368 Inc., Chicago, IL, USA) was used to compare the difference between the effect of
369 control and metal addition treatments on plant biomass. Two-way analysis of variance
370 (ANOVAs) performed by SPSS was used to examine the effects of N addition, lime
371 addition and the interactive effects of N and lime addition on AGB and cover. Linear
372 and nonlinear regression was performed by SPSS to identify the different responses of
373 functional groups to N addition rates. Nonlinear regression was performed to evaluate
374 the effects of N-addition rates on photosynthate distribution. Linear regression was used
375 to evaluate the effects of N-addition rates on photosynthetic rates, bud numbers,
376 inhibitory ratios, expansion potentials, inorganic N and metal ion concentrations in soils.
377 We chose a significance level of 0.05. The LOWESS method performed by R Version
378 3.3.3 was used to assess the changes in AGB of different functional groups over time.

Results

Effects of N enrichment on aboveground biomass and species richness were phase-dependent

We first monitored the temporal changes in aboveground biomass (AGB) of different functional groups in response to N addition. At the early phase of N addition (≤ 3 years), N addition led to increases in AGB grasses and sedges, while it had little effect on AGB of forbs (Fig. 1A, B). With progression of N addition (≥ 3 years, middle phase), AGB of forbs decreased, while AGB of grasses and sedges increased (Fig. 1A, B). When N addition proceeded longer than 8-10 years (later phase), no significant variations in AGB of forbs, and grasses and sedges were detected (Fig. 1A, B). A contrasting change in species richness of forbs, and grasses and sedges by N addition was observed. For example, N addition at early phase did not affect species richness for both forbs, and grasses and sedges (Fig. 1C, D), but a decline in species richness of forbs by N addition longer than 3 years was found (Fig. 1C). After 10 years of N addition, a decline in species richness of grasses and sedges also occurred (Fig. 1D).

We further classified the forbs and grasses into different growth forms and compared their responses to N addition. Nitrogen enrichment first decreased AGB of N_2 -fixing forbs, followed by broad-leaf-non-rhizomatous (BLNR) forbs (Fig. 2). AGB of narrow-leaf-non-rhizomatous (NLNR) forbs was enhanced during the early phase of N addition,

thereafter it rapidly declined, reaching a constant, and low level during the middle phase of N addition (Fig. 2). Nitrogen addition during the early and middle phases increased AGB of bunch grasses, while it inhibited their growth, and concurrently stimulated growth of rhizomatous species (narrow-leaf-rhizomatous (NLR) forbs, rhizomatous grasses and sedges) at the late phase (Fig. 2). In addition to AGB, N addition-induced changes in frequency of occurrence for forbs and grasses also displayed similar phase-dependent characteristics (Fig. 2). Moreover, the changes in frequency of plant species with different growth forms also displayed different trends in response to N addition (Fig. 2). For example, N_2 -fixing forbs were the first species to disappear in the community, whose frequency decreased at the initial phase of N addition. The significant decreases in frequency of BLNR forbs and NLNR forbs were detected after N addition for 4 and 6 years, respectively (Fig. 2). The frequency of bunch grasses began to decrease after N addition for 8 years, while that of rhizomatous species, NLR forbs and rhizomatous monocots, increased during this period of N addition (Fig. 2).

Effects of N-addition regimes on aboveground biomass and species frequency were dependent on functional groups

Aboveground biomass and species frequency of different growth forms within the forbs and grasses also differed in their responses to N-addition rates. For example, AGB of N_2 -fixing, BLNR and NLNR forbs decreased with increasing N-addition rates, and the magnitude of decrease was greatest for the N_2 -fixing forbs (Fig. 3A). AGB of NLR

forbs was enhanced by lower rates of N-addition ($\leq 8 \text{ g N m}^{-2} \text{ yr}^{-1}$), but it was reduced by higher N-addition rates (Fig. 3A). AGB of bunch grasses and rhizomatous grasses and sedges also exhibited different responses to N addition, such that AGB of rhizomatous grasses and sedges significantly increased with increases in N-addition rates, while AGB of bunch grasses displayed a pattern of single-peak curve with increase in N-addition rates (Fig. 3B). Species frequency among the four growth forms of forbs showed similar responses to N-addition rates to those of AGB (Fig. 3C). Species frequency for bunch grasses declined with increases in N-addition rates, while species frequency of rhizomatous grasses and sedges increased with increasing N-addition rates (Fig. 3D).

To test whether the differential responses of the different functional groups and growth forms to N addition is a general phenomenon, we further analyzed data from temperate steppes in a different site of northeast of China. Similar to results in Figure 3, short-term N addition reduced AGB of the N_2 -fixing and BLNR forbs, while AGB of the NLNR forbs, bunch grasses, and rhizomatous grasses and sedges was enhanced by N addition in the temperate steppe of Hulun Buir of northeastern China ($119^{\circ}35'-119^{\circ}41' \text{ E}$; $48^{\circ}27'-48^{\circ}35' \text{ N}$) (Fig. 4). These results consolidate that the differential responses of functional groups occur independent of grassland locations.

Effects of Nitrogen enrichment on soil properties were also phase-dependent

Short-term N addition led to rapid increases in soil $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations (Fig. 5). Both $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ as well as total inorganic N concentrations peaked after two days of N addition, and then gradually decreased with extending period of N addition (Fig. 5). Despite the decreases in $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations after 2 days of N addition, the total inorganic N concentrations in soils remained significantly higher values than those in the control soils following N addition up to 30 days (Fig. 5).

Compared with soil inorganic N, the variations in other soil properties displayed a relatively slow response to N addition. At the early phase of N addition at a rate of $8 \text{ g N m}^{-2} \text{ yr}^{-1}$ (i.e., $< \sim 4$ years), soil pH and concentrations of several mineral nutrients in soils were relatively unchanged (Fig. 6). When N enrichment was longer than 4-5 years, soil pH and potassium (K^+), calcium (Ca^{2+}) and magnesium (Mg^{2+}) concentrations began to decline, and those metal ions (Fe^{3+} and Mn^{2+}) began to increase (Fig. 6). During the period of 4 to 11 years of N addition, soil pH decreased from 6.82 to 5.68, while the exchangeable Mn^{2+} concentrations in soils increased from 18.0 mg kg^{-1} to 43.0 mg kg^{-1} . With further progression of N enrichment, soil pH and Mn^{2+} concentrations remained relatively constant, with the values in the range of 5.68-5.85, and $35.7\text{-}43.0 \text{ mg kg}^{-1}$, respectively (Fig. 6).

Short-term effects of N-addition on grassland community were independent of soil acidification

467

468 Soil acidification evoked by N enrichment has been implicated in decline in species
469 diversity of grasslands (Crawley et al. 2005; Horswill et al. 2008). To test whether the
470 soil acidification is involved in the change in community dynamics by short-term N
471 addition, we conducted additional experiments by supplying lime in the N-added plots
472 to minimize soil acidification. Short-term N Nitrogen addition at $8 \text{ g N m}^{-2} \text{ yr}^{-1}$
473 increased AGB of NLNR forbs and bunch grasses (Fig. 7B, C), but decreased that of
474 BLNR forbs after 3 years of N addition (Fig. 7A), which are consistent with the results
475 of the above long-term N-addition experiment (cf. Fig. 2). Although lime addition
476 buffered soil pH reduced by N addition, lime addition could not change the effect of N
477 addition on functional groups. For example, N addition-induced reduction in AGB of
478 BLNR forbs and cover of forbs was not recovered by lime addition (Fig. 7A; Fig. S1).
479 Lime addition did not suppress the increase in AGB of NLNR forbs and bunch grasses
480 driven by N addition (Fig. 7B, C), moreover, lime addition did not affect the AGB of
481 NLR forbs and rhizome grasses (Fig. S2). However, Lime addition increased soil pH
482 and alleviated the inhibitory effect of N addition on soil pH (Fig. 7D). These results
483 suggest that N itself, but not soil acidification or the second effects induced by soil
484 acidification, may be a key driver for loss of the N-sensitive forbs, such as the BLNR
485 forbs, during the early stage of N addition.

486

487 We further evaluated the roles of soil pH in N-induced changes in grassland community
488 by monitoring the effects of N addition on AGB and soil pH across the three different

grasslands in Xinjiang (42°53.1'N, 83°42.5'E, soil pH, 7.7), Qinghai (37°18'N, 100°15'E, soil pH, 9.5) and Inner Mongolia (116°17'E, 42°02'N, soil pH, 7.0) with different initial soil pH values in northern China. In general, short-term N addition (1 to 4 years) enhanced AGB of grasses across the three sites, while N addition significantly reduced AGB of forbs after N addition longer than 3 years in grasslands of Xinjing and Qinhai (Fig. 8). An increase in AGB of forbs in Inner Mongolia grassland within 2-yr of N addition was observed, and this stimulatory effect diminished by the 3-rd year of N addition (Fig. 8). The N-induced increase in AGB of grasses and decrease in AGB of forbs were independent of soil pH, as evidenced by no significant changes in soil pH across the three grasslands following short-term N addition (Fig. 8). These results further suggest that soil acidification cannot account for the variations in functional groups during short-term N addition.

The phased changes in aboveground linked to belowground processes

To search for the drivers responsible for the N-induced changes in the grassland community, we linked the phased changes in aboveground to belowground processes. The three distinct changes in aboveground can be summarized as inhibition of AGB of BLNR forbs (early phase), inhibition of AGB of bunch grasses and NLNR forbs (middle phase), and enhancement of AGB of rhizomatous species (late phase) (Fig. 9A, B, C). The three phases were driven by inorganic N concentrations in soil, toxic mental concentrations in soil, and the sum of the N and toxic mental concentrations,

511 respectively (Fig. 9E, F, G). The three drivers in soils underlying the phased changes
512 in aboveground patterns were depicted in Figure 9D. Short-term and first phase of N
513 addition inhibited BLNR forb growth. For instance, the N-induced inhibition of BLNR
514 forbs showed positive correlation with inorganic N concentrations in soils ($R^2=0.79$,
515 $P<0.01$, Fig. 9A), while no such inhibitory effect on NLNR forbs and bunch grasses
516 was detected (Fig. 9A). Nitrogen addition rapidly enhanced inorganic N concentrations
517 in soil, such that the total inorganic N concentrations in soils exhibited a positively
518 linear correlation with N addition rates ($R^2=0.78$, $P<0.01$, Fig. 9E).

519

520 During the middle phase of N addition, the toxic metal-ions, particularly Mn^{2+} in soil,
521 were mobilized by soil acidification, which in turn suppressed NLNR forb growth due
522 to Mn toxicity resulting from their excessive foliar accumulation of Mn (Fig. 9B, F;
523 Tian et al. 2016). The inhibition of aboveground biomass for bunch grasses ($R^2=0.50$,
524 $P=0.003$) and NLNR forbs ($R^2=0.96$, $P<0.01$) showed positive correlation with the
525 metal ion concentrations in soils, with the magnitude of inhibition for NLNR forbs
526 being greater than that for bunch grasses (Fig. 9B). After long-term N addition, soil pH
527 was linearly reduced with increases in N addition rates ($R^2=0.72$, $P<0.01$, Fig. 9F),
528 while soil Mn^{2+} concentrations linearly increased with increasing N-addition rate
529 ($R^2=0.84$, $P<0.01$, Fig. 9F).

530

531 During the late stage of N addition, the expansion potential for space in rhizomatous
532 species was enhanced by N addition, while that of bunch grasses did not significantly

change (Fig. 9C). The expansion potential of rhizomatous species showed positive correlation with the concentrations of inorganic N and metal-ions in soils ($R^2=0.75$, $P<0.01$), and the correlation between the two variables in bunch grasses was not significant (Fig. 9C). The increase in the expansion potential of rhizomatous species resulted from the increase in their bud numbers of rhizomes underground (Fig. 9G). The bud numbers of rhizomatous monocots ($R^2=0.53$, $P=0.01$) and bunch grasses ($R^2=0.48$, $P<0.01$), positively correlated with N addition rates (Fig. 9G). However, rhizomatous species developed more buds than bunch grasses with increase in N-addition rates (Fig. 9G). These results suggest that accumulation of nutrients in soils, mainly including inorganic N and metal ions, is responsible for the bud differentiation-induced increase in their expansion potential (Fig. 9C).

Ecophysiological mechanisms for the phase-dependent community changes

The negative effect of N enrichment on BLNR forbs at the early phase of N addition was evidenced by distinct chlorosis in their blade edges, and suppression of their photosynthetic rates (Fig. 10A). In contrast, the same N-addition regimes had no such effects on NLNR forbs and grasses; rather, it enhanced the photosynthetic rates of grasses (Fig. 10A). The decline in AGB of BLNR forbs by N addition may result from ammonium toxicity. Indeed, NH_4^+ -N addition inhibited the photosynthetic rates in BLNR forbs, while the same treatment did not affect the photosynthetic rates in other functional groups including NLNR forbs, bunch grasses and rhizomatous

grasses/sedges (Fig. S3). The rapid accumulation of $\text{NH}_4^+\text{-N}$ was the early driver of loss of some sensitive species to N addition, because N addition led to a rapid increase in $\text{NH}_4^+\text{-N}$ concentrations (Fig. 5A).

In contrast, soil Mn^{2+} concentrations did not change in the first three years of N addition, and they were gradually increased to the peak level of 43.0 mg kg^{-1} from 6.0 mg kg^{-1} along with soil acidification (Fig. 6). Addition of the similar levels of Mn^{2+} to that in the field under N-added conditions suppressed belowground and aboveground growth of the NLNR forbs, but it had little effect on growth of bunch grasses (Fig. 9B; Fig. S4). Exogenous application of Mn to growth medium not only inhibited root elongation of NLNR forbs, but also suppressed their occurrence of adventitious roots, while it had no effect on root growth of grasses (Fig. 10B). Despite a high availability of other nutrients, NLNR forbs displayed greater sensitivity to Mn^{2+} than grasses (Fig. S5). However, the dominant NLNR forb, *Artemisia frigida*, displayed a similarly positive response to increases in N availability to the bunch grass, *Stipa krylovii* (Fig. S6). These results suggest that Mn^{2+} toxicity, but not increases in N and deficiency in other nutrients, may account for the N addition-induced loss of NLNR forbs during the middle phase of N addition.

The rhizome/shoot ratios and root/shoot ratios of rhizomatous monocots (rhizomatous grasses and sedges) and increased with increasing N-addition rates (Fig. 10C). However, the root/shoot ratios of bunch grasses showed negative correlations with N-addition

rates (Fig. 10C). These results suggest that nutrient enrichment in soil with progression of N addition favors allocation of photoassimilate to belowground parts in rhizomatous plants, especially to the reproductive organs of rhizomes. Therefore, an increase in rhizomatous bud number driven by assimilate allocation may contribute to the dominance of rhizome species in later phase of N addition.

To further illustrate the linkage between underground drivers and changes in aboveground biomass during the three phases of N addition, we examined the time-dependent changes in AGB among different functional groups in response to N addition by the method LOWESS (Fig. 11A). A rapid increase in inorganic soil N concentrations during the first 3 years of N addition may result in ammonia toxicity, thus driving the first phase of N-induced decreases in aboveground biomass of the N₂-fixing forbs and BLNR forbs (Fig. 11A, B). With the progression of N addition longer than 3 years, an accumulative increase in soil metal ions, mainly Mn²⁺, due to N-induced soil acidification drove the loss of those Mn-sensitive NLNR forbs from plant community (Fig. 11A, C). During the early and middle phases of N addition, rhizome-bud numbers of rhizomatous species did not show significant variations (Fig. 11D). However, we found that N addition for 15 years led to a significant increase in bud numbers on the rhizomes of rhizomatous grasses and sedges due to greater allocation of photoassimilate to the rhizomes, thus driving the dominance of rhizomatous grasses and sedges in the community after prolonged N addition (Fig. 11A, D).

Discussion

Numerous theories have been used to explain dynamics of grassland community in response to N enrichment. These include interspecific competition (Grime 1973; Tilman 1987), phosphorus limitation (Aerts, Wallen, & Malmer 1992; Phoenix *et al.* 2004; Thayer, St Clair, Field, & Somerville 2008), soil acidification (Crawley *et al.* 2005; Horswill, O'sullivan, Phoenix, & Leake 2008), ion toxicity (i.e. NH_4^+ , Al^{3+}) (Stevens, Dise, Gowing, & Mountford 2006; Maskell, Smart, Bullock, Thompson, & Stevens 2010), and decline in niche dimensionality (McKane *et al.* 2002; Harpole & Tilman 2007). However, as for temporal dimensions, we know little about the relative importance and the linkage of these hypotheses in explaining the N-induced shift of the community composition. As for spatial dimensions, it also lacks the eco-physiological dataset to dissect the mechanisms by linking aboveground to underground processes under the scenarios of N enrichment. To fill this knowledge gap, we compiled data from *in situ* field N addition experiments in grasslands across different sites in China with varying N addition regimes, and combined with results in greenhouse experiments. We discovered that the phase- and belowground-driven processes underlay the N-evoked periodic loss of plant species in grasslands. We further explored the ecophysiological mechanisms underlying the phase-based, functional group-dependent and belowground-driven changes in the community composition by integrating the temporal and spatial features under varying regimes of N addition.

One important finding in the present study is that loss of plant biodiversity in temperate

grasslands by N enrichment exhibits distinct phase- and functional group-dependent features. For example, we found that N addition firstly suppressed growth of N₂-fixing plants and broad-leaf-non-rhizome forbs due to an increase in inorganic N availability in soils. These species may have been adapted physiologically to low soil N availability, and exposure to high soil N disrupts their physiological processes due to suppression of the nitrogenase activity (Halvorson, Franz, Smith, & Black, 1992; Hungria, Franchini, Campo, & Graham 2005) and/or NH₄⁺ toxicity (van den Berg *et al.* 2005; Li, Li, Kronzucker, Baluska, & Shi 2014). In this context, we observed that the broad-leaf-non-rhizome (BLNR) forbs were more sensitive to NH₄⁺ among the NLNR forbs, bunch and rhizomatous grasses in terms of inhibitory effects of NH₄⁺ on photosynthetic rates (Fig. S3). The reduction in photosynthetic rates in these species would render them less competitive for other resources including nutrients and light (Borer *et al.* 2014; Farrer & Suding 2016), thus driving their loss under N-enriched conditions. Data from our lime-addition experiments (cf. Fig. 7) also support that N itself, but not soil acidification and/or the soil acidification-induced second effects, is a key driver for loss of the N-sensitive forbs during the early phase of N addition. The decline in plant biodiversity during early phase of N addition is independent of soil pH was further consolidated by our results from three N-addition experiments in grasslands across different geographic locations with varying initial soil pH values (Fig. 8). The observation that N enrichment significantly altered forb and grass growth prior to soil acidification (Fig. 8) is also in line with this argument.

643 Compared with increased N availability in soil, soil acidification driven by N
644 enrichment is a relatively slow process, e.g., a significant decrease in soil pH was
645 observed after ~ 5 years of N addition at moderate rate of 8 g N m² yr⁻¹ (Fig. 6). One
646 consequence of soil acidification is the changes in availability of metal ions. For
647 example, a marginal reduction in soil pH by moderate N enrichment can lead to an
648 increase soil Mn²⁺ concentrations substantially (Fig. 9F). The increases in soil Mn²⁺
649 concentration led to greater foliar accumulation of Mn by forbs than by grasses (Tian
650 *et al.* 2016). The excessive Mn accumulation suppressed their photosynthetic rates
651 (Tian *et al.* 2016) and inhibited root elongation (Fig. 10B). These observations may
652 provide a mechanistic explanation for the loss of NLNR forbs in the community in the
653 middle phase (~ 5- 10 years) of N addition.

654
655 Chronic N deposition has been suggested to result in P limitation for plant growth
656 (Phoenix *et al.* 2004; Menge & Field 2007; Thayer, St Clair, Field, & Somerville 2008),
657 which in turn may evoke exudation of carboxylates to mobilize soil P, and concurrently
658 enhance Mn accumulation by plants (Lambers, Hayes, Laliberté, Oliveira, & Turner
659 2015). In the present study, an increase in foliar N:P ratio was detected for both forbs
660 and grasses in response to N addition, but the magnitude of the increase was greater for
661 grasses than forbs (Fig. S7). The highest N:P value for forbs under N addition was 14.0
662 $\pm$ 0.8, suggesting N limitation, rather than P-limitation, for the species growth
663 (Güsewell & Koerselman 2002). The N:P value for grasses increased above 16.0 when
664 N-addition rate exceeded 4 g m⁻² yr⁻¹ (Fig. S7), suggesting that P-limitation might occur

in grasses under N addition. We are aware that the N:P values at the species- and/or functional group-levels may not be reliable indicators to truly reflect P limitation at the community level (Jiang, Caldararu, Zaehle, Ellsworth, & Medlyn 2019). Therefore, future studies on the N-induced P limitation need to consider P mobilization, P demand and redistribution at different levels.

We demonstrated that ammonium toxicity, Mn^{2+} toxicity, and meristem differentiation induced by nutrient enrichment act as different filters to drive the phase-dependent variations of grassland communities. However, we do not claim that these mechanisms operate separately from each other in a temporal sense. For example, the nutrient-enrichment-driven increases in expansion potential of rhizomatous grasses and sedges can also play a role in the middle phase of N addition, as increases in soil inorganic N availability during this phase also occurred (Fig. 11B). Inorganic N availability in soil can regulate root system architecture by promoting meristem differentiation (Forde 2002; Desnos 2008). Schoolmaster, Mittelbach, and Gross (2014) have emphasized the importance of spatial strategies on the community response to fertilization by modeling. Unfortunately, they did not provide the physiological evidence for this process (Schoolmaster, Mittelbach, & Gross 2014). Our results indicate that greater allocation of photoassimilate to reproductive organs (rhizomes and buds) in belowground may allow the rhizomatous monocots to occupy more space by increasing rhizome buds, thus driving the dominance of rhizomatous species in the community after prolonged N addition (Fig. 9; Fig. 10). These findings highlight that the linkage between

underground and aboveground processes plays an important role in the shift of plant composition in response to N enrichment.

Our results revealed the shift in relative importance of the three mechanisms across varying periods (1 to 15 year) of N addition. However, we do not negate the roles of other factors in the N-enrichment-induced decline in plant diversity, such as litter accumulation (Foster & Gross 1998), variation in soil microbial communities (Dean *et al.* 2014; Leff *et al.* 2015) and aluminum (Al) toxicity (Maskell, Smart, Bullock, Thompson, & Stevens 2010). On the contrary, our results may provide explanations for the roles of these factors in the overall decline in plant diversity under elevated N regimes. For example, the N-induced damages to leaves of BLNR forbs in an early phase of N addition may contribute to an increase in litter accumulation. The enhanced litter accumulation would further inhibit the growth of some low stature plants in the community by lowering light penetration (Foster & Gross 1998). However, litter accumulation does not affect rhizomatous plants in the same way, because differentiation of rhizomatous buds occurs in soil, and the rhizomatous buds can break through soil resistance and develop into mature plants.

Metal toxicity to plants is closely dependent on soil properties. Most acidic grasslands have reached a potentially toxic level of aluminum (Al^{3+}) (Bowman, Cleveland, Halada, Hresko, & Baron 2008). Therefore, Al toxicity is often a key driver contributing to N deposition-induced loss of plant species in acidic grasslands (Bowman, Cleveland,

Halada, Hresko, & Baron 2008). However, soils of most temperate grasslands are buffered by base cations (Bowman, Cleveland, Halada, Hresko, & Baron 2008) and characterized by low soil Mn availability. Plant species in the temperate grasslands may have evolved strategies to adapt to the low soil Mn^{2+} availability. The enhanced Mn^{2+} due to soil acidification and/or potentially P mobilization exposed the native temperate grassland species to Mn toxicity, particularly for forb species (Tian *et al.* 2016). We expected the ecosystem would shift from Mn to Al toxicity with a further decline of soil pH, because Al^{3+} will replace base cations to buffer soil acidification at pH ranges below pH 4.5 (Bowman, Cleveland, Halada, Hresko, & Baron 2008).

The temporal dimensions of the proposed three mechanisms depend on soil traits (Bowman, Cleveland, Halada, Hresko, & Baron 2008), plant composition (Gross & Mittelbach 2017) and background levels of atmospheric N deposition in the ecosystems (Stevens, Dise, Mountford, Gowing 2004). In the ecosystems with a relatively high soil pH, the early phase associated with N effects may last longer, and can be a major mechanism to determine N-induced changes in community composition. For example, no soil acidification was detected in the alpine grasslands of the Qinghai-Tibet Plateau with soil pH > 8 after N addition for 4 years (Fig. 8). In contrast, for the acidic grasslands in Europe with soil pH < 5, the first phase involved in direct effect of N itself would be relatively short and less important. Moreover, the second phase of metal toxicity for these acidic grasslands, mainly Al toxicity, may dominate the overall mechanisms of N-evoked community change. For the steppe community co-dominated

by bunch grasses and rhizomatous grasses and sedges, the stimulation of clonal propagation underground by prolonged N enrichment may be a key mechanism responsible for the dominance of rhizomatous species.

Conclusion

We discovered that a phase-based, functional group-specific and belowground-driven mechanism accounts for the commonly observed community shift evoked by N enrichment in grassland ecosystems. These findings are of general significance for our mechanistic understanding of N-dependent changes in plant diversity of grassland community, and can be extended to other terrestrial ecosystems suffering from the enhanced N input.

Acknowledgements

This work is supported by National Natural Science Foundation of China (31870457, 31830011 and 31570403). We thank Dr. Kaihui Li (Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences) and Prof. Yuanhe Yang (Institute of Botany, Chinese Academy of Sciences), for providing data of soil pH and aboveground biomass of the N addition experiments in Xinjiang and Qinghai province. We are grateful to Prof. Dianlin Yang at Agro-Environment Protection Institute Ministry of Agriculture for providing biomass data of plant species of N addition experiment in Hulun Buir, Inner Mongolia. We thank the staff at the Duolun Restoration Ecology Research Station, Institute of Botany, Chinese Academy of Sciences, for their help in field investigation.

Authorship

Q.T. and W.H.Z conceived and designed the experiments. Q.T., L.Y., P.M., N.L., H.W. W.B., W.H., P. L., and L. R. performed the experiments. Q.T., H.L. and W.H.Z. wrote the manuscript. P.A.S., J.D.B. and F.S.Z. commented on the details of the manuscript drafts.

Supporting Information

References

- Aerts, R., Wallen, B., & Malmer, N. (1992). Growth-limiting nutrients in *Sphagnum*-dominated bogs subject to low and high atmospheric nitrogen supply. *Journal of Ecology* **80**, 131-140. DOI: 10.2307/2261070
- Baer, S. G., Blair, J. M., & Collins, S. L. (2015). Environmental heterogeneity has a weak effect on diversity during community assembly in tallgrass prairie. *Ecological Monographs* **86**, 94-106. DOI: 10.1890/15-0888.1
- Bai, W. M., Guo, D. L., Tian, Q. Y., Liu, N. N., Cheng, W. X., Li, L. H., & Zhang, W. H. (2015). Differential responses of grasses and forbs led to marked reduction in below-ground productivity in temperate steppe following chronic N deposition. *Journal of Ecology* **103**, 1570-1579. DOI: 10.1111/1365-2745.12468
- Bobbink, R., Ashmore, M. Braun, S., Flückiger, W., & van den Wyngaert, I. J. J. (2003). Empirical nitrogen critical loads for natural and semi-natural ecosystems: 2002 update. In B. Achermann & R. Bobbink (Eds), *Empirical critical loads for nitrogen. Swiss Agency for Environment, Forest and Landscape SAEFL* (pp.143-170). Berne, Switzerland.
- Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., ... De Vries, W. (2010). Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecological Applications* **20**, 30-59. DOI: 10.1890/08-1140.1
- Borer, E. T., Seabloom, E.W., Gruner, D.S., Harpole, W.S., Hillebrand, H., Lind, E. M., ... Yang, L. H. (2014). Herbivores and nutrients control grassland plant diversity via light limitation. *Nature*, 508, 517-521. Herbivores and nutrients control grassland plant diversity via light limitation. *Nature* **508**, 517-520. DOI: 10.1038/nature13144
- Bowman, W. D., Cleveland, C. C., Halada, L., Hresko, J., & Baron, J. S. (2008). Negative impact of nitrogen deposition on soil buffering capacity. *Nature Geoscience* **1**, 767-770. DOI: 10.1038/ngeo339
- Cardinale, B. J. Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., ... Naeem, D. (2012). Biodiversity loss and its impact on humanity. *Nature* **486**, 59-67. DOI: 10.1038/nature11148
- Clark, C. M., & Tilman, D (2008). Loss of plant species after chronic low-level nitrogen deposition to prairie grasslands. *Nature* **451**, 712-715. DOI: 10.1038/nature06503
- Clark, C. M., Cleland, E. E., Collins, S. L., Fargione, J. E., Gough, L., Gross, K.L., ... Grace, J. B. (2007). Environmental and plant community determinants of species loss

- 798 following nitrogen enrichment. *Ecological Letters* **10**, 596-607. DOI:
799 10.1111/j.1461-0248.2007.01053.x
- 800 Crawley, M. J., Johnstone, S. E., Silvertown, J., Dodd, M., de Mazancourt, C., Heard,
801 M. S., ... Edwards, G. R. (2005). Determinants of species richness in the Park Grass
802 experiment. *American Naturalist* **165**, 179-192. DOI: 10.1086/427270
- 803 Dean, S. L., Farrer, E. C., Taylor, D. L., Porras-Alfaro, A., Suding, K. N., & Sinsabaugh,
804 R. L. (2014). Nitrogen deposition alters plant-fungal relationships: linking
805 belowground dynamics to aboveground vegetation change. *Molecular Ecology* **23**,
806 1364-1378. DOI: 10.1111/mec.12541
- 807 Desnos, T. (2008). Root branching responses to phosphate and nitrate. *Current Opinion*
808 *in Plant Biology* **11**, 82-87. DOI: 10.1016/j.pbi.2007.10.003
- 809 Dickson, T. L., & Foster, B. L. (2011) Fertilization decreases plant biodiversity even
810 when light is not limiting. *Ecology Letters* **14**, 380-388. DOI: 10.1111/j.1461-
811 0248.2011.01599.x
- 812 Duprè, C., Stevens, C. J., Ranke, T., Bleeker, A., Pepler-Lisbach, C., Gowing, D. J.
813 G., ... Diekmann, M. (2010). Changes in species richness and composition in
814 European acidic grasslands over the past 70 years: the contribution of cumulative
815 atmospheric nitrogen deposition. *Global Change Biology* **16**, 344-357. DOI:
816 10.1111/j.1365-2486.2009.01982.x
- 817 Farrer, E. C., & Suding, K. N. (2016). Testing apart plant community responses to N
818 enrichment: the roles of resource limitation, competition and soil microbes. *Ecology*
819 *Letters* **19**, 1287-1296. DOI: 10.1111/ele.12665
- 820 Forde, B. G. (2002). Local and long-range signaling pathways regulating plant
821 responses to nitrate. *Annual Review of Plant Biology* **53**, 203-224. DOI:
822 10.1146/annurev.arplant.53.100301.135256
- 823 Foster, B. L., & Gross, K. L. (1998). Species richness in a successional grassland:
824 effects of nitrogen enrichment and litter. *Ecology* **79**, 2593-2602.
- 825 Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z. C., Freney, J.
826 R., ... Sutton, M. A. (2008). Transformation of the nitrogen cycle: recent trends,
827 questions, and potential solutions. *Science* **320**, 889-892. DOI:
828 10.1126/science.1136674
- 829 Gough, L., Gross, K. L., Cleland, E. E., Clark, C. M., Collins, S. L., Fargione, J. E., ...
830 Suding, K. N. (2012). Incorporating clonal growth form clarifies the role of plant

- 831 height in response to nitrogen addition. *Oecologia* **169**, 1053-1062. DOI:
 832 *10.1007/s00442-012-2264-5*
- 833 Grime, J. P. (1973). Competitive exclusion in herbaceous vegetation. *Nature* **242**, 344-
 834 347. DOI: 10.1038/242344a0
- 835 Gross, K. L., & Mittelbach, G. G. (2017). Negative effects of fertilization on grassland
 836 species richness are stronger when tall clonal species are present. *Folia Geobotanica*
 837 **52**, 401-409. DOI: 10.1007/s12224-017-9300-5
- 838 G ü sewell, S., & Koerselman W. (2002). Variation in nitrogen and phosphorus
 839 concentrations of wetland plants. *Perspective in Plant Ecology and Evolution System*
 840 **5**, 37-61. DOI: 10.1078/1433-8319-0000022
- 841 Halvorson, J. J., Franz, E. H., Smith, J. L., & Black, R. A. (1992). Nitrogenase activity,
 842 nitrogen-fixation, and nitrogen inputs by lupines at mount St, Helens. *Ecology* **72**,
 843 87-98. DOI: 10.2307/1938723
- 844 Harpole, W. S., & Tilman, D. (2007). Grassland species loss resulting from reduced
 845 niche dimension. *Nature* **446**, 791-793. DOI: 10.1038/nature05684
- 846 Hautier, Y., Niklaus, P. A., & Hector, A. (2009). Competition for light causes plant
 847 biodiversity loss after eutrophication. *Science* **324**, 636-638. DOI:
 848 10.1126/science.1169640
- 849 Horswill, P., O'sullivan, O., Phoenix, G. K., & Leake, J. R. (2008). Base cation
 850 depletion, eutrophication and acidification of species-rich grasslands in response to
 851 long-term simulated nitrogen deposition. *Environmental Pollution* **155**, 336-349.
 852 DOI: 10.1016/j.envpol.2007.11.006
- 853 Hou, X. Y. (1982) *Vegetation Geography of China and Chemical Composition of*
 854 *Dominant Species*. Science Press, Beijing.
- 855 Houdijk, A. L. F. M., Verbeek, P. J. M., Vandijk, H. F. G., & Roelofs, J. G. M. (1993)
 856 Distribution and decline of endangered herbaceous heathland species in relation to
 857 the chemical-composition of the soil. *Plant and Soil* **148**, 137-143. DOI:
 858 *10.1007/BF02185393*
- 859 Hungria, M., Franchini, J. C., Campo, R. J., & Graham, P. H. (2005). The importance
 860 of nitrogen fixation to soybean cropping in South America. In D. Werner and W. E.
 861 Newton (Eds). *Nitrogen Fixation in Agriculture, Forestry, Ecology, and the*
 862 *Environment* (pp. 25-42.). Springer, Dordrecht, The Netherlands,.
- 863 Inouye, R. S., & Tilman, D. (1995). Convergence and divergence of old-field vegetation
 864 after 11 yr of nitrogen addition. *Ecology* **76**, 1872-1887. DOI: 10.2307/1940720
- 865 Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., ...
 866 Eisenhauer, N. (2015). Biodiversity increases the resistance of ecosystem

- productivity to climate extremes. *Nature* **526**, 574-577. DOI: 10.1038/nature15374
- Jiang, M., Caldararu, S., Zaehle, S., Ellsworth, D. S., & Medlyn, B. E. (2019) Towards a more physiological representation of vegetation phosphorus processes in land surface models. *New Phytologist* **222**, 1223-1229. DOI: 10.1111/nph.15688
- Kimball, S., Goulden, M. L., Suding, K. N., & Parker, S. (2014). Altered water and nitrogen input shifts succession in a southern California coastal sage community. *Ecological Applications* **24**, 1390-1404. DOI: 10.1890/13-1313.1
- Kleijn, D., Bekker, R. M., Bobbink, R., de Graaf, M. C. C., & Roelofs, J. G. M. (2008) In search for key biogeochemical factors affecting plant species persistence in heathland and acidic grasslands: a comparison of common and rare species. *Journal of Applied Ecology* **45**, 680-687. DOI: 10.1111/j.1365-2664.2007.01444.x
- Laliberté, E., Zemunik, G., & Turner, B. L. (2014) Environmental filtering explains variation in plant diversity along resource gradients. *Science* **345**, 1602-1605. DOI: 10.1126/science.1256330
- Lambers, H., Hayes, P. E., Laliberté, E., Oliveira, R. S., & Turner, B. L. (2015). Leaf manganese accumulation and phosphorus-acquisition efficiency. *Trends in Plant Science* **20**, 83-90. DOI: 10.1016/j.tplants.2014.10.007
- Larsen, S., Chase, J. M., Durance, I., & Ormerod, S. J. (2018). Lifting the veil: richness measurements fail to detect systematic biodiversity change over three decades. *Ecology* **99**, 1316-1326. DOI: 10.1002/ecy.2213
- LeBauer, D. S., & Treseder, K. K. (2008). Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* **89**, 371-379. DOI: 10.1890/06-2057.1
- Leff, J. W., Jones, S. E., Prober, S. E., Barberan, A., Boret, E.T., Firn, J. L.,... Fierer, N. (2015). Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of National Academy of Science USA* **112**, 10967-10972. DOI: 10.1073/pnas.1508382112
- Levin, S. A. (1974). Dispersion and population interactions. *American Naturalist* **106**, 207-228. DOI: 10.1086/282900
- Li, B. H., Li, G. J., Kronzucker, H. J., Baluska, F., & Shi, W. M. (2014). Ammonium stress in Arabidopsis: signaling, genetic loci, and physiological targets. *Trends in Plant Science* **19**, 107-114. DOI: 10.1016/j.tplants.2013.09.004
- Li, Y., Niu, S., & Yu, G. (2016). Aggravated phosphorus limitation on biomass production under increasing nitrogen loading: a meta-analysis. *Global Change Biology* **22**, 934-943. DOI: 10.1111/gcb.13125
- Mariotte, P., Mehrabi, Z., Bezemer, T. M., De Dryn, G. B., Kulmatiski, A., Drigo, B., ... Kardol, P. (2018). Plant-soil feedback: bridging natural and agricultural sciences. *Trends in Ecology and Evolution* **33**, 129-142.
- Maskell, L. C., Smart, S. M., Bullock, J. M., Thompson, K., & Stevens, C. J. (2010)

- 906 Nitrogen deposition causes widespread loss of species richness in British habitats.
 907 *Global Chang Biology* **16**, 671-679. DOI: 10.1016/j.tree.2017.11.005
- 908 McKane, R. B., Johnson, L.C., Shaver, G. R., Nadelhoffer, K. J., Rastetter, E. B., Fry,
 909 B.,... Murray, G. (2002). Resource-based niches provide a basis for plant species
 910 diversity and dominance in arctic tundra. *Nature* **415**, 68-71. **DOI:**
 911 **10.1038/415068a**
- 912 Menge, D. N. L., & Field, C .B. (2007). Simulated global changes alter phosphorus
 913 demand in annual grassland. *Global Chang Biology* **13**, 2582-2591. DOI:
 914 10.1111/j.1365-2486.2007.01456.x
- 915 Payne, R. J., Dise, N. B., Stevens, C. J., & Gowing, D. J. (2013). Impact of nitrogen
 916 deposition at the species level. *Proceedings of National Academy of Science USA*
 917 **110**, 984-987. DOI: 10.1073/pnas.1214299109
- 918 Peng, Y., Li, F., Zhou, G. Y., Fang, K., Zhang, D. Y., Li, C. B., ...Yang, Y. H. (2017).
 919 Nonlinear response of soil respiration to increasing nitrogen additions in a Tibetan
 920 alpine steppe. *Environmental Research Letters* **12**, 024018. DOI: 10.1088/1748-
 921 9326/aa5ba6
- 922 Phoenix, G. K., Booth, R. E., Leake, J. R., Read, D. J., Grime, J. P., & Lee, J. A. (2004)
 923 Simulated pollutant nitrogen deposition increases P demand and enhances root-
 924 surface phosphatase activities of three plant functional types in a calcareous
 925 grassland. *New Phytologist* **161**, 279-289. DOI: 10.1046/j.1469-8137.2003.00910.x
- 926 Reich, P. B. (2009). Elevated CO₂ reduces losses of plant diversity caused by nitrogen
 927 deposition. *Science* **326**, 1399-1402. DOI: 10.1126/science.1178820
- 928 Schoolmaster, D. R., Mittelbach, G. G., & Gross, K. L. (2014). Resource competition
 929 and community response to fertilization: the outcome depends on spatial strategies.
 930 *Theoretical Ecology* **7**, 127-135. DOI: 10.1007/s12080-013-0205-5
- 931 Stein, A., Gerstner, K. & Kreft, H. (2014). Environmental heterogeneity as a universal
 932 driver of species richness across taxa, biomes and spatial scales. *Ecology Letter* **10**,
 933 677-684. DOI: 10.1111/ele.12277
- 934 Stevens, C. J., Dise, N. B., Gowing, D. J., & Mountford, J. O. (2006). Loss of forb
 935 diversity in relation to nitrogen deposition in the UK: regional trends and potential
 936 controls. *Global Chang Biology* **12**, 1823-1833. DOI: 10.1111/j.1365-
 937 2486.2006.01217.x
- 938 Stevens, C. J., Dise, N. B., Mountford, J. O., & Gowing, D. J. (2004). Impact of nitrogen
 939 deposition on the species richness of grasslands. *Science* **303**, 1876-1879. DOI:
 940 10.1126/science.1094678
- 941 Suding, K. N., Collins, S. L., Gough, L., Clark, C., Cleland, E. E., Gross, K. L., ...
 942 Pennings, S. (2005). Functional- and abundance-based mechanisms explain diversity

- loss due to N fertilization. *Proceedings of National Academy of Sciences USA* **102**, 4387-4392. DOI: 10.1073/pnas.0408648102
- Thayer, S. S., St Clair, S. B., Field, C. B., & Somerville S. C. (2008). Accentuation of phosphorus limitation in *Geranium dissectum* by nitrogen: an ecological genomic study. *Global Chang Biology* **14**, 1877-1890. DOI: 10.1111/j.1365-2486.2008.01618.x
- Tian, D., & Niu S. (2015). A global analysis of soil acidification caused by nitrogen addition. *Environmental Research Letters* **10**, 024019. DOI: 10.1088/1748-9326/10/2/024019
- Tian, Q. Y., Liu, N. N., Bai, W. M., Li, L. H., & Zhang, W. H. (2015). Disruption of metal ion homeostasis in soils is associated with nitrogen deposition-induced species loss in an Inner Mongolia steppe. *Biogeosciences* **12**, 3499-3512. DOI: 10.5194/bg-12-3499-2015
- Tian, Q. Y., Liu, N. N., Bai, W. M., Li, L. H., Chen, J. Q., Reich, P. R., ... Zhang, W. H. (2016). A novel soil manganese mechanism drives plant species loss with increased nitrogen deposition in a temperate steppe. *Ecology* **97**, 65-74. DOI: 10.1890/15-0917.1
- Tilman, D. (1987). On the meaning of competition and the mechanisms of competitive superiority. *Functional Ecology* **1**, 304-315. DOI: 10.2307/2389785
- Tilman, D. (1993). Species richness of experimental productivity gradients: how important is colonization limitation. *Ecology* **74**, 2179-2191. DOI: 10.2307/1939572
- van den Berg, L. J. L., Dorland, E., Vergeer, P., Hart, M. A., Bobbink, B., & Roelofs, J. G. M. (2005). Decline of acid-sensitive plant species in heathland can be attributed to ammonium toxicity in combination with low pH. *New Phytologist* **166**, 551-564. DOI: 10.1111/j.1469-8137.2005.01338.x
- Vitousek, P. M., Abert, J. D., Howarth, R. W., Likens, G. E., Matson, P. A., Schlesinger, W. H., & Tilman, D. (1997) Human alteration of the global nitrogen cycle: sources and consequences. *Ecological Applications* **7**, 737-750. DOI: 10.2307/2269431
- Xu, W., Luo, X. S., Pan, Y. P., Zhang, L., Tang, A. H., Shen, J. L., ... Liu, X. J. (2015). Quantifying atmospheric nitrogen deposition through a nationwide monitoring network across China. *Atmospheric Chemistry and Physics* **15**, 18365-18405. DOI: 10.5194/acp-15-12345-2015
- Yang, H. J., Wu, M. Y., Liu, W. X., Zhang, Z., Zhang, N. L., & Wan, S. Q. (2011) Community structure and composition in response to climate change in a temperate steppe. *Global Chang Biology* **17**, 452-465. DOI: 10.1111/j.1365-2486.2010.02253.x
- Zuppinger-Dingley, D., Schmid, B., Petermann, J. S., Yadav, V., de Deyn, G. B., & Flynn, D. F. B. (2014). Selection for niche differentiation in plant communities

981 increases biodiversity effects. *Nature* **515**, 108-111. DOI: 10.1038/nature13869
982
983

Figure legends

Figure 1 Changes in aboveground biomass (AGB) and species richness of the two main functional groups: forbs, grasses and sedges, in response to nitrogen (N) addition across 1 to 15 years. Left, forbs; right, grasses and sedges. (A) AGB of forbs; (B) AGB of grasses and sedges (C) species richness of forbs; (D) species richness of grasses and sedges. The “-N” and “+N” were the treatments with N-addition rate of 0 g N m⁻² yr⁻¹ and 8 g N m⁻² yr⁻¹. Species AGB and species number (species richness) were determined in quadrats (1 m × 1 m) at the peak biomass in the middle of August each year. Data are means ± SE (n=8). Significant differences are denoted by “*” ($P < 0.05$).

Figure 2. Time- and functional group-dependent changes in aboveground biomass and species frequency of functional groups in response to N addition ranging from 1 to 15 years. The “-N” and “+N” were the treatments with N-addition rate of 0 g N m⁻² yr⁻¹ and 8 g N m⁻² yr⁻¹. The N₂-fixing forbs, BLNR forbs, NLNR forbs and NLR forbs respectively were leguminous forbs, broad-leaf-non-rhizomatous forbs, narrow-leaf-non-rhizomatous forbs, and narrow-leaf-rhizomatous forbs. Data are means of eight replicates. * indicates significant difference between the treatments of N addition with those of the control at $P < 0.05$.

Figure 3. Changes in aboveground biomass and species frequency of functional groups in responses to varying N-addition rates. The effect of N-addition rate on aboveground biomass of forbs (A), grasses and sedges (B). The effect N-addition rate on species frequency of forbs (C) grasses and sedges (D). Aboveground biomass and species frequency was determined in 1 m² quadrats and pooled over a 15-yr period (2004-2018). Curves represent significant nonlinear regression fits between aboveground biomass, species frequency and N-addition rate. The predictor variables are aboveground biomass changes and species frequency. The response variables are the logarithm of sum of 1 and N-addition rates (0, 1, 2, 4, 8, 16, 32, 64 g N m⁻² yr⁻¹). The significance of the nonlinear regression and correlation coefficients are shown for each relationship.

Figure 4. Different responses of functional groups to early period of N addition in Hulun Buir grassland in Inner Mongolia of China (119°35'-119°41' E, 48°27'-48°35' N). The treatments of N1, N2 and N3, respectively, represented N-addition rate being 3, 5 and 10 g N m⁻² yr⁻¹. Responses were calculated as the log response ratio: $\ln[\text{mean}(\text{biomass treatment})/\text{mean}(\text{biomass control})]$. The N₂-fixing forbs mainly included *Medicago ruthenica*, *Thermopsis lanceolata* and *Viciamul ticaulis*. The broad-leaf-non-rhizomatous (BLNR) forbs included *Potentilla acaulis*, *Potentilla tanacetifolia*, *Potentilla betonicifolia*, *Phlomis umbrosa*, and *Sibbaldia adpressa*. The narrow-leaf-non-rhizomatous (NLNR) forbs mainly included *Artemisia frigida*, *Filifolium sibiricum*, and *Serratula yamasutana*. Bunch grasses included *Stipa baicalensis*, *Achnatherum sibiricum* and *Cleistogenes squarrosa*. The rhizomatous monocots were *Carex duriuscula* and *Leymus chinensis*.

Figure 5 Effect of short-term nitrogen (N) addition on inorganic nitrogen concentrations in soils. Time-dependent changes in soil NH₄⁺-N (A), NO₃⁻-N (B) and total inorganic N (C) in response to varying N addition rates. The rates of N addition were 0 (control), 2 (N2), 8 (N8) g N m⁻² yr⁻¹. Data are means ± SE (n = 6). * indicates significant difference between the treatments of N addition and the control without N addition at $P < 0.05$.

Figure 6 Time-dependent changes in soil pH and metal ion concentrations in response to nitrogen (N) addition from 2004 to 2017. The rates of N addition were 0 (control) and 8 (+N) g N m⁻² yr⁻¹. Soil samples were collected in the middle of August each year. Data are the means ± SE (n=5 or 6). * indicates significant difference between the treatments of N addition and the control at $P < 0.05$.

Figure 7 Effect of lime addition and N addition from 1 to 3 years on aboveground biomass (AGB) of functional groups and soil pH. (A) AGB of the broad-leaf-non-rhizomatous (BLNR) forbs; (C) AGB of the narrow-leaf-non-rhizomatous (NLNR); (C) AGB of the bunch grasses; (D) soil pH. CK, control; L, lime addition; N, N addition;

NL, N plus lime addition. The levels of N and lime addition were $8 \text{ g N m}^{-2} \text{ yr}^{-1}$ and $50 \text{ g m}^{-2} \text{ yr}^{-1}$, respectively. Each treatment included six replicates. Two-way ANOVA was conducted to determine the effects of N addition and lime addition on aboveground biomass of forbs and grasses.

Figure 8 Effect of short-term nitrogen (N) addition on aboveground biomass of different functional groups and soil pH of grasslands in different regions. Top, Xinjiang; Middle, Qinghai; Bottom, Inner Mongolia. N1 and N2 refer to two levels of N-addition rate. In Xinjiang, N1 was $3 \text{ g N m}^{-2} \text{ yr}^{-1}$ and N2 was $9 \text{ g N m}^{-2} \text{ yr}^{-1}$. In Qinghai and Inner Mongolia, N1 and N2 were 2 and $8 \text{ g m}^{-2} \text{ yr}^{-1}$, respectively. The replicates of experimental plots were six in Xinjiang and Inner Mongolia, and five in Qinghai. Data are means $\pm$ SE. * indicates significant difference between the results of N-addition treatments and those of the control at $P < 0.05$. The same letters indicate insignificant difference between the results of N addition treatments at $P < 0.05$.

Figure 9 Illustration of linkage between the aboveground and underground processes to explain the phased responses of different functional groups to N addition. (A) Relative changes in aboveground biomass of BLNR forbs (broad-leaf-non-rhizome forbs), NLNR forbs (narrow-leaf-non-rhizome forbs) and bunch grasses in response to soil inorganic N levels during short-term N addition (<3 year) . (B) Relationships between relative changes in AGB of NLNR forbs and the bunch grasses and N addition-induced metal ion accumulation in soils by supplementing Mn similar to Mn level by N addition treatments in the field experiments. (C) Responses of expansion potential of bunch and rhizome grasses to inorganic N and metal ion accumulation in soils during the later period of N addition. (D) Changes in plant composition of community with prolongation of N addition. The loss sequence of functional groups at early, middle and later stage was N_2 -fixing forbs/BLNR forbs, NLNR forbs and bunch grasses. (E) Effect of N addition on inorganic N accumulation. (f) Effect of N addition on metal ion (Mn^{2+} , Cu^{2+} , Al^{3+}) concentrations in soils. (g) Effect of N addition on bud numbers in roots of grasses and rhizomes of rhizomatous grasses and sedges. Significant linear regression,

nonlinear regression and correlation coefficients are given in the figures.

Figure 10 Ecophysiological processes of different growth forms in response to N nutrient enrichment at different stages of N addition. (A) Effect of N addition on photosynthetic rates of grasses and narrow-leaf non-rhizomatous (NLNR) forbs. The broad-leaf-non-rhizomatous (BLNR) forbs, NLNR forbs and bunch grasses were the species, which mainly included *Potentilla acaulis*, *Potentilla tanacetifolia*, *Potentilla betonicifolia*, *Artemisia frigida*, *Cleistogenes squarrosa* and *Stipa krylovii*. (B) Effects of exogenous addition of metal (Mn^{2+}) on root growth of bunch grasses and narrow-leaf-non-rhizomatous (NLNR) forbs in pot experiments. Responses of root length were calculated as the log response ratio: $\ln[\text{mean}(\text{root length treatment})/\text{mean}(\text{root length control})]$. (C) At late stages, effect of N addition on biomass allocation of bunch grasses and rhizomatous monocots. Two dominant species of rhizomatous grass and sedge, *Leymus chinensis* and *Carex korshinskyi*, and two dominant species of bunch grass, *Stipa krylovii* and *Cleistogenes squarrosa*, were selected. The significance of the linear/nonlinear regression and correlation coefficients are shown for each relationship. FIG. 11 Time-dependent changes in plant composition of aboveground and corresponding drivers of underground. (A) Responses of aboveground biomass of different functional groups to N addition ($8 \text{ g N m}^{-2} \text{ yr}^{-1}$) over time. Changes in aboveground biomass of different functional groups with prolongation of N addition were analyzed by LOWESS method. (B) Changes in inorganic N changes in soils after N addition. Data are means of six replicates. (C) Effects of N addition at varying periods on concentrations of toxic metal Mn^{2+} in soils. Data are means of six replicates. (D) Effects of N addition up to 15 years on bud numbers in belowground. ***, **, and * indicate significant differences between the treatments of N addition with those of the control at $P < 0.001$, $P < 0.01$ and $P < 0.05$, respectively.

Figure 11 Time-dependent changes in plant composition of aboveground and corresponding drivers of underground. (A) Responses of aboveground biomass of different functional groups to N addition ($8 \text{ g N m}^{-2} \text{ yr}^{-1}$) over time. Changes in

1104 aboveground biomass of different functional groups with prolongation of N addition
1105 were analyzed by LOWESS method. (B) Changes in inorganic N changes in soils after
1106 N addition. Data are means of six replicates. (C) Effects of N addition at varying periods
1107 on concentrations of toxic metal Mn^{2+} in soils. Data are means of six replicates. (D)
1108 Effects of N addition up to 15 years on bud numbers in belowground. ***, **, and *
1109 indicate significant differences between the treatments of N addition with those of the
1110 control at $P<0.001$, $P<0.01$ and $P<0.05$, respectively.

1111

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 9

Figure 10

Figure 11

Fig. S1. Inter-annual variation in cover of forbs and grasses under the treatments of lime and N addition in the first three years. CK, control plots; L, lime addition plots; N, N addition plots; NL, both N and lime addition plots. The levels of N and lime addition were 8 g N m⁻² yr⁻¹ and 50 g m⁻² yr⁻¹. Each treatment included six replicates. Two-way ANOVA was conducted to determine the effects of N enrichment and lime addition on aboveground biomass and cover of forbs and grasses.

Fig. S2 Effect of lime addition and short-term N addition on aboveground biomass (AGB) of narrow-leaf-rhizomatous (NLR) forbs (A) and rhizome grasses and sedges (B). CK, control plots; L, lime addition plots; N, N addition plots; NL, both N and lime addition plots. The levels of N and lime addition were $8 \text{ g N m}^{-2} \text{ yr}^{-1}$ and $50 \text{ g m}^{-2} \text{ yr}^{-1}$, respectively. Each treatment included six replicates. Two-way ANOVA was conducted to determine the effects of N enrichment and lime addition on aboveground biomass and cover of forbs and grasses.

Fig. S3. Effect of different forms of N on leaf photosynthetic rates of a range of plant species. Data are means $\pm$ SE ($n > 15$). The leaf photosynthetic rates of different functional groups were measured after 7-10 d of N addition. The rates of N addition were 0 (Control) and 8 g N m⁻² yr⁻¹. Different letters indicate significant difference between the results of N addition treatments at $P < 0.05$.

Fig. S4 Different responses of the NLNR forb and the bunch grass to simulate Mn concentrations occurred in the field N addition experiment. The dominant species of NLNR forbs and bunch grasses were *Artemisia frigida* and *Stipa krylovii*. The plants were grown in soil that was collected from the control plots in the N addition experiment in field. The Mn²⁺ concentrations in soils were controlled to the levels of 20 mg kg⁻¹ (Mn 20) and 55 mg kg⁻¹ (Mn 55). The shoot and root biomass of these plants were measured after about two month. Data are means $\pm$ SE (n=6). Different letters indicate significant difference between the results of Mn addition treatments at $P < 0.05$.

Fig. S5. Different responses of forb and grass species to Mn addition under the conditions of other nutrients being sufficient. Shoot and root biomass of forbs and grasses were determined after exposure of the seedlings to nutrient solution containing 0 and 500 μM MnCl_2 . Data are means $\pm$ SE ($n=5$). * indicates significant difference between the results of Mn treatment with those of the control at $P < 0.05$.

Fig. S6. Different responses of the narrow-leaf-non-rhizomatous (NLNR) forb, *Artemisia frigida*, and the bunch grass, *Stipa krylovii*, to short-term N addition. Nitrogen levels were the same as those of N-addition experiment in the field. The replicates of each treatment were six. Shoot biomass, plant height, total root length and number of ramets and tillers were measured following treatment with different concentrations of N (0, 1, 2, 4, 8, 16, 32 g N m⁻²) for two month. Response ratios of biomass were calculated as the log response ratio: $\ln[\text{mean}(\text{biomass treatment})/\text{mean}(\text{biomass control})]$.

Fig. S7. Foliar nitrogen (N): phosphorus (P) ratios of forbs and grasses in response to N-addition rate. The forbs included dominant forb species of N₂-fixing forbs, broad-leaf-non-rhizomatous forbs (BLNR forbs), narrow-leaf-non-rhizomatous forbs (NLNR forbs) and narrow-leaf-rhizomatous forbs (NLR forbs). The grasses included dominant grass species of bunch grasses and rhizomatous grasses. Significant differences are denoted by different letters ($P<0.05$).

Table S1. Common plant species in the temperate grasslands in our studies of northern China. Aboveground biomass, occurrence frequency and seedling density were measured by a quadrat (1 m×1 m) from 2004 to 2018 in the experimental plots. Plant species of which frequency is more than 0.10 was considered to be a common species.

Functional groups	species	Biomass (g m ⁻²)	Frequency	Density (No. m ⁻²)
Leguminous forbs	<i>Astragalus galactites</i>	6.6	0.41	4.2
	<i>Astragalus scaberrimus</i>	5.5	0.21	7.1
	<i>Melissitus rutenica</i>	4.5	0.31	4.8
Broad leaf forbs	<i>Potentilla acaulis</i>	21.1	0.93	7.6
	<i>Potentilla tanacetifolia</i>	3.0	0.17	2.6
	<i>Potentilla verticillaris</i>	1.2	0.10	1.1
	<i>Potentilla betonicifolia</i>	0.7	0.12	1.8
	<i>Phlomis umbrosa</i>	3.5	0.15	3.9
	<i>Sibbaldia adpressa</i>	1.2	0.10	8.6
Narrow leaf forbs	<i>Artemisia frigida</i>	71.2	0.99	5.8
	<i>Allium bidentatum</i>	13.6	0.90	24.7
	<i>Allium ramosum</i>	3.4	0.37	10.4
	<i>Allium tenuissimum</i>	1.8	0.74	8.8
	<i>Allium neriniflorum</i>	0.6	0.65	15.9
	<i>Heteropappus altaicus</i>	2.0	0.43	3.8
	<i>Dianthus chinensis</i>	1.9	0.47	3.4
	<i>Thalictrum petaloideum</i>	1.5	0.29	2.0
	<i>Silene jensisensis</i>	2.8	0.23	3.8
	<i>Artemisia scoparia</i>	4.9	0.10	5.3
	<i>Artemisia pubescens</i>	8.5	0.10	8.5
Deep rhizome forbs	<i>Potentilla bifurca</i>	5.7	0.72	11.2
Annual/biennial forbs	<i>Dontostemon dentatus</i>	3.0	0.31	10.2
	<i>Gentiana squarrosa</i>	0.5	0.18	10.1
	<i>Chenopodium aristatum</i>	0.7	0.13	3.0
	<i>Salsolacollina pall</i>	0.3	0.10	2.3
Bunch graminoids	<i>Stipa krylovii</i>	114.1	0.99	66.1
	<i>Cleistogenes squarrosa</i>	8.4	0.99	28.2
	<i>Poa annua</i>	8.2	0.25	2.6
Rhizomatous graminoids	<i>Carex korshinskyi</i>	10.3	0.96	50.2
	<i>Agropyron cristatum</i>	19.8	0.59	18.4
	<i>Leymus chinensis</i>	16.4	0.29	20.4

Table S2. Rare plant species in the temperate grasslands in our studies of northern China. Aboveground biomass, occurrence frequency and seedling density were measured by a quadrat (1 m×1 m) from 2004 to 2018 in our experimental plots. Plant species of which frequency is less than 0.10 was taken as a rare species.

Functional groups	species	Biomass (g m ⁻²)	Frequency	Density (No. m ⁻²)
Perennial forbs	<i>Oxytropis leptophylla</i>	0.3	0.07	1.3
	<i>Astragalus adsurgens</i>	0.01	0.01	0.1
	<i>Gentiana deburica</i>	2.2	0.05	3.0
	<i>Potentilla multifida</i>	0.4	0.03	0.6
	<i>Iris tenuifolia</i>	1.6	0.04	2.4
	<i>Convolvulus ammannii</i>	0.2	0.01	0.1
	<i>Artemisia gmelinii</i>	0.3	0.01	0.1
	<i>Cymbaria dahurica</i>	1.8	0.04	2.4
	<i>Scorzonera austriaca</i>	0.2	0.03	0.4
	<i>Orosta.chys fimbriata</i>	0.1	0.04	2.0
	<i>Thalictrum squarrosum</i>	0.3	0.02	0.5
	<i>Gueldenstaedtia verna</i>	1.2	0.02	0.9
	<i>Allium wallichii</i>	0.9	0.04	0.8
	<i>Stellaria petraea</i>	3.9	0.08	1.3
	<i>Androsace umbellate</i>	1.0	0.07	1.4
	<i>Tephroseris kirilowii</i>	0.1	0.01	0.3
Annual/biennial forbs	<i>Chenopodium glaucum</i>	0.8	0.06	3.0
	<i>Lepidum apetalum</i>	0.01	0.01	0.1
Perennial graminoids	<i>Koeleria glauca</i>	6.8	0.02	1.3
Annual/biennial graminoids	<i>Setaria viridis</i>	0.2	0.01	0.3