The role of central nervous system mechanisms in resistant hypertension

Dagmara Hering¹, Markus Schlaich¹

¹School of Medicine and Pharmacology - Royal Perth Hospital Unit, The University of Western Australia

Address for correspondence:

Dr habil. Dagmara Hering, MD

School of Medicine and Pharmacology - Royal Perth Hospital Unit

The University of Western Australia

Level 3 MRF Building, Rear 50 Murray Street, Perth WA 6000 MDBP: M570

P +61 8 9224 0316

F +61 8 9224 0374

E dagmara.hering@uwa.edu.au

Abstract

Arterial hypertension remains a primary global health problem with significant impact on cardiovascular morbidity and mortality. The low rate of hypertension control and failure to achieve target blood pressure levels particularly among high risk patients with resistant hypertension has triggered renewed interest in unraveling the underlying mechanisms to implement therapeutic approaches for better patient management. Here we summarize the crucial role of neurogenic mechanisms in drug-resistant hypertension, with a specific focus on central control of blood pressure, the factors involved in central integration of afferent signaling to increase sympathetic drive in resistant hypertension, and briefly review recently introduced interventional strategies distinctively targeting sympathetic activation.

Key words: resistant hypertension, central sympathetic nervous system, noradrenaline spillover, muscle sympathetic nerve activity, neural reflex mechanisms, oxidative stress, therapeutic intervention

Role of the sympathetic nervous system in human hypertension

The sympathetic nervous system (SNS) is activated in human hypertension, with an increase in sympathetic outflow to the kidney, the heart and the skeletal muscle vascular bed as convincingly demonstrated by the use of isotope dilution method for quantifying noradrenaline (NA) spillover rates and postganglionic efferent sympathetic nerve recording with microneurography. NA spillover has allowed for assessment of regional differentiation of sympathetic outflow. While the rate of overflow of NA into plasma has been documented for most internal organs, neurotransmitter release from the heart and the kidney are most relevant in initiating and maintaining arterial hypertension [1, 2]. Increased NA release from the renal sympathetic nerves is present in patients with untreated essential hypertension (EH), particularly below the age of 40 [3] and is a primary trigger for BP elevation [4]. Augmented cardiac NA spillover and reduced neuronal NA reuptake further contribute to sympathetic activation in EH [5].

In the healthy heart the amount of adrenaline release from the sympathetic nerves is negligible as the majority (80%) of adrenaline is secreted from the adrenal medulla into the blood stream, with only a small amount released from sympathetic nerve endings. However, in patients with untreated EH, adrenaline release has been reported from cardiac sympathetic nerves providing a possible additional mechanisms of potentiated NA release through adrenaline co-transmission [6]. Additionally, stress *per se* induces adrenaline plasma release and is likely to enhance neuronal uptake of adrenaline in cardiac sympathetic nerve endings leading to discharge of adrenaline as co-transmitter supporting the 'adrenaline hypothesis' in EH [6].

In fact, patients with uncontrolled hypertension are at high risk for developing heart disease and kidney disease with the progression to heart failure (HF) and kidney failure over time. Accordingly, the prognosis in HF has been directly linked to the magnitude of NA release

from the cardiac sympathetic nerves [7], while increased NA release from the renal sympathetic nerves has been shown to be a strong independent predictor of all-cause mortality and heart transplantation in HF patients [8]. Clearly, elevated cardiac and renal sympathetic tone are core features of human hypertension and associated with disease progression and mortality. High levels of cardiac and renal NA spillover accompanying increased muscle sympathetic nerve activity (MSNA) measured by microneurography have been related to heart damage including left ventricular (LV) hypertrophy [9] and LV dysfunction [10] in untreated EH. However, it is not only elevated blood pressure (BP) and hypertension-related ventricular remodelling that is characterized by sympathetic activation, increased rate of sympathetic nerve firing is evident in low risk subjects with high-normal BP [11], EH [5] with a further 2-3 fold increase in patients with true resistant hypertension (RH) irrespective of the combination of multi drug-therapy which supposed to oppose chronic sympathetic outflow [12-14]. Indeed, our own experience indicates that patients with RH are characterized by high levels of MSNA when compared to healthy controls and EH patients, with burst activity evident with every heartbeat, often superimposed within one cardiac cycle (*sharp “M” shape burst*) due to high sympathetic nerve firing. While there is an increased incidence of sympathetically-mediated associated co-morbidities in patients with RH including diabetes, CKD, obesity and obstructive sleep apnoea (OSA) [15], the mechanism implicated in drug resistance are not fully understood. Sympathetic excitation is primarily derived from specific autonomic brain regions and/or impaired peripheral neural reflex mechanisms including arterial baroreceptors, arterial chemoreceptors and cardiopulmonary mechanoreceptors. The regulatory process underlying sympathetic activation in RH remains uncertain, though it is likely to involve an imbalance between catecholaminergic neurons widely distributed in the brain stem and possible loss of inhibition transmitted from non-catecholaminergic structures in the rostral and caudal ventrolateral medulla.

Despite substantial progress in the management of hypertension and a better understanding of the mechanisms controlling BP, one in every ten adults is likely to fail in achieving target BP with antihypertensive pharmaco-therapy with true RH warranting alternative treatment options to minimize associated cardiovascular (CV) risk [16, 17]. Of clinical relevance, therapeutic approaches aimed at SNS inhibition via modulation of interrelated pathways appears pivotal for attaining BP control and reducing adverse complications associated with chronic elevation of sympathetic drive in neurogenic hypertension.

Role of autonomic brain regions in sympathetic and blood pressure control

Tonic sympathetic activation and tonic arterial pressure control critically depend on central integrative structures in the brain stem, specifically the rostral ventrolateral medulla (RVLM) (Figure 1). Descending projections to the RVLM arise among others from the neurons in the peri-aqueductal grey (PAG) and hypothalamic paraventricular nucleus (PVN). The RVLM integrates reflex neural mechanisms from arterial baroreceptors, chemoreceptors and various afferent sensory visceral receptors via direct connection with the upper part of the medulla through the nucleus tractus solitarius (NTS) and PVN which modulate vasomotor sympathetic nerve discharge and BP. Evidence from experimental models has indicated that up to 70% of spinally projecting RVLM neurons are sympathoexcitatory C1 neurons. These play a critical role in tonic drive and reflex control of sympathetic vasomotor activity to the preganglionic motor neurons of the spinal cord, thereby directly modulating peripheral SNS activity. Consequently, chemical, electrical or photo stimulation of most of the RVLM neurons increases BP, which subsequently decreases with experimental cooling, destruction of the region or GABA agonists injection [18-21]. Under physiological conditions, arterial baroreceptors play a fundamental role in preventing excessive variation in BP levels. Afferent signals from baroreceptors stimulate the NTS in the upper part of the medulla in response to the distension of the vessel wall caused by transmural pressure. Subsequently, a signal arising

from the NTS exerts a (1) parasympathetic vagal effect resulting in slowing HR and (2) direct activation of the lower caudal ventrolateral medulla (CVLM) (3) and reduced tonic sympathetic activity generated in the RVLM. Depressor neurons of the CVLM provide direct tonic inhibition to the RVLM neurons (Figure 1). In this context, a reduction in tonic sympathoinhibitory function of the CVLM to RVLM is likely to contribute to the development of hypertension. In fact the parasympathetic cardiac component of the baroreflex is impaired in patients with hypertension and a family history of hypertension [22]. While the lack of activity of depressor neurons of the CVLM on the RVLM has been shown to be attenuated in spontaneous hypertensive rats and renovascular hypertensive rats [23, 24], the activity of CVLM and RVLM neurons in human arterial hypertension has not been tested directly.

Central nervous system (CNS) NA turnover in humans can be estimated from the combined overflow of brain NA and its lipophilic metabolites into the internal jugular veins [25, 26]. Indeed, direct blood sampling from the internal jugular veins and simultaneous cerebral blood flow scans revealed that CNS mechanisms involving suprabulbar noradrenergic projections from the brain stem to the hypothalamus play an important role in sympathetic activation in untreated EH [27]. Augmented NA release from subcortical brain regions but not cerebral cortex corroborates the critical contribution of sympathoexcitatory and pressor noradrenergic brain neurons to elevated BP in patients with EH [25], mediating increased sympathetic outflow to the periphery. Subcortical NA turnover in brain regions is significantly higher in EH when compared to healthy subjects and directly related to MSNA in healthy subjects, total body NA spillover in normal and hypertensive subjects combined, and renal NA spillover in EH [25, 27]. Notably, amongst noradrenergic neurons in the brain stem, the locus coeruleus (LC) is the largest source of NA production [28, 27]. In this context the role of kidney in centrally-mediated cardiovascular regulation is of significant relevance. The kidney

has a dense network of chemoreceptors and baroreceptors that transmit afferent signals to the brain stem. These afferent signals originating from unmyelinated renal sensory fibres traveling along arteries, enter the dorsal roots and project to spinal and supra spinal neurons and the hypothalamus causing increase in BP and sympathetic outflow to peripheral organs [29, 30]. A series of experimental studies showed that kidney failure following 5/6 nephrectomy in rats causes subsequent hypertension through sympathetic activation. Dorsal rhizotomy in the dorsolateral part of the thoraco-lumbar vertebral bodies (T10-L2) led to a significant BP reduction and prevented an increase in NA turnover rate in the LC (A6), and posterior and lateral hypothalamus where the majority of brain NA producing neurons reside [31].

In humans, bilateral nephrectomy reduced or “normalized” MSNA in hemodialysis patients [32] and renal transplant recipients [33]. Furthermore bilateral nephrectomy improved BP control in hemodialysis patients with RH [34]. This observation affirms that afferent renal sensory fibres originating from the failing kidney cause sympathetic activation and BP elevation [32-34].

Role of central processes in sympathetic outflow

The association between obstructive sleep apnea (OSA) and hypertension is well recognized [16]. Accordingly, OSA is associated with augmented sympathetic excitation contributing to neurogenic hypertension. Very recently, a study using functional Magnetic Resonance Imaging (MRI) assessed brain activity and regional grey matter changes in OSA patients with simultaneous MSNA recordings [35]. In comparison to healthy controls, high levels of MSNA in OSA patients appear to be associated with functional (not anatomical) changes within the higher cortical and subcortical brain regions which are involved in the modulation of sympathetic outflow and BP via the brainstem regulatory nuclei. Increased signal intensity in the medial prefrontal cortex and covariation with MSNA levels in OSA patients when

compared to controls [35] may have important implications for understanding the mechanisms involved in reduced baroreflex gain in OSA and RH. Accordingly, the medial prefrontal cortex has projections to the NTS and RVLM which are responsible for sympathetic and BP regulation [35].

In view of recent meta-analysis demonstrating favorable effects of renal denervation (RDN) in reducing apnea-hypopnea index and BP in OSA patients [36], it is reasonable to speculate that interrupting afferent signaling from the kidney to brain stem via renal nerve ablation may reverse altered functional brain changes thereby improving outcomes in OSA patients. At this stage the functional changes in blood oxygenation and flow in response to brain neural activity in RH patients remain unknown. Likewise, the potential impact of therapeutic interventional strategies opposing chronic sympathetic drive on brain functional imaging in RH need to be determined.

Role of renal denervation

In view of the well established contribution of the SNS and renal nerves to the initiation, development and maintenance of elevated BP, the interruption of renal sympathetic and afferent nerves appears as a reasonable treatment approach to improve BP control in RH and renal failure patients. Accordingly, RDN has improved BP control over the longer term [37, 38] and has been introduced to clinical practice [16]. The recent Symplicity HTN-3 study was unable to confirm a BP lowering effect above that of the sham control [39] although it has been questioned whether RDN was actually achieved in this study [40]. In an initial proof-of-concept pilot study of a total of 12 patients with end stage renal disease (ESRD) and uncontrolled BP [41], 8 out of 9 patients from the study cohort who had eligible anatomy of renal arteries to ablate the nerves experienced a substantial improvement in BP control. The reduction in renal NA spillover directly after bilateral RDN and in MSNA at follow-up was observed in the 2 patients who also underwent assessment of sympathetic activation [41]. The

reduction of whole body NA spillover, renal NA spillover and postganglionic efferent multi-unit MSNA in addition to BP improvement has also been demonstrated in patients with RH in the short-term [42, 12, 13] and longer term [14] out to 12 months post RDN. Interestingly, the reduction in multi-unit MSNA after RDN was accompanied by even more pronounced reduction in all properties of single active vasoconstrictors neurons including firing rate, firing probability and the incidence of multiple spikes within one cardiac cycle which may be reflective of central mechanisms integrating afferent signalling from the kidneys and efferent sympathetic outflow resulting in sympathetic inhibition and BP control [12].

Additional relevant findings involving brain noradrenergic neurons, hypothalamic and amygdala projections in behavioral responses mediated autonomic activation were observed after RDN [43, 44]. Patients with RH experienced a significant improvement in health related quality of life aspects including the mental components (i.e. the vitality, social function, role emotion, mental health). These effects were unrelated to the BP lowering effect post procedure. Additionally, symptoms indicative of depression including sadness, tiredness and libido were considerably attenuated with RDN [43]. The potential impact of RDN to favorably modulate brain regions involved in cognitive processing was confirmed in another study demonstrating a substantial reduction in anxiety, depressive symptoms, the intensity of headache and stress tolerance after the procedure [44].

Role of neurovascular compression in resistant hypertension

Pulsatile vascular compression between arterial and neural tissue in the medulla has been shown to be implicated in sympathetically-mediated hypertension. Neurovascular compression (NVC) related to hypertension results from the compression and subsequent irritation of the left RVLM regions at the levels of the C1 neurons and/or the root entry zone of cranial nerves by looping arteries, mostly posterior inferior cerebellar artery (PICA) and/or vertebral artery [18]. Among several proposed theories for the association between NVC on

the left side and BP rise, oxygen deficiency resulting from the compression of nerve IX and X is likely to cause deafferentation of the NTS leading to loss of inhibitory effect transmitted from the NTS and sympathoexcitation of C1 neurons. While NVC may potentially involve various cranial nerves in that area (i. e. V, VII, VIII, XII), compression of the cranial glossopharyngeal and vagus nerves appear to predominantly contribute to BP elevation. Visceral afferent fibres of the IX nerve provide sensory information from the carotid body baroreceptors to the NTS while a signal arising from the aortic body baroreceptors is transmitted via X afferent fibres modulating BP control. While screening for posterior fossa NVC in the management of RH is not routinely performed, it should be noted that neurovascular pulsatile compression, commonly present on the left side at the RVLM has been documented in patients with RH. Previous studies have demonstrated a substantial decrease in BP and antihypertensive regimens in 7 out of 8 patients with uncontrolled BP 3 months following microvascular decompression (MVD) of IX and X cranial nerves root-entry zone causing pulsatile neurovascular compression, with sustained BP reduction in 4 patients who were followed out to 12 months post procedure [45]. Additional interesting observation for a beneficial effect of a single MVD comes from a case report of a 72-year old women demonstrating typical clinical symptoms including trigeminal neuralgia, hemifacial spasm and tinnitus in addition to hypertension and paroxysmal supraventricular tachycardia (PSVT) caused by vertebral artery and subsequent compression of cranial nerves V, VII, VIII, IX and X. Of note, decompression of the posterior cranial fossa alleviated all symptoms immediately after the surgery with corresponding BP improvement and the absence of PSVT out to 8 months post procedure [46]. Further proof for a favourable BP lowering effect and improved sensitivity to antihypertensive medication after successful neurosurgical decompression of the brain stem has been documented retrospectively in 9 patients with severe primary hypertension not responding to conventional drug therapy and 7 non-operated

patients with neurovascular compression who were followed over 2 years period [47]. While MVD resulted in a significant and permanent fall in BP accompanied by an improved response to medication over 2 years following the procedure, there were no noticeable changes in BP in controls at that point in time [47]. Additional retrospective analysis of 13 patients with RH revealed that MVD independently of the types of NVC (brain stem and/or the nerves IX-X) and/or side (left and/or right) led to a reduction in BP in all types of NVC, with a more pronounced BP lowering effect when NVC affected both the brain stem and the nerves on the left side [48]. Whether NVC of its origin is a leading contributor to sympathetic activation and thereby causing BP elevation or remains a secondary cause of HT is debated. Nevertheless, studies based on microneurography have shown that NVC is associated with augmented MSNA in both normotensive and hypertensive subjects [49]. Moreover, the prevalence of NVC and the magnitude of MSNA was greater in EH when compared to normal BP and high-normal BP suggesting that pulsatile compression may play a critical role in triggering sympathetic activation and may precede the development of hypertension [49]. Further observation has shown that only hypertensive patients with signs of NVC of the RVLM on the MRI have high MSNA but not hypertensives without deformation or only vascular contact visible on the MRI [50]. While MVD results in sustained reduction in BP, MSNA and medication use from baseline to 12 month follow-up in RH, surprisingly it appeared that resting MSNA returned to the baseline BP and MSNA levels from 18 to 24 months post procedure [51]. Whether the rebound of hypertension and sympathetic activation in RH is associated with counter-regulatory neurohumoral mechanisms, increased hyper-responsiveness of beta-adrenergic receptors during sympathetic inhibition, occurrence of other sympathetically-mediated co-morbidities or long history of primary hypertension requires further investigation.

In view of available data, NVC of the RVLM by a looping artery could be a potential underpinning cause of drug RH when BP remains elevated despite multi-drug antihypertensive combination therapy, medication adherence and patient compliance. Furthermore, NVC associated with hypertension is often not accompanied by clinical symptoms such as cranial rhizopathies. At this stage there is no supportive evidence for MVD in long term BP control in RH. Nevertheless, therapy with clonidine appears to be a useful approach in NVC associated with drug-resistance hypertension and CKD [52] and should perhaps be offered to patients until a suitable group of RH patients who may benefit from the MVD procedure can be identified.

Deep brain stimulation

Deep brain stimulation (DBS) is a promising interventional therapy designed to alter pathological and sympathetic activity within the CNS and has gained significant recognition in the treatment of Parkinson's disease. In addition to a wide range of neurological disorders, DBS of the ventrolateral (PAG)/periventricular grey matter has been successfully demonstrated in RH [53, 54]. While the central grey is the primary control centre for descending pain modality, it also sends descending projections to the RVLM. DBS was initially performed to treat chronic central pain syndrome unresponsive to pain-relief drugs, however it unexpectedly produced a sustained BP reduction in the two patient cases. While the effect of DBS of MSNA in RH is unknown, more recently acute DBS of specific midbrain nuclei and ventrolateral PAG improved acutely vasomotor baroreflex sensitivity, reduced MSNA and BP in patients with chronic neuropathic pain and Parkinson's disease [55]. Although costly and associated with a potential 1% stroke risk, DBS appears to be an alternative therapeutic approach for treating severe forms of uncontrolled hypertension and/or possibly patients unresponsive to other currently available interventional strategies. However,

whether BP control achieved with DBS may improve patient outcomes needs to be investigated further.

Baroreceptor stimulation

Arterial baroreceptors are an important centrally-mediated mechanism involved in the short-term BP regulation. However, in the presence of sustained BP elevation, baroreflex control becomes less sensitive in mediating changes in sympathetic activity to the heart and blood vessels. In this context, prolonged electric stimulation of baroreflex afferent nerves has been shown to be associated with a marked BP lowering effect via baroreceptor-induced central sympathoinhibition [56]. Although in humans the role of baroreflex control in long-term regulation is not entirely understood, recent data from electrical stimulation of carotid sinus baroreceptors via an implantable device has provided novel insights into baroreflex physiology in RH patients. Indeed the safety and efficacy of device-based chronic baroreflex activation therapy (BAT) with the CVRx Rheos System (DEBuT-HT Trial) was demonstrated in high CV risk patients with RH. The magnitude of the fall in BP 3 months following device implementation was maintained in patients who were followed out to 53 months post procedure [57, 58]. While safe and effective in terms of BP control, BAT with the CVRx Rheos System was associated with serious procedure-related adverse events and short-term battery life when compared to new generation device, the BAROSTIM neo which produced a significant BP reduction at 3 and 6 month follow-up, even in patients previously treated with RDN, with less device-related side effects [59]. Regarding sympathetic activity, the acute effect of BAT has only been determined in 12 patients with RH indicating a rapid reduction in BP and MSNA when the stimulator was switched on, returning to baseline level when switched off [60]. In view of the dominant role of the kidney in long-term BP and MSNA improvement [14], further clinical studies need to determine the applicability of BAT on hypertension-induced end organ damage and neural function.

Role of oxidative stress in central brainstem

Oxidative stress is implicated in the progression of CV disease. There is also supportive evidence from numerous experimental and clinical studies linking oxidative stress to the development and maintenance of hypertension [61]. Recently, increased oxidative stress has been suggested to promote transition from pre-hypertension to established hypertension [62], while therapy with nebivolol via inducing endothelium-dependent vasodilation via nitric oxide (NO)-pathway led to a substantial increase in nitric oxide production in pre-hypertension and arterial hypertension [63]. While the mechanisms inducing the reduction in NO, bioavailability or impairment NO-mediated vasodilation in hypertension are not entirely understood, supportive evidence from experimental models is indicative of a bidirectional interaction between NO and SNS in modulating BP levels [61]. Accordingly, central production of oxidative stress in autonomic brain regions, particularly in the RVLM increases central sympathetic activation thereby triggering BP elevation [64]. In this context, a better understanding of brain mechanisms linking oxidative stress to human neurogenic hypertension may help to implement novel therapies for drug-RH.

Conclusions

Arterial hypertension remains a growing and challenging clinical problem. Special attention should be given to drug- RH affecting ~10% of hypertensive adults. Currently available clinical interventional therapeutic strategies that have the potential to favourably modulate autonomic neural mechanisms such as RDN and carotid baroreceptor stimulation, if performed by experienced operators, can be offered to RH patients in the case of ineffective drug therapy. Given the importance of neurogenic hypertension, unravelling the CNS mechanisms implicated in human RH is likely to have a substantial impact in the application of specific treatment approaches for attaining BP control and reducing adverse complications associated with chronically elevated sympathetic drive.

DISCLOSURE

Dr Schlaich is supported by career fellowships from the NHMRC, is an investigator in studies sponsored by Medtronic, serves on scientific advisory boards for Abbott (formerly Solvay) Pharmaceuticals, BI, Novartis Pharmaceuticals, BI and Medtronic and has received honoraria and travel support from Abbott, BI, Servier, Novartis and Medtronic. The laboratories of Dr Schlaich receive research funding from Medtronic, Abbott Pharmaceuticals, Otsuka and Servier Australia.

Figure 1.

Central tonic sympathetic outflow and blood pressure (BP) control in central integrative structures in the brain stem; the upper part of the medulla, rostral ventrolateral medulla (RVLM) and lower part of the medulla, caudal ventrolateral medulla (CVLM).

Parasympathetic outflow to the heart arises mainly from neurons in the nucleus ambiguous (AMB). Descending projections to the RVLM from the peri-aqueductal grey (PAG) and hypothalamic paraventricular nucleus (PVN). Projections from the amygdala (A) and locus coeruleus (LC) involved in behavioural, psychological and cognitive control.

References

Papers of particular interest, published recently, have been highlighted as:

•• Of major importance

1. •• Esler M, Lambert G, Jennings G. Regional norepinephrine turnover in human hypertension. Clinical and experimental hypertension Part A, Theory and practice. 1989;11 Suppl 1:75-89. *This study presents first evidence for organ-specific regional differentiation of sympathetic outflow indicating that increased NA release from the heart and kidney are pivotal in human essential hypertension.*
2. Esler M. The 2009 Carl Ludwig Lecture: pathophysiology of the human sympathetic nervous system in cardiovascular diseases: the transition from mechanisms to medical management. Journal of Applied Physiology. 2010;108(2):227-37. doi:DOI 10.1152/jappphysiol.00832.2009.
3. •• Esler M, Lambert G, Jennings G. Increased Regional Sympathetic Nervous Activity in Human Hypertension - Causes and Consequences. Journal of Hypertension. 1990;8:S53-S7. *The importance of increased renal NA spillover to the pathophysiology of essential hypertension.*
4. Parati G, Esler M. The human sympathetic nervous system: its relevance in hypertension and heart failure. European Heart Journal. 2012;33(9):1058-U39. doi:DOI 10.1093/eurheartj/ehs041.
5. Schlaich MP, Lambert E, Kaye DM, Krozowski Z, Campbell DJ, Lambert G et al. Sympathetic augmentation in hypertension - Role of nerve firing, norepinephrine reuptake, and angiotensin neuromodulation. Hypertension. 2004;43(2):169-75. doi:Doi 10.1161/01.Hyp.0000103160.35395.9e.
6. •• Rumantir MS, Jennings GL, Lambert GW, Kaye DM, Seals DR, Esler MD. The 'adrenaline hypothesis' of hypertension revisited: evidence for adrenaline release from

- the heart of patients with essential hypertension. *Journal of Hypertension*. 2000;18(6):717-23. doi:Doi 10.1097/00004872-200018060-00009. *First study indicating augmented adrenaline release from cardiac sympathetic nerves and the relevant contribution of adrenaline co-transmission in triggering noradrenaline release in essential hypertension development.*
7. •• Kaye DM, Lefkovits J, Jennings GL, Bergin P, Broughton A, Esler MD. Adverse Consequences of High Sympathetic Nervous Activity in the Failing Human Heart. *Journal of the American College of Cardiology*. 1995;26(5):1257-63. doi:Doi 10.1016/0735-1097(95)00332-0. *This study demonstrated that increased cardiac noradrenaline spillover predicted poor CV outcomes in heart failure patients.*
 8. •• Petersson M, Friberg P, Eisenhofer G, Lambert G, Rundqvist B. Long-term outcome in relation to renal sympathetic activity in patients with chronic heart failure. *European Heart Journal*. 2005;26(9):906-13. doi:DOI 10.1093/eurheartj/ehi184. *This study documented the association of increased renal noradrenaline spillover and all-cause mortality, and heart transplantation in heart failure patients.*
 9. Schlaich MP, Kaye DM, Lambert E, Sommerville M, Socratous F, Esler MD. Relation between cardiac sympathetic activity and hypertensive left ventricular hypertrophy. *Circulation*. 2003;108(5):560-5. doi:Doi 10.1161/01.Cir.0000081775.72651.B6.
 10. Grassi G, Seravalle G, Quarti-Trevano F, Dell'Oro R, Arenare F, Spaziani D et al. Sympathetic and baroreflex cardiovascular control in hypertension-related left ventricular dysfunction. *Hypertension*. 2009;53(2):205-9. doi:10.1161/HYPERTENSIONAHA.108.121467.
 11. Hering D, Kara T, Kucharska W, Somers VK, Narkiewicz K. High-normal blood pressure is associated with increased resting sympathetic activity but normal responses

- to stress tests. Blood Pressure. 2013;22(3):183-7. doi:Doi 10.3109/08037051.2012.759689.
12. •• Hering D, Lambert EA, Marusic P, Walton AS, Krum H, Lambert GW et al. Substantial Reduction in Single Sympathetic Nerve Firing After Renal Denervation in Patients With Resistant Hypertension. Hypertension. 2013;61(2):457-+. doi:Doi 10.1161/Hypertensionaha.111.00194. *First study documenting resting multi-unit and single-unit muscle sympathetic nerve activity in patients with resistant hypertension and the effect of renal denervation on sympathetic nerve firing.*
 13. Hering D, Lambert EA, Marusic P, Ika-Sari C, Walton AS, Krum H et al. Renal nerve ablation reduces augmentation index in patients with resistant hypertension. Journal of Hypertension. 2013;31(9):1893-900. doi:Doi 10.1097/Hjh.0b013e3283622e58.
 14. •• Hering D, Marusic P, Walton AS, Lambert EA, Krum H, Narkiewicz K et al. Sustained Sympathetic and Blood Pressure Reduction 1 Year After Renal Denervation in Patients With Resistant Hypertension. Hypertension. 2014;64(1):118-24. doi:Doi 10.1161/Hypertensionaha.113.03098. *First study demonstrating the sustained effect of renal denervation on blood pressure and sympathetic nerve firing out to 12 months post procedure.*
 15. Calhoun DA, Jones D, Textor S, Goff DC, Murphy TP, Toto RD et al. Resistant hypertension: Diagnosis, evaluation, and treatment - A scientific statement from the American Heart Association Professional Education Committee of the Council for High Blood Pressure Research (Reprinted from Hypertension, vol 51, pg 1403-1419, 2008). Circulation. 2008;117(25):E510-E26. doi:Doi 10.1161/Circulationaha.108.189141.
 16. Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Bohm M et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension

- (ESH) and of the European Society of Cardiology (ESC). *J Hypertens.* 2013;31(7):1281-357. doi:10.1097/01.hjh.0000431740.32696.cc.
17. Judd E, Calhoun DA. Apparent and true resistant hypertension: definition, prevalence and outcomes. *J Hum Hypertens.* 2014;28(8):463-8. doi:10.1038/jhh.2013.140.
 18. •• Jannetta PJ, Segal R, Wolfson SK. Neurogenic Hypertension - Etiology and Surgical-Treatment .1. Observations in 53 Patients. *Ann Surg.* 1985;201(3):391-8. doi:Doi 10.1097/00000658-198503000-00023. *Study demonstrating the contribution of pulsatile compression of the nerve at the level of medulla oblongata to neurogenic hypertension and the potential for microvascular decompression in the treatment of associated hypertension.*
 19. Naraghi R, Gaab MR, Walter GF, Kleineberg B. Arterial-Hypertension and Neurovascular Compression at the Ventrolateral Medulla - a Comparative Microanatomical and Pathological-Study. *J Neurosurg.* 1992;77(1):103-12. doi:DOI 10.3171/jns.1992.77.1.0103.
 20. Jannetta PJ, Gendell HM. Clinical observations on etiology of essential hypertension. *Surgical forum.* 1979;30:431-2.
 21. Amano M, Kubo T. Involvement of Both Gaba-a and Gaba-B Receptors in Tonic Inhibitory Control of Blood-Pressure at the Rostral Ventrolateral Medulla of the Rat. *N-S Arch Pharmacol.* 1993;348(2):146-53. doi:Doi 10.1007/Bf00164791.
 22. Parmer RJ, Cervenka JH, Stone RA. Baroreflex Sensitivity and Heredity in Essential-Hypertension. *Circulation.* 1992;85(2):497-503.
 23. Smith JK, Barron KW. The Rostral and Caudal Ventrolateral Medulla in Young Spontaneously Hypertensive Rats. *Brain research.* 1990;506(1):153-8. doi:Doi 10.1016/0006-8993(90)91213-Z.

24. Colombari E, Sato MA, Cravo SL, Bergamaschi CT, Campos RR, Lopes OU. Role of the medulla oblongata in hypertension. *Hypertension*. 2001;38(3):549-54.
25. •• Ferrier C, Jennings GL, Eisenhofer G, Lambert G, Cox HS, Kalff V et al. Evidence for Increased Noradrenaline Release from Subcortical Brain-Regions in Essential-Hypertension. *Journal of Hypertension*. 1993;11(11):1217-27. *First study demonstrating increased brain noradrenaline turnover in the pathophysiology of human essential hypertension.*
26. Lambert GW, Ferrier C, Kaye DM, Kalff V, Kelly MJ, Cox HS et al. Monoaminergic neuronal activity in subcortical brain regions in essential hypertension. *Blood Press*. 1994;3(1-2):55-66.
27. Esler M, Lambert G, Vaz M, Thompson J, Kaye D, Kalff V et al. Central nervous system monoamine neurotransmitter turnover in primary and obesity-related human hypertension. *Clinical and Experimental Hypertension*. 1997;19(5-6):577-90. doi:10.3109/10641969709083171.
28. Foote SL, Bloom FE, Astonjones G. Nucleus Locus Ceruleus - New Evidence of Anatomical and Physiological Specificity. *Physiol Rev*. 1983;63(3):844-914.
29. Stella A, Zanchetti A. Functional-Role of Renal Afferents. *Physiol Rev*. 1991;71(3):659-82.
30. DiBona GF, Kopp UC. Neural control of renal function. *Physiol Rev*. 1997;77(1):75-197.
31. •• Campese VM, Kogosov E. Renal Afferent Denervation Prevents Hypertension in Rats with Chronic-Renal-Failure. *Hypertension*. 1995;25(4):878-82. *First study demonstrating the contribution of afferent renal nerves from the failing kidney to the brain stem resulting in blood pressure elevation and sympathetic activation.*

32. •• Converse RL, Jacobsen TN, Toto RD, Jost CMT, Cosentino F, Fouadtarazi F et al. Sympathetic Overactivity in Patients with Chronic-Renal-Failure. *New Engl J Med.* 1992;327(27):1912-8. doi:Doi 10.1056/Nejm199212313272704. *First study demonstrating the effect of nephrectomy on sympathetic nerve firing in chronic renal failure indicating that sympathetic excitation is associated with afferent signals arising from the failing kidney.*
33. Hausberg M, Kosch M, Harmelink P, Barenbrock M, Hohage H, Kisters K et al. Sympathetic nerve activity in end-stage renal disease. *Circulation.* 2002;106(15):1974-9. doi:Doi 10.1161/01.Cir.0000034043.16664.96.
34. •• Zazgornik J, Biesenbach G, Janko O, Gross C, Mair R, Brucke P et al. Bilateral nephrectomy: The best, but often overlooked treatment for refractory hypertension in hemodialysis patients. *American Journal of Hypertension.* 1998;11(11):1364-70. doi:Doi 10.1016/S0895-7061(98)00154-X. *First study demonstrating the beneficial effect of bilateral nephrectomy in the treatment of resistant hypertension in hemodialysis patients.*
35. •• Fatouleh RH, Hammam E, Lundblad LC, Macey PM, McKenzie DK, Henderson LA et al. Functional and structural changes in the brain associated with the increase in muscle sympathetic nerve activity in obstructive sleep apnoea. *NeuroImage Clinical.* 2014;6:275-83. doi:10.1016/j.nicl.2014.08.021. *First study which simultaneously assessed functional brain activity and sympathetic nerve firing in obstructive sleep apnoea.*
36. Shantha GPS, Pancholy SB. Effect of renal sympathetic denervation on apnea-hypopnea index in patients with obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Breath.* 2015;19(1):29-34. doi:DOI 10.1007/s11325-014-0991-z.

37. •• Krum H, Schlaich MP, Sobotka PA, Bohm M, Mahfoud F, Rocha-Singh K et al. Percutaneous renal denervation in patients with treatment-resistant hypertension: final 3-year report of the Symplicity HTN-1 study. *Lancet*. 2014;383(9917):622-9. doi:10.1016/S0140-6736(13)62192-3. *First study demonstrating sustained blood pressure reduction out to 3 years following renal denervation in patients with resistant hypertension.*
38. •• Esler MD, Bohm M, Sievert H, Rump CL, Schmieder RE, Krum H et al. Catheter-based renal denervation for treatment of patients with treatment-resistant hypertension: 36 month results from the SYMPPLICITY HTN-2 randomized clinical trial. *Eur Heart J*. 2014;35(26):1752-9. doi:10.1093/eurheartj/ehu209. *First randomized-controlled study demonstrating the long-term effect of renal denervation on sustained blood pressure reduction in patients with resistant hypertension.*
39. •• Bhatt DL, Kandzari DE, O'Neill WW, D'Agostino R, Flack JM, Katzen BT et al. A controlled trial of renal denervation for resistant hypertension. *N Engl J Med*. 2014;370(15):1393-401. doi:10.1056/NEJMoa1402670. *First prospective, single-blind, randomized, sham-controlled study which did not show a significant reduction in blood pressure in patients with resistant hypertension when compared to sham control.*
40. •• Kandzari DE, Bhatt DL, Brar S, Devireddy CM, Esler M, Fahy M et al. Predictors of blood pressure response in the SYMPPLICITY HTN-3 trial. *Eur Heart J*. 2015;36(4):219-27. doi:10.1093/eurheartj/ehu441. *Post hoc analysis of the Symplicity HTN-3 trial which revealed several potential confounding factors that could explain the lack of a difference in blood pressure responses between the renal denervation and the sham control groups.*
41. •• Schlaich MP, Bart B, Hering D, Walton A, Marusic P, Mahfoud F et al. Feasibility of catheter-based renal nerve ablation and effects on sympathetic nerve activity and blood

- pressure in patients with end-stage renal disease. *International Journal of Cardiology*. 2013;168(3):2214-20. doi:DOI 10.1016/j.ijcard.2013.01.218. *First pilot study demonstrating the feasibility of renal denervation in patients with end-stage renal disease with a substantial reduction in blood pressure and sympathetic activity.*
42. •• Schlaich MP, Sobotka PA, Krum H, Lambert E, Esler MD. Renal Sympathetic-Nerve Ablation for Uncontrolled Hypertension. *New Engl J Med*. 2009;361(9):932-4. doi:Doi 10.1056/Nejmc0904179. *First case report of a patient with uncontrolled hypertension demonstrating a reduction in central sympathetic outflow after renal denervation indicating the potential involvement of afferent nerve fibers in sustained blood pressure reduction.*
 43. •• Lambert GW, Hering D, Esler MD, Marusic P, Lambert EA, Tanamas SK et al. Health-Related Quality of Life After Renal Denervation in Patients With Treatment-Resistant Hypertension. *Hypertension*. 2012;60(6):1479-84. doi:Doi 10.1161/Hypertensionaha.112.200865. *First study demonstrating an improvement in several aspects of quality of life in patients with resistant hypertension following renal denervation.*
 44. Lenski D, Kindermann I, Lenski M, Ukena C, Bunz M, Mahfoud F et al. Anxiety, depression, quality of life and stress in patients with resistant hypertension before and after catheter-based renal sympathetic denervation. *Eurointervention*. 2013;9(6):700-8. doi:Doi 10.4244/Eijv916a114.
 45. Geiger H, Naraghi R, Schobel HP, Frank H, Sterzel RB, Fahlbusch R. Decrease of blood pressure by ventrolateral medullary decompression in essential hypertension. *Lancet*. 1998;352(9126):446-9. doi:Doi 10.1016/S0140-6736(97)11343-5.
 46. Yin J, Wang WH, Zhang QB. A single microvascular decompression surgery cures a patient with trigeminal neuralgia, hemifacial spasm, tinnitus, hypertension, and

- paroxysmal supraventricular tachycardia caused by the compression of a vertebral artery. *Neurol India*. 2013;61(1):73-5. doi:Doi 10.4103/0028-3886.108016.
47. Legrady P, Voros E, Bajcsi D, Sonkodi S, Barzo P, Abraham G. Neurovascular Pulsatile Compression and Neurosurgical Decompression of the Rostral Ventrolateral Medulla in Medically Resistant Hypertensive Patients. *Kidney Blood Press R*. 2008;31(6):433-7. doi:Doi 10.1159/000195696.
 48. Legrady P, Voros E, Bajcsi D, Fejes I, Barzo P, Abraham G. Observations of Changes of Blood Pressure Before and after Neurosurgical Decompression in Hypertensive Patients with Different Types of Neurovascular Compression of Brain Stem. *Kidney Blood Press R*. 2013;37(4-5):451-7. doi:Doi 10.1159/000355725.
 49. •• Smith PA, Meaney JFM, Graham LN, Stoker JB, Mackintosh AF, Mary DASG et al. Relationship of neurovascular compression to central sympathetic discharge and essential hypertension. *Journal of the American College of Cardiology*. 2004;43(8):1453-8. doi:DOI 10.1016/j.jacc.2003.11.047. *First study indicating that increased sympathetic nerve firing is related to neurovascular compression.*
 50. Sendeski MM, Consolim-Colombo FM, Leite CC, Rubira MC, Lessa P, Krieger EM. Increased sympathetic nerve activity correlates with neurovascular compression at the rostral ventrolateral medulla. *Hypertension*. 2006;47(5):988-95. doi:Doi 10.1161/01.Hyp.0000214403.07762.47.
 51. Frank H, Heusser K, Geiger H, Fahlbusch R, Naraghi R, Schobel HP. Temporary Reduction of Blood Pressure and Sympathetic Nerve Activity in Hypertensive Patients After Microvascular Decompression. *Stroke*. 2009;40(1):47-51. doi:Doi 10.1161/Strokeaha.108.518670.
 52. Morimoto S, Aota Y, Sakuma T, Ichibangase A, Ikeda K, Sawada S et al. Efficacy of clonidine in a patient with refractory hypertension and chronic renal failure exhibiting

- neurovascular compression of the rostral ventrolateral medulla. *Hypertension Research*. 2009;32(3):227-8. doi:Doi 10.1038/Hr.2009.1.
53. •• Pereira EAC, Wang SY, Paterson DJ, Stein JF, Aziz TZ, Green AL. Sustained reduction of hypertension by deep brain stimulation. *Journal of Clinical Neuroscience*. 2010;17(1):124-7. doi:DOI 10.1016/j.jocn.2009.02.041. *First case report demonstrating blood pressure reduction and pain relief following deep brain stimulation in a patient with resistant hypertension.*
 54. Patel NK, Javed S, Khan S, Papouchado M, Malizia AL, Pickering AE et al. Deep Brain Stimulation Relieves Refractory Hypertension. *Neurology*. 2011;76(4):405-7. doi:Doi 10.1212/Wnl.0b013e3182088108.
 55. Sverrisdottir Y, Green A, Aziz T, Bahuri NF, Hyam J, Basnayake S et al. Effect of Deep Brain Stimulation on Sympathetic outflow in humans. *Faseb J*. 2014;28(1).
 56. Lohmeier TE, Irwin ED, Rossing MA, Serdar DJ, Kieval RS. Prolonged activation of the baroreflex produces sustained hypotension. *Hypertension*. 2004;43(2):306-11. doi:Doi 10.1161/01.Hyp.00001111837.73693.9b.
 57. •• Scheffers IJ, Kroon AA, Schmidli J, Jordan J, Tordoir JJ, Mohaupt MG et al. Novel baroreflex activation therapy in resistant hypertension: results of a European multi-center feasibility study. *J Am Coll Cardiol*. 2010;56(15):1254-8. doi:10.1016/j.jacc.2010.03.089. *First study demonstrating sustained blood pressure reduction following carotid sinus baroreflex stimulation in patients with resistant hypertension.*
 58. Bakris GL, Nadim MK, Haller H, Lovett EG, Schafer JE, Bisognano JD. Baroreflex activation therapy provides durable benefit in patients with resistant hypertension:

- results of long-term follow-up in the Rheos Pivotal Trial. *J Am Soc Hypertens.* 2012;6(2):152-8. doi:10.1016/j.jash.2012.01.003.
59. Hoppe UC, Brandt MC, Wachter R, Beige J, Rump LC, Kroon AA et al. Minimally invasive system for baroreflex activation therapy chronically lowers blood pressure with pacemaker-like safety profile: results from the Barostim neo trial. *Journal of the American Society of Hypertension.* 2012;6(4):270-6. doi:DOI 10.1016/j.jash.2012.04.004.
 60. Heusser K, Tank J, Engeli S, Diedrich A, Menne J, Eckert S et al. Carotid Baroreceptor Stimulation, Sympathetic Activity, Baroreflex Function, and Blood Pressure in Hypertensive Patients. *Hypertension.* 2010;55(3):619-26. doi:Doi 10.1161/Hypertensionaha.109.140665.
 61. Kishi T, Hirooka Y. Oxidative stress in the brain causes hypertension via sympathoexcitation. *Frontiers in Physiology.* 2012;3. doi:Unsp 335 Doi 10.3389/Fphys.2012.00335.
 62. Chrysohoou C, Panagiotakos DB, Pitsavos C, Skoumas J, Economou M, Papadimitriou L et al. The association between pre-hypertension status and oxidative stress markers related to atherosclerotic disease: The ATTICA study. *Atherosclerosis.* 2007;192(1):169-76. doi:DOI 10.1016/j.atherosclerosis.2006.04.030.
 63. Davis JT, Pasha DN, Khandrika S, Fung MM, Milic M, O'Connor DT. Central Hemodynamics in Prehypertension: Effect of the ss-Adrenergic Antagonist Nebivolol. *J Clin Hypertens.* 2013;15(1):69-74. doi:Doi 10.1111/Jch.12031.
 64. Hirooka Y, Kishi T, Ito K, Sunagawa K. Potential Clinical Application of Recently Discovered Brain Mechanisms Involved in Hypertension. *Hypertension.* 2013;62(6):995-1002. doi:Doi 10.1161/Hypertensionaha.113.00801.