Leaf manganese accumulation and phosphorus-acquisition efficiency

Hans Lambers¹, Patrick E. Hayes¹, Etienne Laliberté¹, Rafael S. Oliveira^{2,1} and Benjamin L. Turner^{3,1}

¹School of Plant Biology, The University of Western Australia, Stirling Highway, Crawley (Perth), WA 6009, Australia

²Departamento de Biologia Vegetal, Universidade Estadual de Campinas. Rua Monteiro Lobato 255, Campinas 13083-862, Brazil

³Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Republic of Panama

Corresponding author: Lambers, H. (hans.lambers@uwa.edu.au)

Keywords Carboxylates, exudation, manganese, phosphorus, phosphorus-acquisition efficiency

Plants that deploy a phosphorus- (P) mobilising strategy based on the release of carboxylates tend to have high leaf manganese concentrations ([Mn]). This occurs because the carboxylates not only mobilise soil inorganic and organic P, but also a range of micronutrients, including Mn. Concentrations of most other micronutrients increase to a small extent, but Mn accumulates to significant levels, even when plants grow in soil with low concentrations of exchangeable Mn availability. Here, we propose that leaf [Mn] can be used to select for genotypes that are more efficient at acquiring P when soil P availability is low. Likewise, leaf [Mn] can be used to screen for belowground functional traits related to nutrient-acquisition strategies among species in low-P habitats.

Phosphorus-acquisition strategies

Here we explore the idea of using leaf manganese concentrations ([Mn]) to indicate a carboxylate-releasing phosphorus- (P) acquisition strategy. The rationale behind this contention is that both the availability of P and that of Mn are increased when roots release carboxylates into the rhizosphere [1] (Figure 1). The availability of some other micronutrients is also enhanced, but most of these do not lead to such a strong signal as that provided by Mn. The release of carboxylates into the rhizosphere is important for P-acquisition, because they mobilise not only inorganic P, but also organic P, which can be a major fraction of soil P, especially when P availability is low [2].

Addressing this topic is timely, because there is a growing interest among plant ecologists in belowground functional traits, to complement the suite of 'easy-to-measure' aboveground traits [3]. Furthermore, because of the gradual decline in phosphate rock that is used to produce P fertilisers [4], there is an increasing need for more P-efficient cropping systems [5]. Therefore, a simple tool to screen for P-acquisition efficiency in crop species will be welcomed by agronomists and plant breeders.

Manganese as a plant nutrient

The significance of Mn as an essential plant nutrient was firmly established in 1922 [6]. More recent work has revealed the role of Mn in redox processes, as an activator of a large range of enzymes, and as a cofactor of a small number of enzymes, including proteins required for light-induced water oxidation in photosystem II [7, 8]. Crop plants that contain 50 $\mu\text{g Mn g}^{-1}$ dry weight (DW) in their leaves are considered to have sufficient Mn for maximum growth and yield [9]. Conversely, Mn toxicity can occur when plants are grown at moderately low soil pH or in flooded soils, when Mn availability is increased [10-13] and its uptake is not tightly regulated. Critical toxicity concentrations in leaves range from 200 to 3,500 $\mu\text{g Mn g}^{-1}$ DW [14], but some hyperaccumulators, such as Proteaceae species in New Caledonia, may contain >10,000 $\mu\text{g Mn g}^{-1}$ DW without harmful effects

[14, 15]. Mechanisms that counter Mn toxicity in plants involve Mn export from the cytoplasm, across the tonoplast for sequestration into the vacuole, or across the plasma membrane out of the cell [16]. The *Arabidopsis AtMTP* family of genes encode proteins of the cation diffusion facilitator family, some of which play a role in metal tolerance [17]. Expression of cation diffusion facilitators in a Mn-hypersensitive yeast mutant restores Mn tolerance to wild-type levels, showing the importance of this transport system for Mn tolerance [18].

High leaf manganese concentrations in non-mycorrhizal species with cluster roots

Relatively high leaf [Mn] are typically found in species that produce cluster roots, particularly Proteaceae species, which are almost all non-mycorrhizal and occur on soils with very low P availability [19-22]. Cluster roots also occur in actinorhizal species and in many Fabaceae [1, 23]. These specialised roots release large amounts of carboxylates in an 'exudative burst' to mobilise P, and this also mobilises Mn [23, 24]. For a large number of Proteaceae, the range of [Mn] is 126-10,000 $\mu\text{g Mn g}^{-1}\text{ DW}$ [20, 25-29]. In New Caledonia, no Proteaceae species exhibit leaf [Mn] <100 $\mu\text{g Mn g}^{-1}\text{ DW}$ [20]. Likewise, in Fabaceae species with cluster roots, relatively high leaf [Mn] have been observed: 7,370 $\mu\text{g Mn g}^{-1}\text{ DW}$ in *Lupinus albus*, which is also non-mycorrhizal [30] and 120 $\mu\text{g Mn g}^{-1}\text{ DW}$ in *Aspalathus linearis* [31]. These high concentrations can be explained by the ability of cluster roots to mobilise Mn as well as P [24, 32, 33]. For example, in a glasshouse experiment with *Hakea prostrata* (Proteaceae), variation in leaf [Mn] was positively correlated with investment in cluster roots [34], and similar results were found for *L. albus* [35]. Given that a concentration of 50 $\mu\text{g Mn g}^{-1}\text{ DW}$ is considered sufficient for maximum growth of crop plants [9], concentrations >100 $\mu\text{g Mn g}^{-1}\text{ DW}$ are considered 'high', especially for species with scleromorphic (see Glossary) leaves (e.g., many Proteaceae). However, the exact concentrations will also depend on Mn availability in the soil (which is strongly pH-dependent), and hence the concentration in leaves should be compared with that of other species growing at the same location [22].

Manganese is taken up as Mn^{2+} by roots from the rhizosphere, partly involving broad-specificity transporters [16, 36] (Box 1). The broad specificity of these transporters accounts for Mn accumulation and toxicity in plants where soil Mn availability is high. The availability of soil Mn increases with decreasing soil pH, until approximately pH 5, at which point the availability declines again [13, 37]. Importantly, Mn availability is also increased by root exudation of carboxylates, which chelate Mn and reduce Mn^{4+} to Mn^{2+} in either acidic or alkaline soils [38].

The literature reviewed for non-mycorrhizal, cluster-rooted species discussed above leads to the hypothesis that leaf [Mn] can be used as a proxy for the carboxylate-releasing P-mobilising strategy and as a screening tool for P-acquisition efficiency when soil P availability is low. Here, we explore this hypothesis.

High leaf manganese concentrations in other non-mycorrhizal species

A high leaf [Mn] has been found in *Phytolacca acinosa* and *P. americana*, with up to 19,300 $\mu g Mn g^{-1}$ DW [39-43]; these Mn-hyperaccumulating species belong to Phytolaccaceae, a non-mycorrhizal and non-cluster-rooting family [44, 45]. Manganese-hyperaccumulating species contain about 100 times more Mn than non-accumulator species, reaching at least 1,000 $\mu g Mn g^{-1}$ DW [46]. There is no information in the literature on the rhizosphere chemistry of *Phytolacca* species, and so it is unknown if Mn accumulation depends on the release of carboxylates, protons or both. Based on the high oxalate concentration in leaves of *P. americana* [47], we surmise that this species releases protons generated in the production of oxalic acid, and that the high internal concentration of carboxylate anions (oxalate) is used to internally chelate and detoxify Mn.

Polygonum perfoliatum and *P. hydropiper* (Polygonaceae) are also Mn-hyperaccumulating herbaceous species, with shoot concentrations up to 18,340 $\mu g Mn g^{-1}$ DW [42]. Polygonaceae species lack cluster roots and are considered non-mycorrhizal [44]; however, *P. viviparum* has been

found to be ectomycorrhizal [48]. No mechanism(s) accounting for their Mn hyperaccumulation are known.

Like the cluster-root forming *Lupinus albus*, the non-cluster-root-producing non-mycorrhizal *L. angustifolius* also accumulates high concentrations of Mn in its leaves: 1,108 $\mu\text{g Mn g}^{-1}$ DW [49]. Such Mn accumulation is thought to be due to reducing conditions in the rhizosphere, allowing for the increased availability of Mn [49]. Both *L. albus* and *L. angustifolius* are non-mycorrhizal [1] and release relatively large amounts of carboxylates and protons into their rhizosphere [50]. High leaf [Mn] is therefore not restricted to species producing cluster roots.

Across a coastal dune chronosequence in Jurien Bay in Western Australia, leaf [Mn] is consistently greater in non-mycorrhizal species compared with co-occurring mycorrhizal species, with most non-mycorrhizal species known to release carboxylates [22]. Interestingly, this occurs across all soils along this 2-million year dune chronosequence, despite the soils showing a wide range of pH values (~5–9). At the community level and within individual non-mycorrhizal species, leaf [Mn] also increases with increasing soil age and associated declines in soil P availability (Figure 2A) and pH, but is not influenced by total or exchangeable soil [Mn] [22]. One species with high leaf [Mn], *Conostylis candicans* (100 $\mu\text{g Mn g}^{-1}$ DW), has been subsequently shown to release a range of carboxylates in its rhizosphere (F. Albornoz and E. Laliberté, unpubl.) in soils spanning a wide range of pH (6–8; determined in 10 mM CaCl_2) and total [P] of 20–430 mg P kg^{-1} . Other non-cluster rooted non-mycorrhizal species with high leaf [Mn] along the same chronosequence remain to be further investigated.

High shoot [Mn] (451 to 1156 $\mu\text{g Mn g}^{-1}$ DW) have been observed in *Discocactus placentiformis*, a non-mycorrhizal cactus species, abundant in low-P soils in the *campos rupestres* of Central Brazil (Figure 2B) [51]. When grown in a low-P nutrient solution, the roots of this cactus increase their exudation of carboxylates, predominantly oxalate, but also malate and citrate. Other non-mycorrhizal species in the *campos rupestres* also show high leaf [Mn], when compared with

mycorrhizal species in the same communities (Figure 2B). We speculate that the high leaf [Mn] of non-mycorrhizal *campos rupestres* species is accounted for by root carboxylate release.

High leaf manganese concentrations in species of typically mycorrhizal families

Thirty one *Alyxia* species (Apocynaceae) from New Caledonia all contain high leaf [Mn] [15], with no information on their Mn-uptake mechanisms. *Alyxia* species are considered arbuscular mycorrhizal [52]. *Cupania tenuivalvis* (Sapindaceae) is a species native on acidic soils in cerradão vegetation in Brazil, with leaf [Mn] of 3,300 $\mu\text{g Mn g}^{-1}$ DW [53]. We can only speculate that the mechanism accounting for Mn hyperaccumulation in *Alyxia* and *Cupania* species are similar to those we discuss for other species in this section. If so, carboxylates would be released by the plant, as there is no conclusive evidence that arbuscular mycorrhizal fungi access significant amounts of insoluble P sources in soil [54].

Within the genus *Eucalyptus* (Myrtaceae), the two subgenera, *Symphyomyrtus* and *Monocalyptus* differ nutritionally, based on [Mn] in their leaves, stem and bark; [Mn] are consistently greater in symphyomyrts than in monocalypts [55]. These two subgenera are likely either arbuscular mycorrhizal or both arbuscular mycorrhizal and ectomycorrhizal [52]. There is no evidence that these differences in foliar [Mn] are accounted for by differences in soil [Mn], and hence it is hypothesised that they are most likely due to differences in Mn mobilisation in the rhizosphere. Leaf [Mn] in eucalypts are so high (800 $\mu\text{g Mn g}^{-1}$ DW in green leaf-fall and up to 2,800 $\mu\text{g Mn g}^{-1}$ DW in leaves of glasshouse-grown seedlings) that they are within the toxic range for most plants [14]. The mechanism for Mn accumulation is unknown for these and other eucalypts [56], but we do know that *Eucalyptus gummifera* can access poorly soluble forms of inorganic P (aluminium phosphate and iron phosphate) [57], presumably through the releases of carboxylates.

Austromyrtus bidwillii (Myrtaceae), a tropical rainforest tree in north-eastern Australia, hyperaccumulates Mn in its leaves (up to 19,200 $\mu\text{g Mn g}^{-1}$ DW) and bark (up to 26,500 $\mu\text{g Mn g}^{-1}$

DW) [58]. Since its leaves contain about three times more carboxylates (up to 123,000 $\mu\text{g g}^{-1}$ DW) than required to chelate all Mn, it is likely that, as in the *Phytolacca* species discussed above, it releases predominantly protons, and not carboxylates, thereby acidifying the rhizosphere and consequently mobilising soil Mn [58].

Gossia bidwillii (Myrtaceae) is a Mn-hyperaccumulating eastern Australian subtropical rainforest tree [59, 60]. There is no published information on the mechanism accounting for the Mn accumulation of this species. In another Australian Mn-hyperaccumulating species, *Denhamia fournieri* (previously known as *Maytenus founieri*) (Celastraceae), Mn appears to be associated with carboxylates in leaves [61], suggesting Mn accumulation is associated with proton release as in *Phytolacca* species, as discussed above.

Chengiopanax sciadophylloides (synonyms: *Eleutherococcus sciadophylloides* and *Acanthopanax sciadophylloides*) (Araliaceae) is a Mn-hyperaccumulating Japanese tree; Mn accumulation is based on acidification of the rhizosphere, but not carboxylate release, as in the other species discussed above [62, 63]. Hyperaccumulation in this species occurs in non-contaminated forest soils and is specific to Mn, not other metals [64-66]. A ZIP gene analog (ZIP: ZRT, IRT-like protein; Box 1) encodes a protein with 65% or less sequence identity with ZIPs of other herbaceous species. Expression of this gene is induced in the callus of Mn-deficient *C. sciadophylloides*, but it does not show Zn or Fe transport [67].

Schima superba (Theaceae) is a Mn-accumulating subtropical tree species native to China [68], without known mechanisms accounting for its Mn accumulation.

In summary, Mn hyperaccumulation in mycorrhizal species does not reflect soil [Mn] or [P] or soil pH, but is associated with Mn mobilisation in the rhizosphere, most likely due to the release of protons and subsequent acidification of the rhizosphere. The carboxylates generated to produce

the protons released into the rhizosphere are used internally to bind Mn inside plant tissues, thus reducing the toxic effects of Mn hyperaccumulation.

Variation in leaf [Mn] or root [Mn] as dependent on P treatments

In barley (*Hordeum vulgare*), an elevated P supply reduces Mn acquisition [69], suggesting a role for carboxylates in mobilising both P and Mn in this species. For example, at a high P supply, carboxylate release would be suppressed, thus reducing Mn uptake. It has been shown that mycorrhizal plants of subclover (*Trifolium subterraneum*) grown at a limiting P supply had higher leaf [P] , but a relatively lower leaf [Mn] than their non-mycorrhizal counterparts [70], indicating alternative strategies to acquire P, depending on the presence of mycorrhizal inoculum [71]. Following a pulse of P, the root [Mn] declined, suggesting a role for P- and Mn-mobilising exudates, as indicated by the reduced uptake of Mn at a high P supply.

In summary, plants that exhibit a P-mobilising strategy dependent on the release of carboxylates or protons, show lower leaf [Mn] when supplied with sufficient P in comparison with P-limited plants.

Effects of carboxylate-releasing plants on leaf [Mn] in neighbouring plants

When wheat (*Triticum aestivum*) is grown in a cropping situation together with the non-mycorrhizal cluster-rooted *Lupinus albus*, its leaves contain higher leaf [Mn] [30]. This shows that wheat can enhance its Mn uptake when neighbouring white lupins mobilise Mn in soil. Similarly, when grown in pots together with *Banksia attenuata* (Proteaceae), leaf [Mn] and growth of the ectomycorrhizal species *Scholzia involucreata* (Myrtaceae) are enhanced, indicating facilitation of Mn uptake by a cluster-rooted species [72].

Concluding remarks

Plants that release relatively large amounts of carboxylates tend to have relatively high leaf [Mn], *e.g.*, Proteaceae and some lupin species [20]. However, some species that hyperaccumulate Mn may not release carboxylates, but strongly acidify their rhizosphere instead, such as *Chengiopanax sciadophylloides* and *Phytolacca* species [62, 63]. Therefore, we suggest that high leaf [Mn] in an environment with a low P availability should be taken only as a strong indication of root carboxylate exudation, primarily for non-mycorrhizal species. High leaf [Mn] by itself does not provide firm evidence for carboxylate exudation. Results showing high leaf [Mn] must be followed up with analyses of rhizosphere carboxylates, or possibly other exudates, before inferring that the studied species must have a specialised P-mobilising strategy. High leaf [Mn] is possibly only associated with carboxylate release into the rhizosphere in non-mycorrhizal species of the ‘Proteaceae type’, *sensu* Lambers & Teste [73] and in mycorrhizal species that can switch to a carboxylate-releasing strategy [71]. These are typically associated with soils of low P availability, *e.g.*, along the Jurien Bay chronosequence [22] and on the sandplains of the Brazilian cerrado [51]. On the other hand, high leaf [Mn] in species of the ‘Brassicaceae type’, which occur on nutrient-rich soils, might be associated with release of protons and internal carboxylate accumulation. Acidification of the rhizosphere at low soil pH will render P less available, rather than more [74]. This holds for both inorganic and organic P [75].

We propose leaf [Mn] as a valuable tool to screen for P-efficient crop genotypes in a common environment with low soil P availability, provided the promising genotypes are subsequently further investigated, focusing on rhizosphere exudates, and primarily carboxylates (Box 2). Within a plant community on soils with low P availability, leaf [Mn] also provides a strong indication of specific species utilising P-mobilising carboxylate-releasing strategies [22] (Box 2). When plants that lack such a strategy are analysed, as dependent on neighbouring species, some of which do and others that do not depend on the P-mobilising carboxylate-releasing strategy, some evidence might be obtained for facilitation of nutrient acquisition, as shown in a mixed cropping

situation [30] and a pot experiment [72]. The approach is not advocated to be used between sites, as these may differ in soil pH and Mn availability [53].

We also propose leaf [Mn] as a useful trait for within-site comparisons to add to the standard set of traits typically considered in trait-based community ecology studies and comparative functional ecology [3]. In these two fields of study, a small number of traits are typically measured across a wide range of co-occurring species. Consequently, easily-measured aboveground traits (*e.g.*, specific leaf area, leaf dry matter content) are favoured, with the underlying (but generally untested) assumption that above- and belowground traits are coordinated among species [76]. Therefore, most trait-based community ecology and comparative functional ecology studies are currently biased towards aboveground traits. Leaf [Mn] could provide an easily-measured aboveground trait that reflects belowground functioning as a time-integrated proxy for P acquisition via carboxylate release. This is especially important because carboxylate release can occur in pulses over a short period of time [77, 78].

In conclusion, we propose that leaf [Mn] analysis is a valuable screening tool, both in breeding crops for a high P-acquisition efficiency and in identifying species in a community that use a P-mobilising strategy.

Acknowledgements

HL was supported by the Australian Research Council (ARC; DP0985685 and DP110101120); PEH was supported by an Australian Postgraduate Award through the University of Western Australia, EL was supported by a DECRA (DE120100352) from the ARC; RSO was supported by São Paulo Research Foundation (Fapesp 2010/172040 and Fapesp 2011/520720).

References

1 Lambers, H., *et al.* (2013) How a phosphorus-acquisition strategy based on carboxylate exudation powers the success and agronomic potential of lupines (*Lupinus*, Fabaceae). *Am. J. Bot.* 100, 263-288

2 Turner, B.L., *et al.* (2013) Soil microbial biomass and the fate of phosphorus during long-term ecosystem development. *Plant Soil* 367, 225-234

3 Pérez-Harguindeguy, N., *et al.* (2013) New handbook for standardised measurement of plant functional traits worldwide. *Aust. J. Bot.* 61, 167-234

4 Scholz, R.W. and Wellmer, F.-W. (2013) Approaching a dynamic view on the availability of mineral resources: what we may learn from the case of phosphorus? *Global Environ. Change* 23, 11-27

5 Simpson, R.J., *et al.* (2011) Strategies and agronomic interventions to improve the phosphorus-use efficiency of farming systems. *Plant Soil* 349, 89-120

6 McHargue, J.S. (1922) The role of manganese in plants. *J. Am. Chem. Soc.* 44, 1592-1598

7 Broadley, M., *et al.* (2012) Function of nutrients: micronutrients. In *Marschner's Mineral Nutrition of Higher Plants, Third Edition* (Marschner, P., ed), pp. 191-248, Academic Press

8 Stout, P.R. and Arnon, D.I. (1939) Experimental methods for the study of the role of copper, manganese, and zinc in the nutrition of higher plants. *Am. J. Bot.* 26, 144-149

9 Epstein, E. and Bloom, A.J. (2005) *Mineral Nutrition of Plants: Principles and Perspectives*. Sinauer

10 El-Jaoual, T. and Cox, D.A. (1998) Manganese toxicity in plants. *J. Plant Nutr.* 21, 353-386

11 Horiguchi, T. (1988) Mechanism of manganese toxicity and tolerance of plants. *Soil Sci. Plant Nutr.* 34, 65-73

12 Foy, C.D., *et al.* (1978) The physiology of metal toxicity in plants. *Annu. Rev. Plant Physiol.* 29, 511-566

13 Lambers, H., *et al.* (2008) *Plant Physiological Ecology, second edition*. Springer

14 Krämer, U. (2010) Metal hyperaccumulation in plants. *Annu. Rev. Plant Biol.* 61, 517-534

15 Brooks, R.R., *et al.* (1981) Studies on manganese-accumulating *Alyxia* species from New Caledonia *Taxon* 30, 420-423

16 Pittman, J.K. (2005) Managing the manganese: molecular mechanisms of manganese transport and homeostasis. *New Phytol.* 167, 733-742

17 Delhaize, E., *et al.* (2007) A role for the *AtMTP11* gene of *Arabidopsis* in manganese transport and tolerance. *Plant J.* 51, 198-210

18 Peiter, E., *et al.* (2007) A secretory pathway-localized cation diffusion facilitator confers plant manganese tolerance. *Proceedings of the National Academy of Sciences* 104, 8532-8537

19 Lambers, H., *et al.* (2006) Root structure and functioning for efficient acquisition of phosphorus: matching morphological and physiological traits. *Ann. Bot.* 98, 693-713

20 Jaffré, T. (1979) Accumulation du manganèse par les Protéacées de Nouvelle Calédonie. *C.R. Acad. Sci. Paris D* 289, 425-428

21 Lambers, H., *et al.* (2008) Plant nutrient-acquisition strategies change with soil age. *Trends Ecol. Evol.* 23, 95-103

22 Hayes, P., *et al.* (2014) Foliar nutrient concentrations and resorption efficiency in plants of contrasting nutrient-acquisition strategies along a 2-million-year dune chronosequence. *J. Ecol.* 102, 396-410

23 Shane, M.W. and Lambers, H. (2005) Cluster roots: a curiosity in context. *Plant Soil* 274, 101-125

24 Gardner, W.K., *et al.* (1982) The acquisition of phosphorus by *Lupinus albus* L. I. Some characteristics of the soil/root interface. *Plant Soil* 68, 19-32

25 Fernando, D.R., *et al.* (2009) Foliar Mn accumulation in eastern Australian herbarium specimens: prospecting for 'new' Mn hyperaccumulators and potential applications in taxonomy. *Ann. Bot.* 103, 931-939

26 Denton, M.D., *et al.* (2007) *Banksia* species (Proteaceae) from severely phosphorus-impooverished soils exhibit extreme efficiency in the use and re-mobilization of phosphorus. *Plant Cell Environ.* 30, 1557-1565

27 Fernando, D.R., *et al.* (2006) Manganese accumulation in the leaf mesophyll of four tree species: a PIXE/EDAX localization study. *New Phytol.* 171, 751-758

28 Rabier, J., *et al.* (2007) Characterization of metal tolerance and accumulation in *Grevillea exul* var *exul*. *Int. J. Phytorem.* 9, 419-435

287 29 Child, R. and Smith, A.N. (1960) Manganese toxicity in *Grevillea robusta*. *Nature* 186, 1067
288 30 Gardner, W.K. and Boundy, K.A. (1983) The acquisition of phosphorus by *Lupinus albus* L. IV. The
289 effect of interplanting wheat and white lupin on the growth and mineral composition of the two
290 species. *Plant Soil* 70, 391-402
291 31 Morton, J.F. (1983) Rooibos tea, *Aspalathus linearis*, a caffeineless, low-tannin beverage. *Econ.*
292 *Bot.* 37, 164-173
293 32 Grierson, P.F. and Attiwill, P.M. (1989) Chemical characteristics of the proteoid root mat of
294 *Banksia integrifolia* L. *Aust. J. Bot.* 37, 137-143
295 33 Dinkelaker, B., et al. (1995) Distribution and function of proteoid roots and other root clusters.
296 *Bot. Acta* 108, 193-200
297 34 Shane, M.W. and Lambers, H. (2005) Manganese accumulation in leaves of *Hakea prostrata*
298 (Proteaceae) and the significance of cluster roots for micronutrient uptake as dependent on
299 phosphorus supply. *Physiol. Plant.* 124, 441-450
300 35 Moraghan, J. (1992) Iron-manganese relationships in white lupine grown on a calciaquoll. *Soil Sci.*
301 *Soc. Am. J.* 56, 471-475
302 36 Korshunova, Y.O., et al. (1999) The IRT1 protein from *Arabidopsis thaliana* is a metal transporter
303 with a broad substrate range. *Plant Mol. Biol.* 40, 37-44
304 37 Godo, G.H. and Reisenauer, H.M. (1980) Plant effects on soil manganese availability. *Soil Sci. Soc.*
305 *Am. J.* 44, 993-995
306 38 Jauregui, M.A. and Reisenauer, H.M. (1982) Dissolution of oxides of manganese and iron by root
307 exudate components. *Soil Sci. Soc. Am. J.* 46, 314-317
308 39 Xue, S.G., et al. (2004) Manganese uptake and accumulation by the hyperaccumulator plant
309 *Phytolacca acinosa* Roxb. (Phytolaccaceae). *Environ. Pollut.* 131, 393-399
310 40 Xue, S.G., et al. (2005) Manganese uptake and accumulation by two populations of *Phytolacca*
311 *acinosa* Roxb. (Phytolaccaceae). *Water, Air, and Soil Pollution* 160, 3-14
312 41 Xu, X., et al. (2006) Distribution and mobility of manganese in the hyperaccumulator plant
313 *Phytolacca acinosa* Roxb. (Phytolaccaceae). *Plant Soil* 285, 323-331
314 42 Liu, P., et al. (2010) Manganese tolerance and accumulation in six Mn hyperaccumulators or
315 accumulators. *Plant Soil* 335, 385-395
316 43 Min, Y., et al. (2007) Accumulation and uptake of manganese in a hyperaccumulator *Phytolacca*
317 *americana*. *Miner. Eng.* 20, 188-190
318 44 Gerdemann, J.W. (1968) Vesicular-arbuscular mycorrhiza and plant growth. *Annu. Rev.*
319 *Phytopathol.* 6, 397-418
320 45 Janos, D.P. (1980) Vesicular-arbuscular mycorrhizae affect lowland tropical rain forest plant
321 growth. *Ecology* 61, 151-162
322 46 Brooks, R.R., et al. (1998) Phytomining. *Trends Plant Sci.* 3, 359-362
323 47 Dou, C.M., et al. (2009) Accumulation and detoxification of manganese in hyperaccumulator
324 *Phytolacca americana*. *Plant Biol.* 11, 664-670
325 48 Massicotte, H.B., et al. (1998) Anatomical aspects of field ectomycorrhizas on *Polygonum*
326 *viviparum* (Polygonaceae) and *Kobresia bellardii* (Cyperaceae). *Mycorrhiza* 7, 287-292
327 49 Reay, P.F. and Waugh, C. (1981) Mineral-element composition of *Lupinus albus* and *Lupinus*
328 *angustifolius* in relation to manganese accumulation. *Plant Soil* 60, 435 - 444
329 50 Pearse, S.J., et al. (2006) Carboxylate release of wheat, canola and 11 grain legume species as
330 affected by phosphorus status. *Plant Soil* 288, 127-139
331 51 Abrahão, A., et al. (2014) Convergence of a specialized root trait in plants from nutrient-
332 impoverished soils: phosphorus-acquisition strategy in a nonmycorrhizal cactus. *Oecologia* 176, 345-
333 355
334 52 Wang, B. and Qiu, Y.-L. (2006) Phylogenetic distribution and evolution of mycorrhizas in land
335 plants. *Mycorrhiza* 16, 299-363

336 53 Viani, R.A.G., *et al.* (2014) Soil pH accounts for differences in species distribution and leaf nutrient
 337 concentrations of Brazilian woodland savannah and Seasonally Dry forest species. *Perspect. Plant*
 338 *Ecol. Evol. Syst.*
 339 54 Smith, S.E. and Read, D.J. (2008) *Mycorrhizal Symbiosis*. Academic Press and Elsevier
 340 55 Hill, J., *et al.* (2001) Does manganese play a role in the distribution of the eucalypts? *Aust. J. Bot.*
 341 49, 1-8
 342 56 Grigg, A.M., *et al.* (2008) Water relations and mineral nutrition of closely related woody plant
 343 species on desert dunes and interdunes. *Aust. J. Bot.* 56, 27-43
 344 57 Mullette, K.J., *et al.* (1974) Insoluble phosphorus usage by *Eucalyptus*. *Plant Soil* 41, 199-205
 345 58 Bidwell, S.D., *et al.* (2002) Hyperaccumulation of manganese in the rainforest tree *Austromyrtus*
 346 *bidwillii* (Myrtaceae) from Queensland, Australia. *Funct. Plant Biol.* 29, 899-905
 347 59 Fernando, D.R., *et al.* (2006) In vivo localization of manganese in the hyperaccumulator *Gossia*
 348 *bidwillii* (Benth.) N. Snow & Guymier (Myrtaceae) by cryo-SEM/EDAX. *Plant Cell Environ.* 29, 1012-
 349 1020
 350 60 Fernando, D.R., *et al.* (2007) Variability of Mn hyperaccumulation in the Australian rainforest tree
 351 *Gossia bidwillii* (Myrtaceae). *Plant Soil* 293, 145-152
 352 61 Fernando, D.R., *et al.* (2008) Foliar manganese accumulation by *Maytenus founieri* (Celastraceae)
 353 in its native New Caledonian habitats: populational variation and localization by X-ray microanalysis.
 354 *New Phytol.* 177, 178-185
 355 62 Mizuno, T., *et al.* (2008) Age-dependent manganese hyperaccumulation in *Chengiopanax*
 356 *sciadophylloides* (Araliaceae). *J. Plant Nutr.* 31, 1811-1819
 357 63 Mizuno, T., *et al.* (2006) Continual pH lowering and manganese dioxide solubilization in the
 358 rhizosphere of the Mn-hyperaccumulator plant *Chengiopanax sciadophylloides*. *Soil Sci. Plant Nutr.*
 359 52, 726-733
 360 64 Memon, A.R. and Yatazawa, M. (1984) Nature of manganese complexes in manganese
 361 accumulator plant - *Acanthopanax sciadophylloides*. *J. Plant Nutr.* 7, 961-974
 362 65 Memon, A.R., *et al.* (1979) Absorption and accumulation of iron, manganese and copper in plants
 363 in the temperate forest of central Japan. *Soil Sci. Plant Nutr.* 25, 611-620
 364 66 Mizuno, T., *et al.* (2013) Manganese hyperaccumulation from non-contaminated soil in
 365 *Chengiopanax sciadophylloides* Franch. et Sav. and its correlation with calcium accumulation. *Soil Sci.*
 366 *Plant Nutr.* 59, 591-602
 367 67 Mizuno, T., *et al.* (2008) Cloning of ZIP family metal transporter genes from the manganese
 368 hyperaccumulator plant *Chengiopanax sciadophylloides*, and its metal transport and resistance
 369 abilities in yeast. *Soil Sci. Plant Nutr.* 54, 86-94
 370 68 Yang, S.X., *et al.* (2008) Manganese uptake and accumulation in a woody hyperaccumulator,
 371 *Schima superba*. *Plant, Soil and Environment* 54, 441-446
 372 69 Pedas, P., *et al.* (2011) Elevated phosphorus impedes manganese acquisition by barley plants.
 373 *Frontiers in Plant Science* 2, 37
 374 70 Nazeri, N., *et al.* (2013) Do arbuscular mycorrhizas or heterotrophic soil microbes contribute
 375 toward plant acquisition of a pulse of mineral phosphate? *Plant Soil* 373 699-710
 376 71 Ryan, M.H., *et al.* (2012) Carbon trading for phosphorus gain: the balance between rhizosphere
 377 carboxylates and mycorrhizal symbiosis in plant phosphorus acquisition. *Plant Cell Environ.* 35, 2061-
 378 2220
 379 72 Muler, A.L., *et al.* (2014) Does cluster-root activity of *Banksia attenuata* (Proteaceae) benefit
 380 phosphorus or micronutrient uptake and growth of neighbouring shrubs? *Oecologia* 174, 23-31
 381 73 Lambers, H. and Teste, F.P. (2013) Interactions between arbuscular mycorrhizal and non-
 382 mycorrhizal plants: do non-mycorrhizal species at both extremes of nutrient-availability play the
 383 same game? *Plant Cell Environ.* 36, 1911-2070
 384 74 Lindsay, W.L. (1979) *Chemical Equilibria in Soils*. John Wiley & Sons
 385 75 Celi, L. and Barberis, E. (2005) Abiotic stabilization of organic phosphorus in the environment. In
 386 *Organic Phosphorus in the Environment* (Turner, B.L., *et al.*, eds), pp. 113-132, CAB International

387 76 Reich, P.B. (2014) The world-wide 'fast-slow' plant economics spectrum: a traits manifesto. *J.*
388 *Ecol.* 102, 275-301

389 77 Watt, M. and Evans, J.R. (1999) Proteoid roots. Physiology and development. *Plant Physiol.* 121,
390 317-323

391 78 Shane, M.W., *et al.* (2004) Developmental physiology of cluster-root carboxylate synthesis and
392 exudation in harsh hakea. Expression of phosphoenolpyruvate carboxylase and the alternative
393 oxidase. *Plant Physiol.* 135, 549-560

394 79 Nagarajah, S., *et al.* (1970) Competitive adsorption of phosphate with polygalacturonate and
395 other organic anions on kaolinite and oxide surfaces. *Nature* 228, 83-85

396 80 Turner, B.L. and Laliberté, E. (2014) Soil development and nutrient availability along a two million
397 year coastal dune chronosequence under species-rich Mediterranean shrubland in southwestern
398 Australia. *Ecosystems* in press

399 81 McBain, J.W. (1909) The mechanism of the adsorption ("sorption") of hydrogen by carbon.
400 *Philosophical Magazine Series 6* 18, 916-935

401 82 Guerinot, M.L. (2000) The ZIP family of metal transporters. *Biochim. Biophys. Acta -*
402 *Biomembranes* 1465, 190-198

403 83 Hall, J.L. and Williams, L.E. (2003) Transition metal transporters in plants. *J. Exp. Bot.* 54, 2601-
404 2613

405 84 Kobayashi, T. and Nishizawa, N.K. (2012) Iron uptake, translocation, and regulation in higher
406 plants. *Annu. Rev. Plant Biol.* 63, 131-152

407 85 Connolly, E.L., *et al.* (2002) Expression of the IRT1 metal transporter is controlled by metals at the
408 levels of transcript and protein accumulation. *Plant Cell* 14, 1347-1357

409 86 Eckhardt, U., *et al.* (2001) Two iron-regulated cation transporters from tomato complement metal
410 uptake-deficient yeast mutants. *Plant Mol. Biol.* 45, 437-448

411 87 Bughio, N., *et al.* (2002) Cloning an iron-regulated metal transporter from rice. *J. Exp. Bot.* 53,
412 1677-1682

413 88 Römheld, V. and Schaaf, G. (2004) Iron transport in plants: future research in view of a plant
414 nutritionist and a molecular biologist. *Soil Sci. Plant Nutr.* 50, 1003-1012

415 89 Martínez-Cuenca, M.-R., *et al.* (2013) Effects of high levels of zinc and manganese ions on
416 Strategy I responses to iron deficiency in citrus. *Plant Soil* 373, 943-953

417 90 Venturas, M., *et al.* (2014) Root iron uptake efficiency of *Ulmus laevis* and *U. minor* and their
418 distribution in soils of the Iberian Peninsula. *Frontiers in Plant Science* 5

419 91 Conte, S.S. and Walker, E.L. (2011) Transporters contributing to iron trafficking in plants.
420 *Molecular Plant* 4, 464-476

421 92 Baxter, I.R., *et al.* (2008) The leaf ionome as a multivariable system to detect a plant's
422 physiological status. *Proc. Natl. Acad. Sci. USA* 105, 12081-12086

423 93 Cailliatte, R., *et al.* (2010) High-affinity manganese uptake by the metal transporter NRAMP1 is
424 essential for *Arabidopsis* growth in low manganese conditions. *Plant Cell* 22, 904-917

425 94 Ishimaru, Y., *et al.* (2012) OsNRAMP5, a major player for constitutive iron and manganese uptake
426 in rice. *Plant Signal. Behav.* 7, 763-766

427 95 Ishimaru, Y., *et al.* (2012) Characterizing the role of rice NRAMP5 in manganese, iron and
428 cadmium transport. *Nat. Comm.* 2, 286

429 96 Wulandari, C., *et al.* (2014) Effect of iron deficiency on root ferric chelate reductase, proton
430 extrusion, biomass production and mineral absorption of citrus root stock orange jasmine (*Murraya*
431 *exotica* L.). *J. Plant Nutr.* 37, 50-64

432 97 Sasaki, A., *et al.* (2012) Nramp5 is a major transporter responsible for manganese and cadmium
433 uptake in rice. *Plant Cell* 24, 2155-2167

434 98 Vitousek, P.M., *et al.* (2010) Terrestrial phosphorus limitation: mechanisms, implications, and
435 nitrogen-phosphorus interactions. *Ecol. Appl.* 20, 5-15

436 99 Veneklaas, E.J., *et al.* (2003) Chickpea and white lupin rhizosphere carboxylates vary with soil
437 properties and enhance phosphorus uptake. *Plant Soil* 248, 187-197

438 100 Han, W.X., *et al.* (2011) Biogeography and variability of eleven mineral elements in plant leaves
 439 across gradients of climate, soil and plant functional type in China. *Ecol. Lett.* 14, 788-796
 440 101 Mitchell, P.J., *et al.* (2008) Using multiple trait associations to define hydraulic functional types
 441 in plant communities of south-western Australia. *Oecologia* 158, 385-397
 442 102 Van der Heijden, M.G.A. and Scheublin, T.R. (2007) Functional traits in mycorrhizal ecology: their
 443 use for predicting the impact of arbuscular mycorrhizal fungal communities on plant growth and
 444 ecosystem functioning. *New Phytol.* 174, 244-250
 445 103 Oliveira, R.S., *et al.* (2015) Mineral nutrition of *campos rupestres* plant species on contrasting
 446 nutrient-impoverished soil types. *New Phytol.*
 447 104 Roelofs, R.F.R., *et al.* (2001) Exudation of carboxylates in Australian Proteaceae: chemical
 448 composition. *Plant Cell Environ.* 24, 891-904
 449 105 Zhu, Y., *et al.* (2005) A link between citrate and proton release by proteoid roots of white lupin
 450 (*Lupinus albus* L.) grown under phosphorus-deficient conditions? *Plant Cell Physiol.* 46, 892-901

451

452

453

Figure legends

Figure 1. Effects of carboxylates (and other exudates with similar effects, *e.g.*, polygalacturonate [79]) on mobilisation of phosphorus (P) and transition metals. Carboxylates (organic anions) are released via a carboxylate channel. The manner in which phosphatases are released is unknown. Carboxylates mobilise both inorganic (Pi) and organic (Po) P, which are both sorbed onto soil particles. At acid pH, Pi and Po bind to oxides and hydroxides of iron (Fe) and aluminium (Al); at alkaline pH, these compounds are precipitated by calcium (Ca). The carboxylates effectively take the place of Pi or Po, thus pushing this into solution. The released phosphatase enzymes hydrolyse Po compounds, after they have been mobilised by carboxylates. Carboxylates also mobilise some of the transition metal cations, especially Fe, manganese (Mn), zinc (Zn) and copper (Cu). Chelated Fe moves to the root surface, where it is reduced, followed by uptake via a Fe^{2+} transporter (IRT, for iron-regulated transporter). This transporter is not specific and also transports other micronutrients, such as Mn, Cu and Zn, which have been mobilised by carboxylates in soil. Alternatively, these transition metals can be taken up by a transporter referred to as NRAMP (for Natural Resistance Associated Macrophage Protein). For further explanation, see text; modified after Lambers *et al.* [13].

Figure 2. (A) Relationships between leaf manganese (Mn) concentration and total soil phosphorus (P) concentration along the Jurien Bay dune chronosequence, Western Australia [22]. Leaf Mn concentrations are calculated either at the community level (*i.e.* cover-weighted) or individually for the three non-mycorrhizal species that occurred across at least two chronosequence stages. Lines of best fit are shown for each panel. The increase in cover-weighted leaf [Mn] with declining soil [P] partly reflects the greater relative cover of non-mycorrhizal Proteaceae (which had high leaf [Mn]) in these soils. Differences in cover-weighted leaf [Mn] were only accounted for by differences in soil [P] [22], not by differences in soil pH or soil Mn availability, which is low on all dunes [80]. Soil [P] was also the strongest predictor of intra-specific differences in leaf [Mn] for the three non-mycorrhizal species shown in panels 2-4, although pH also had a significant (but smaller) effect. **(B)** Leaf Mn concentrations of *campos rupestres* sandplain species of central Brazil with different nutrient-acquisition strategies. (left) Each bar represents average values for 3-5 individual plants. An M above bars indicates arbuscular mycorrhizal species. (right) AM, arbuscular mycorrhizal; NM, non-mycorrhizal species. The central vertical bar in each box represents the median, the box represents the interquartile range, and the whiskers represent the most extreme data points that are still within 1.5 of the upper or lower quartiles. The circles outside the whiskers are values that are more than 1.5 from the upper and lower quartiles. The notches represent confidence intervals around the median. The dashed line indicates the leaf [Mn] adequate for crop growth [9].

491 **Glossary**

492 **Arbuscular mycorrhiza:** a type of mycorrhizal association that forms arbuscules or coiled hyphae
493 (highly branched exchange structures) within cortical cells of the root.

494 **Carboxylate:** organic anion; organic acid minus the proton(s). For example, citrate is the
495 carboxylate, released from the deprotonation of the organic acid, citric acid.

496 **Chelate:** compound that combines reversibly, usually with high affinity, with a metal ion (*e.g.*, iron,
497 copper or manganese).

498 **Cluster roots:** bottle-brush-like or Christmas-tree-like structures in roots with a dense packing of
499 root hairs, releasing carboxylates into the rhizosphere, thus solubilising poorly available nutrients
500 (*e.g.*, phosphate) in the soil.

501 **Ectomycorrhiza:** mycorrhizal association, mostly in woody species, in which a large part of the fungal
502 tissue is found outside the root.

503 **Heavy metal:** metal with a mass density exceeding 5 g ml⁻¹

504 **Hyperaccumulating plant species:** plants that typically accumulate 100 times more of a specific
505 heavy metal than the concentrations that occur in non-accumulator plants, growing in the same
506 substrates. For most elements, including manganese, the threshold concentration is 1,000 µg g⁻¹ dry
507 weight, except for zinc (10,000 µg g⁻¹), gold (1 µg g⁻¹) and cadmium (100 µg g⁻¹).

508 **IRT:** iron-regulated transporter associated with the uptake of iron from the rhizosphere into root
509 cells. It is not highly specific, but also transports other micronutrients.

510 **Micronutrient:** inorganic nutrients that a plant requires in relatively small quantities, *e.g.*, copper,
511 iron, manganese, molybdenum and zinc.

512 **Mycorrhiza:** structure arising from a symbiotic association between a mycorrhizal fungus and the
513 root of a higher plant (from the Greek words for fungus and root, respectively; the Greek plural
514 would be mycorrhizas, but the Latin plural (mycorrhizae) is also used).

515 **Non-mycorrhizal plant family:** plant family whose members predominantly are unable to establish a
516 symbiotic association with a mycorrhizal fungus.

517 **NRAMP (for Natural Resistance Associated Macrophage Protein):** divalent cation transporter
518 associated with the uptake of transition metals, such as copper, iron, manganese and zinc.

519 **Rhizosphere:** zone of soil influenced by the presence of a root.

520 **Scleromorphic:** containing a relatively large amount of tough structures (sclerenchyma)

521 **Sorption:** process referring to the binding, *e.g.*, of phosphate, onto the surface of (*i.e.* adsorption)
522 and inside (*i.e.* absorption) soil particles. The term was coined by McBain, going back to 1909 [81].
523 In soil science the non-committal term sorption is used to indicate all processes that result in the
524 transfer of material from the soil solution to the solid phase.

525 **Transition metal:** any metal in the d-block of the periodic table, which includes groups 3 to 12 of the
526 periodic table; the f-block lanthanide and actinide series are also considered transition metals and
527 are referred to as 'inner transition metals'.

528 **ZIP:** ZRT, IRT-like protein.

529 **ZRT:** zinc-regulated transporter.

Box 1. Manganese transport from the rhizosphere into roots

Plants use transition metal transporters to take up metals such as iron (Fe), copper (Cu), manganese (Mn), nickel (Ni), zinc (Zn) and cadmium (Cd), generally found at low concentration in the soil [82-84]. In *Arabidopsis thaliana* [36, 85], *Solanum lycopersicum* (tomato) [86] and *Oryza sativa* (rice) [87], a cation transporter (IRT1, iron-regulated transporter), has a broad substrate range (*e.g.*, Fe, Zn, Mn, Ni, and Cd) [84]. *Oryza sativa*, a Strategy II plant, which takes up Fe as a chelate, possesses several YSL (for yellow stripe-like) genes. Among them, OsYSL2 transports Fe(II)-nicotianamine as well as Mn(II)-nicotianamine [84]. Strategy II of Fe uptake is found in Gramineae [88]. In *Citrus aurantium* (Seville orange), a Strategy I species, increasing the Zn or Mn concentration in the nutrient solution decreases plant Fe concentrations; likewise, Fe inhibits the uptake of Zn and Mn [89]. In *Ulmus laevis*, leaf [Mn] increases fivefold when plants are grown under Fe-deprived conditions in nutrient solution [90].

The broad specificity of transition metal transporters [91, 92] may partly account for accumulation of Mn (and to a lesser extent Zn and Cu) when Mn is mobilised by exuded carboxylates [34]. However, there are also more specific Mn transporters, which are essential for Mn uptake from soil with low Mn availability, *e.g.*, NRAMP1 (for Natural Resistance Associated Macrophage Protein 1) in *Arabidopsis thaliana* [93] and NRAMP5 in *Oryza sativa* (rice), deployed for constitutive uptake of Fe and Mn as well as Cd [94, 95]. Citrus root stocks differ in their Fe-deficiency tolerance. *Murraya exotica* (orange jasmine) is more tolerant than *Poncirus trifoliata* var. *monstrosa* (flying dragon) [96]. The Mn concentration in *M. exotica* is independent of Fe availability, whereas the Zn concentration in its roots doubles, while that in *P. trifoliata* var. *monstrosa* increases four-fold. *Murraya exotica* appears to have mechanisms for regulating uptake of Mn, and to a lesser extent of Zn, in response to Fe deficiency [96]. We conclude that even closely related species differ in the specificity of transition-metal uptake from the rhizosphere. Manganese uptake appears to be the

554 most tightly linked with Mn availability in the rhizosphere, and thus offers the best tool to screen for
555 a P-mobilising strategy based on carboxylate release.

556 Recently, some of the genes involved in transition metal transport in plants have been identified, but
557 Mn^{2+} transport pathways are only just beginning to be unravelled at the molecular level [16]. Several
558 transporter gene families have been implicated in Mn^{2+} transport into root cells, NRAMP
559 transporters in *O. sativa* [97] and *A. thaliana* [93], and zinc-regulated transporter/iron-regulated
560 transporter (ZRT/IRT1)-related protein (ZIP) transporters in [67]. In addition, the characterization of
561 Mn hyperaccumulator plants allows the identification of genes that confer this trait [67].

562

563

Box 2. Leaf manganese concentrations as a proxy for the exudation of phosphorus-mobilising carboxylates: a tool to screen for efficient crop cultivars and belowground functional traits.

Phosphorus (P) is a macronutrient that is limiting for plant productivity in many natural and managed ecosystems [98]. To sustain crop productivity requires a continuous input of P fertilisers, which are produced from mined phosphate rock, a non-renewable resource that is gradually being depleted [4]. Therefore, there is a growing need to develop crops that are better at acquiring soil P. One such strategy is based on mycorrhizal associations; another, which is particularly effective when soil P availability is very low, is based on the exudation of P-mobilising carboxylates [21]. Measuring carboxylates in the rhizosphere is laborious. In addition, carboxylate exudation depends on climatic conditions and can occur as relatively short pulses, thus complicating sampling. We propose to use leaf [Mn] as a first step to obtain information on the carboxylate-releasing P-mobilising strategy in a range of genotypes screened for variation in P-acquisition efficiency [1]. If the results provide an indication for rhizosphere carboxylates, this can then be followed up by actual measurements of exuded carboxylates [99].

Plant ecologists are increasingly interested in functional traits, *i.e.* traits that allow grouping of species that play similar roles in an ecosystem [3, 100, 101]. Most of these traits pertain to aboveground plant characteristics, because these are easiest to measure. Belowground functional traits are mostly restricted to mycorrhizal status, the capacity to symbiotically fix dinitrogen and morphological traits [3, 102]. Traits related to nutrient acquisition are considered desirable to include, but hard to measure [3]. We propose to use leaf [Mn] as a first step to obtain information on the carboxylate-releasing P-mobilising strategy in natural environments where the P-availability is low [22]. If the results suggest that some species release carboxylates into their rhizosphere, this can then be followed up by actual measurements of exuded carboxylates in the laboratory [99].

The proposed approach is expected to be particularly promising for plants growing in alkaline soil, where carboxylate release is an effective strategy to release P and mobilise Mn [13]. It

should be equally effective on slightly acidic soils, as demonstrated along the Jurien Bay dune chronosequence in south-western Australia [22]. On more acidic soils in *campos rupestres* in Brazil, below pH 5 [51, 103], Mn availability declines with decreasing pH [74]. Here, carboxylate release is expected to be associated with increasing Mn availability only if the accompanying cations are not protons, which is a distinct possibility [104, 105].

In summary, the proposed approach looks promising for a range of soil conditions where P is a major limiting nutrient, but further experimental work is required to determine the exact soil conditions for which the approach is most useful.

Figure 1
[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)