

Treating Insomnia Symptoms with Medicinal Cannabis: A Randomized, Cross-Over Trial of the Efficacy of a Cannabinoid Medicine Compared with Placebo

Jennifer H Walsh^{1,2}
Kathleen J Maddison^{1, 2}
Tim Rankin¹
Kevin Murray³
Nigel McArdle^{1,2}
Melissa J Ree^{1,4}
David R Hillman^{1,2}
Peter R Eastwood^{1,2,5}

¹ Centre for Sleep Science, School of Human Sciences, The University of Western Australia, 35 Stirling Hwy, Crawley, Western Australia 6009

² West Australian Sleep Disorders Research Institute, Department of Pulmonary Physiology & Sleep Medicine, Sir Charles Gairdner Hospital, Hospital Avenue, Nedlands, Western Australia, 6009

³ School of Population and Global Health, The University of Western Australia, Crawley, Western Australia, 6009

⁴ School of Psychological Science, The University of Western Australia, Crawley, Western Australia, 6009

⁵ Flinders Health and Medical Research Institute, College of Medicine and Public Health, Flinders University, Adelaide, South Australia

Corresponding Author

Jennifer Walsh; Centre for Sleep Science, The University of Western Australia, M309, Perth, Western Australia, 6009.

Phone: +61 8 6488 8694

Email: Jennifer.Walsh@health.wa.gov.au

© Sleep Research Society 2021. Published by Oxford University Press on behalf of the Sleep Research Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Study Objectives: This randomized, double-blind, placebo-controlled, cross-over study was conducted to evaluate the safety and efficacy of two-weeks of nightly sublingual cannabinoid extract (ZTL-101) in treating chronic insomnia (symptoms \geq three months).

Methods: Co-primary study endpoints were safety of the medication based on adverse event reporting and global insomnia symptoms (Insomnia Severity Index; ISI). Secondary endpoints included: self-reported (sleep diary), actigraphy-derived and polysomnography measurements of sleep onset latency (SOL), wake after sleep onset (WASO), total sleep time (TST), sleep efficiency (SE); and self-reported assessments of sleep quality (sSQ) and feeling rested upon waking. Adjusted mean differences between placebo and ZTL-101 were calculated.

Results: Twenty-three of 24 randomized participants (n=20 female, mean age 53 ± 9 years) completed the protocol. No serious adverse events were reported. Forty mild, non-serious, adverse events were reported (36 during ZTL-101) with all but one resolving overnight or soon after waking. Compared to placebo, ZTL-101 decreased ISI (-5.07 units [95%CI: -7.28 to -2.86]; $p=0.0001$) and self-reported SOL (-8.45 mins [95%CI: -16.33 to -0.57]; $p=0.04$) and increased self-reported TST (64.6 mins [95%CI: 41.70 to 87.46]; $p<0.0001$), sSQ, (0.74 units [95%CI: 0.51 to 0.97]; $p<0.0001$) and feeling of being rested on waking (0.51 units [95%CI: 0.24 to 0.78]; $p=0.0007$). ZTL-101 also decreased actigraphy-derived WASO (-10.2 mins [95%CI: -16.2 to -4.2]; $p=0.002$), and increased actigraphy-derived TST (33.4 mins [95%CI: 23.07 to 43.76]; $p<0.001$) and SE (2.9% [95%CI: 2.0 to 3.8]; $p=0.005$).

Conclusion: Two-weeks of nightly sublingual administration of a cannabinoid extract (ZTL-101) is well tolerated and improves insomnia symptoms and sleep quality in individuals with chronic insomnia symptoms.

Keywords: Chronic Insomnia; Cannabinoids; Pharmacokinetics

Clinical Trial: ANZCTR; anzctr.org.au; ACTRN12618000078257

Statement of Significance:

Chronic insomnia is present in 6-15% of the population and is associated with adverse health outcomes. Current pharmacologic treatment options are often unsatisfactory. Cannabinoids have been proposed as a potential new therapeutic alternative although evidence regarding their safety and efficacy is limited. In this world-first randomized double-blind placebo-controlled cross-over trial of 24 participants with chronic insomnia, 2-weeks of ZTL-101 significantly decreased insomnia symptoms relative to placebo. While more adverse events were reported while taking ZTL-101 than placebo all were mild and the majority resolved soon after waking. While replication in a larger cohort is required before definitive conclusions can be drawn, these results suggest that the cannabinoid formulation ZTL-101 may provide a useful therapeutic option for individuals with chronic insomnia, at least for short term use.

Accepted Manuscript

Introduction

Chronic insomnia disorder, characterized by difficulty initiating or maintaining sleep at least 3 nights per week for at least 3 months, is present in 6-15% of the population,[1][2] and is associated with poor health outcomes and reduced productivity. Cognitive behavioural therapy for insomnia (CBTi) is the first-line treatment, with improvement reported in approximately 60% of patients.[3] In cases where treatment is ineffective or access to CBTi is limited or delayed, then pharmacological therapies can be useful. However, adverse effects from conventional pharmacological treatments for insomnia are common[4] and include dependence, abuse potential, tolerance, daytime sedation, psychomotor impairment manifesting as falls and cognitive impairment[5] and increased risk of head injury or fracture.[6] These undesirable side-effects drive an ongoing search for alternative therapies. Cannabinoids have emerged as a possible alternative therapy for patients with insomnia who are considering therapeutic options.

Cannabis use in the US was prohibited from 1937 until it was legalized for medical use in the late 1990s. Similar legislative changes have followed in many countries, prompting increased availability and use for medical purposes. Insomnia (or “sleep disorder”) is a common symptom for which people use cannabis.[7] However, few studies have examined the efficacy of cannabinoid formulations in treating insomnia[8] and, a placebo-controlled randomized trial has yet to be undertaken. Improvements in sleep quality have been reported with delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) alone or in combination. However, when taken in combination CBD is known to attenuate the potential psychotropic effects of THC[9] although high doses of CBD have been reported to have potential alerting properties.[10] Greater drowsiness has been reported with the addition of cannabinol (CBN) and THC than with THC alone.[11]

This study therefore employed a double-blind, randomized, placebo-controlled, cross-over design to evaluate the safety and efficacy of a cannabinoid formulation which included THC, CBD and CBN (ZTL-101), for treating insomnia symptoms in patients with chronic insomnia disorder.

Methods

Participants

Men and women aged 25-70 years presenting with chronic insomnia, defined as self-reported difficulty initiating sleep (latency to persistent sleep >30 mins) and/or maintaining sleep (>30 mins awake, or waking >30 mins before desired waking time) on ≥ 3 nights per week, for ≥ 3 months and an Insomnia Severity Index (ISI) score >10.[12] Participants were recruited via email, newspaper or television advertisement asking for volunteers to test a new medication to improve sleep quality. Only after responding to the advertisement were potential participants informed that the medication was medicinal cannabis.

Participants were excluded from the study if they: were unwilling to cease using psychotropic, CNS-depressant (confirmed at baseline with urine drug screen for opioids, amphetamine, cocaine, benzodiazepine and cannabis) or cytochrome P450 inhibitor medications for the study duration, commencing from at least 2-weeks prior to baseline assessments; had untreated cardiovascular, metabolic or significant psychopathologic disorders (self-reported); had other significant sleep disorders (self-reported or identified at baseline); or were participating in a behavioural therapy program to improve sleep (see **Supplementary Methods** for full exclusion criteria). Participants were provided with travel vouchers where required but no other compensation.

Study Design and Procedures

A double-blind, randomized, placebo-controlled, cross-over design study was conducted between May and December 2019 at The University of Western Australia's Centre for Sleep Science. Zelira Therapeutics Ltd (formerly Zeldia Therapeutics) sponsored the study, but the investigators independently designed it and collected, analyzed, and interpreted all data. Ethical review and approval of the study, which was in accordance with the guidelines of the International Council for Harmonisation and principles of the Declaration of Helsinki, was obtained from Bellberry Ltd., (HREC2017-03-226) and The University of Western Australia (RA/4/1/9236). Informed written consent was obtained from each participant following full explanation of procedures, risks and benefits.

Preliminary eligibility was established during a clinical interview at a screening and consent visit, which was followed by a two-week baseline period (**Figure 1**) during which participants wore a wrist-based activity monitor (GT9X, ActiGraph, Pensacola, USA) and completed sleep diaries.[13] On the 14th baseline night a laboratory-based polysomnography (PSG) study (Graef, Compumedics, Sydney, Australia) was performed; participants were encouraged to attempt to sleep at their usual time and were not woken by others in the morning. The PSG montage aligned with AASM criteria[14] and studies were scored according to these criteria by a single experienced scorer who was blinded to treatment.

Potential sensitivity, including anaphalaxis, adverse cardiovascular, hallucinogenic or paranoid responses, to a small dose of the study medication (THC/CBN/CBD; 3/0.3/0.15mg + 0.15ml placebo) was assessed from a 3-hour observation period on the morning following this baseline PSG.

Eligible participants then underwent a one-week washout period following which they were randomly allocated to ZTL-101 or placebo for two-weeks (see **Supplementary Methods** for further detail on randomization procedure). Following a further one-week washout period, participants

crossed-over to the alternate study arm for a further two-weeks. All participants and investigators were blinded as to whether ZTL-101 or placebo was administered. Wrist-based activity monitors were worn throughout each study arm with PSG studies performed on the 14th night of each arm. During each of the two-week study arms, participants sublingually self-administered 0.5ml of ZTL-101 or placebo one, hour prior to their desired sleep time. With physician approval, participants were allowed, but not required, to double the dose (*i.e.*, increase to two syringes of ZTL-101 or placebo = 1.0ml sublingually) from the 4th night of each two-week study period. Further increases in dose were not permitted, although dose decreases were. Participants were contacted daily for the first three days of each two-week study period, and following dose increases, to monitor adverse events. Participants were supplied with sufficient doses for one week at a time and all used and unused syringes were returned to the study site to confirm doses taken.

ZTL-101 contained delta-9-tetrahydrocannabinol (THC) 20mg/ml, cannabidiol (CBD) 1mg/ml and naturally occurring terpenes, extracted from the cannabis plant, in pharmaceutical grade sunflower oil as the diluent. The investigational product was manufactured to an approved specification in a good manufacturing practice certified facility (Eurofins PROXY Laboratories, Leiden, Netherlands). The placebo contained the same terpenes, but no cannabinoids, extracted from the same cannabis plant, to match ZTL-101 as closely as possible for smell, taste and colour (Eurofins PROXY Laboratories, Leiden, Netherlands).

Primary Outcomes

The co-primary outcome measures were: (i) frequency, type and severity of adverse events during each of the two-week treatment periods; and (ii) global insomnia symptoms as assessed by the ISI on the 14th night of each two-week treatment period.

Secondary Outcomes

Measures of sleep quality and quantity were obtained from the following methods: (i) self-report from sleep diary (*s*); (ii) actigraphy (*a*) and; (iii) PSG. Measures of sleep quality/quantity derived from each method, where possible, included sleep onset latency (SOL: time from lights off to falling asleep); total sleep time (TST: total time spent asleep overnight); wake after sleep onset (WASO: time spent awake after initially falling asleep until final out of bed time); sleep efficiency (SE: proportion of time spent asleep between the period of lights out and out of bed time) and awakening index (AI: number of awakenings per hour of sleep from lights out to out of bed time). A rating of perceived sleep quality (*s*SQ) and feeling rested/refreshed on waking was also recorded. Self-report and actigraphy measures were calculated from the mean of each of the two-week baseline and treatment periods. The proportion of the sleep period spent in sleep stages N1, N2, N3 and REM were determined from PSG. Additional detail regarding each outcome measure is provided in **Supplementary Methods**.

A Maintenance of Blinding Questionnaire[15] was administered on the 14th night of each study arm to assess the success of participant blinding to treatment.

Pharmacokinetic Study

All participants who completed both two-week study periods were invited to an overnight study during Feb-Mar 2020 to examine the 12-hour pharmacokinetic responses to a single dose (i.e., THC/CBN/CBD; 10/1/0.5mg), and in some participants, a double dose of ZTL-101 (i.e., THC/CBN/CBD; 20/2/1mg). Venous blood samples were drawn at 0,1,2,4,6,8,10 and 12 hours following dosing and the plasma was analysed for CBD, CBN, THC and the metabolite carboxy-THC (THC-COOH) (see **Supplementary Methods** for further detail on pharmacokinetic study).

Statistical Analysis

Sample Size

No previous study had reported changes in ISI, the primary outcome measure, following cannabinoid therapy for insomnia. Therefore, a sample size calculation was undertaken based on the assumption that a clinically meaningful improvement in ISI was 6 points.[16] Assuming a SD of 6,[17] alpha level of 0.05 and power of 0.9, a minimum of 13 participants were required to detect this difference. A separate sample size calculation for the secondary outcome measure of SE from PSG was also performed. In order to detect an increase in SE of 20 with a SD of 25,[18] power of 0.9 and alpha of 0.05, a minimum of 19 participants was required. To ensure a sufficient sample size to detect both primary and secondary outcome measures in *per protocol* analysis and to allow for a 20% attrition rate the planned sample size was 24 participants. Due to the high prevalence of OSA in the general population, approximately 40% of participants were expected to be excluded following the baseline PSG. Thus, we expected to consent a total of 40 participants, and exclude approximately 16 following consent due to the presence of OSA.

Linear mixed models were used to analyse differences in responses between the two study arms. Fixed effects of treatment (placebo vs ZTL-101) and order and period of treatment were included, along with random individual effects. Estimated differences in least squares means were calculated along with standard errors and confidence intervals. Cohen's *d* (effect size) was calculated based on the method for paired data.[19] Whilst multiple tests were undertaken, and all results presented, the *p*-values presented have not been adjusted for multiplicity.

One participant did not complete the baseline ISI questionnaire. Screening data was imputed for this participant's baseline data. One participant did not complete question three of the seven item ISI questionnaire on the first dosing arm (placebo). A mean of screening and baseline data (which were identical) was imputed for this question. No other data were missing.

Unless specified, data are presented as adjusted mean±SD and adjusted mean difference [95% CI].

Results

Of 167 individuals considered for inclusion, 85 were ineligible due to concomitant medications (n=35), age (n=23), other sleep disorders (n=11), comorbidities (n=4), currently participating in CBTi (n=4), ISI<10 (n=4), self-exclusion for unknown reason (n=3) or history of falling asleep while driving (n=1). In addition, 41 individuals declined participation due to lack of availability (n=31) or unwillingness to participate in a pharmaceutical trial (n=10) leaving 41 participants who consented for the trial. A total of 15 were excluded: 12 with significant other sleep disorders; two with suicidal ideation; and one with contraindicated medication use. A further two participants withdrew prior to initial dosing. Twenty-four participants (n=20 female, mean age 53±9years) proceeded to dosing (**Figure 1**). One female participant withdrew after the 4th night of ZTL-101 due to non-serious adverse events (xerostomia, oral hypoesthesia, swollen tongue, nausea). Twenty-three, predominantly Caucasian (n=1 Asian) participants completed the full protocol (n=19 female; n=5 premenopausal; mean age 52±9years; mean BMI 24.8±3.6kg.m²).

Of the 23 participants who completed the protocol, 12 (52%) were taking a double dose of ZTL-101 on the 14th night. Sixteen (69.5%) were taking a double dose of placebo on the 14th night. Twenty-one of 21 participants (100%) (n=2 missing data) guessed that they were receiving the active medication when taking ZTL-101. “*Improvement in sleep quality*” was the reason for their guess in 17 (81%) with adverse reactions the reason in the remaining four participants. When taking the placebo, 18 of 23 participants (78%) thought they were receiving placebo. Sixteen noted “*lack of improvement in sleep quality*” as the reason for their guess that they were receiving placebo.

Primary Outcomes

Safety

No serious adverse events were reported. During ZTL-101 dosing, 36 non-serious adverse events possibly or likely related to ZTL-101 medication were recorded from 17 participants. Four non-serious adverse events were recorded from four participants during dosing with the placebo medication and one non-serious adverse event was reported during sensitivity testing (**Table 1**). All adverse events were classified as mild and had either resolved overnight or soon after waking (97.5%).

Insomnia Symptoms

ISI scores at the end of two-weeks of ZTL-101 were significantly lower than scores following two-weeks of placebo (adjusted mean difference -5.1 [95%CI -2.9 to -7.3], $p=0.0001$, $d=0.94$; **Figure 2**; **Table 2**). ISI scores for the 12 participants who took a single dose of ZTL-101 and 11 participants who took a double dose are presented in **Supplementary Results, Table S1**.

Secondary Outcomes

Self-reported measures from sleep diary

When taking ZTL-101 vs placebo participants perceived that they went to sleep faster (sSOL) (8.4mins [95%CI -16.3 to -0.6], $p=0.0369$, $d=0.44$), slept for longer (sTST) (64.6mins [95%CI 41.7 to 87.5], $p<0.0001$, $d=1.27$), had improved sleep quality (sSQ) (0.7 [95%CI 0.5 to 1.0], $p<0.0001$, $d=1.37$) and felt more rested/refreshed on waking (0.5 [95%CI 0.2 to 0.8], $p=0.0007$, $d=0.83$) (**Table 2**).

Actigraphy measures

Relative to placebo, ZTL-101 significantly decreased $aWASO$ (-10.2mins [95%CI -16.2 to -4.2], $p=0.0021$, $d=0.68$) and increased $aTST$ (33.4mins [95%CI 23.1 to 43.8], $p<0.001$, $d=1.44$) and aSE (2.9% [95%CI 2.0 to 3.8], $p<0.001$, $d=1.28$) (**Table 2**). Measurements of $aSOL$ (-0.4mins [95%CI -1.2 to 0.4], $p=0.2825$, $d=0.22$) and aAI (0.1 [95%CI -1.6 to 1.8, $p=0.8829$, $d=0.03$]) were unchanged by ZTL-101.

Polysomnography measures

Relative to placebo, ZTL-101 had minimal effect on SOL (0.42mins [95%CI -26.4 to 27.3], $p=0.9743$, $d=0.00$), $WASO$ (15.1mins [95%CI -7.9 to 38.1], $p=0.1872$, $d=0.26$), TST (-3.5mins [95%CI -38.9 to 31.9], $p=0.8396$, $d=0.03$) and SE (-2.4% [95%CI -8.4 to 3.6], $p=0.4124$, $d=0.15$). ZTL-101 had minimal effect on the number of awakenings (4.1 [95%CI -0.4 to 8.6], $p=0.0709$, $d=0.40$) (**Table 2**), proportion of sleep stages or severity of obstructive sleep apnea or periodic limb movements (**Table S2**), compared with placebo.

Pharmacokinetic measures

Pharmacokinetic measures were obtained in all those available to return for an additional night of testing. Pharmacokinetic measures in response to a single dose of ZTL-101 were obtained in nine participants. Time at which maximal plasma concentration was reached (T_{max}) occurred at 4-6 hours for CBD, CBN, THC-COOH and THC (**Figure 3**).

Pharmacokinetic measures in response to a double dose of ZTL-101 ($n=2$) are displayed in **Figure S1** and additional pharmacokinetic parameters are shown **Table S3**.

Discussion

This Phase 1b study demonstrated that nightly sublingual administration of a novel cannabinoid formulation for two-weeks improved insomnia symptoms without significant adverse events in participants with chronic insomnia symptoms. This study used the ISI, a reliable, valid and widely used instrument, to quantify perceived insomnia severity and its impact on daily function.[17] The ISI was lower while taking ZTL-101 relative to placebo. Consistent with this positive treatment response, improvements in self-reported sleep diary and objective actigraphic measures of sleep quality and quantity were also documented.

The novel formulation, ZTL-101, was well tolerated with only one participant withdrawal, due to a non-serious adverse event. Seventeen of the 24 participants experienced at least one adverse event while taking ZTL-101, with dry mouth and dizziness being most frequently reported. Although the number of adverse events is somewhat greater than that commonly reported for contemporary hypnotics[20] it is comparable to other trials using medicinal cannabis.[21] Furthermore, all adverse events were classified as mild and all but one (xerostomia, oral hypoesthesia, swollen tongue, nausea in the participant who withdrew) had resolved upon waking or soon afterwards. The occurrence of dizziness and hallucinations are the most concerning and suggest caution is required in populations such as the elderly or those with psychiatric disorders. Although the occurrence of adverse events may be reduced with a more gradual dose titration [22] it is clear that further research is required to more comprehensively assess benefits and harms of medicinal cannabis use for the treatment of insomnia.

The effects of ZTL-101 on sleep were assessed from self-report, actigraphy and PSG. Each of these methods is widely used to assess sleep, although none are without limitations in individuals with insomnia. For example, while self-reported sleep difficulty is the basis of the clinical diagnosis of insomnia, self-reported measures of total time spent asleep and time taken to fall asleep at the start

of the night are known to be under and overestimated, respectively.[23] Actigraphy can objectively and unobtrusively measure sleep over multiple nights in the home environment. However, it tends to overestimate sleep time and underestimate wake time because the method designates periods of no motion as sleep, whether asleep or not. Polysomnography is considered the 'gold-standard' method of defining sleep, wake, respiration and movement, and can therefore provide an objective measure of sleep quality, quantity and identify other sleep disorders. It is typically performed in a laboratory or clinic on a single night and is more intrusive than actigraphy. However, its use in individuals with insomnia is limited by first night effects (*i.e.*, patients sleeping worse or better than usual) and an inability to capture night-to-night variability in sleep behaviour with a single night measurement. To enable a thorough evaluation of the impact of ZTL-101 on sleep, the present study utilized all three methods.

When taking ZTL-101 participants reported an improvement in the time taken to fall asleep, time spent asleep and feelings of being more rested/refreshed on waking. These self-reported improvements were supported by actigraphy-derived measures of the mean total time spent asleep each night, which increased by 33.5mins; sleep efficiency, which increased by 2.9% to 84.8%; and the time spent awake during the night, which decreased by 10mins. The mean total time spent asleep across the two-week period was over 7 hours while taking ZTL-101, which is the recommended minimum sleep duration for adults,[24] and above average for individuals of comparable age without insomnia.[25] Despite this, the average time spent awake during the night remained high at greater than 70mins, possibly reflecting a persisting tendency to disturbed sleep in individuals with chronic insomnia or that sufficient sleep had been achieved given the other improvements. Notably, the time taken to fall asleep was unchanged by ZTL-101. This has been reported in other studies[26] and could reflect a "floor effect" given the relatively short α SOL values at baseline in the study participants; or the inability of actigraphy to differentiate motionless wakefulness from sleep.

The improvements in self-reported and actigraphy-based measures of sleep with ZTL-101 were not seen in any PSG measure of sleep quantity or quality. This is likely due, at least in part, to the inclusion criteria being based on self-reported diagnostic criteria for insomnia[27] rather than PSG-defined insomnia, as well as the limitations of a single night PSG measure described previously. The main purpose of PSG was to identify and exclude participants with other sleep disorders such as sleep apnea and periodic limb movement disorder, which can coexist with insomnia and potentially confound interpretation of any sleep-related changes ascribed to ZTL-101. It is important to also note that ZTL-101 did not induce these sleep disorders, nor did it alter the proportion of time spent in the different sleep stages. By contrast, alteration of the proportion of sleep stages is common with many hypnotic medications; benzodiazepines, for example, are known to significantly decrease the proportion of REM sleep.[28]

Insomnia is characterized by self-reported difficulties initiating or maintaining sleep.[2] As such, any globally effective pharmacological therapy for it should be capable of targeting either characteristic. To explore this, the pharmacokinetic properties of ZTL-101 overnight were determined after a standardized evening meal, thereby obtaining measurements of drug metabolism under conditions pertaining to its use in the clinical setting. Following a single dose of ZTL-101, peak plasma levels of the major constituents were reached at approximately 4-6 hours. A similar profile was observed following ingestion of a double dose of ZTL-101, albeit with greater maximum plasma levels. Although it is possible that accumulation of cannabinoids might occur with dosing over multiple nights, this has not been demonstrated in dosing with THC/CBD = 21.6/20.0mg out to 9 days. These data suggest that for patients with sleep onset insomnia dosing 2-4 hours before desired bedtime might be optimal, while those with sleep maintenance insomnia should dose one hour before desired bedtime.

Limitations

Due to the possible risk of exacerbating pre-existing conditions, individuals with a history of significant cardiovascular disease or known major psychopathology were excluded. The safety and efficacy of ZTL-101 in these populations remains to be established. Likewise, the possible drug-drug interaction between ZTL-101 and cytochrome P450 inhibitors requires further investigation.

Maintenance of blinding is challenging in hypnotic[29] and cannabinoid trials[30] and this study is one of the first randomized, placebo-controlled medicinal cannabis trials to assess efficacy of blinding. The results of the blinding questionnaire suggest that despite considerable efforts to ensure ZTL-101 and placebo were similar in appearance, odour and taste, effective blinding for a beneficial treatment effect was unable to be achieved; thus the data should be interpreted with this in mind. This is the first study to demonstrate acceptable safety and promising efficacy of a cannabinoid therapy in a randomized, double-blind, placebo-controlled manner, although it was limited to two-weeks and included a relatively small sample of participants. Similarly, the pharmacokinetic component of the study used a subset of participants (n=9), which although small is not unusual for pharmacokinetic studies.[31,32]

We undertook a multi-faceted study of the influence of ZTL-101 on sleep and wakeful function, examining for a consistent trend across many measures beyond the ISI, the co-primary outcome measure. Adjustment for multiple comparisons was not performed as we did not want to artificially discount any of the exploratory analyses, which are all presented.[33] Further dedicated studies are needed to confirm the promising findings of this preliminary investigation of a novel pharmacological approach to a very common and vexing health issue.

Conclusions

This study has demonstrated that ZTL-101, a novel cannabinoid therapy, is well tolerated and improves insomnia symptoms and sleep quality in individuals with chronic insomnia symptoms.

These improvements, observed over a two-week dosing period, are encouraging and support further investigation of ZTL-101 for the treatment of insomnia in studies with larger sample sizes.

Accepted Manuscript

Data Availability Statement

The data that support the findings of this article are available from the corresponding author (JW), upon reasonable request.

Disclosure Statements:

Financial Disclosure: This clinical trial was funded by Zelira Therapeutics Limited although the design and conduct of the study, including analyses, was performed entirely independent of the sponsor. PRE was funded by a National Health and Medical Research Council Senior Research Fellowship [1136548].

Non- Financial Disclosure: nothing to declare.

Accepted Manuscript

References

1. Reynolds AC, Appleton SL, Gill TK, Adams RJ. *Chronic insomnia disorder in Australia: A report to the Sleep Health Foundation*. <https://www.sleephealthfoundation.org.au/news/special-reports/chronic-insomnia-disorder-in-australia.html>2019. Accessed 29 July 2020
2. Morin CM, Benca R. Chronic insomnia. *Lancet*. 2012; 379 (9821): 1129-1141.
3. Morin CM, Vallieres A, Guay B, et al. Cognitive behavioral therapy, singly and combined with medication, for persistent insomnia: a randomized controlled trial. *JAMA*. 2009; 301 (19): 2005-2015.
4. Herring WJ, Connor KM, Ivgy-May N, et al. Suvorexant in Patients With Insomnia: Results From Two 3-Month Randomized Controlled Clinical Trials. *Biological psychiatry*. 2016; 79 (2): 136-148.
5. Schroeck JL, Ford J, Conway EL, et al. Review of Safety and Efficacy of Sleep Medicines in Older Adults. *Clin Ther*. 2016; 37 (11): 2340-2372.
6. Lai MM, Lin CC, Lin CC, Liu CS, Li TC, Kao CH. Long-term use of zolpidem increases the risk of major injury: a population-based cohort study. *Mayo Clinic proceedings*. 2014; 89 (5): 589-594.
7. Hazekamp A, Ware MA, Muller-Vahl KR, Abrams D, Grotenhermen F. The medicinal use of cannabis and cannabinoids-an international cross-sectional survey on administration forms. *J Psychoactive Drugs*. 2013; 45 (3): 199-210.
8. Surav AS, Marshall NS, Vandrey R, et al. Cannabinoid therapies in the management of sleep disorders: A systematic review of preclinical and clinical studies. *Sleep medicine reviews*. 2020; 53: 101339.
9. Bhattacharyya S, Morrison PD, Fusar-Poli P, et al. Opposite effects of delta-9-tetrahydrocannabinol and cannabidiol on human brain function and psychopathology. *Neuropsychopharmacology*. 2010; 35 (3): 764-774.
10. Nicholson AN, Turner C, Stone BM, Robson PJ. Effect of Delta-9-tetrahydrocannabinol and cannabidiol on nocturnal sleep and early-morning behavior in young adults. *Journal of clinical psychopharmacology*. 2004; 24 (3): 305-313.
11. Karniol IG, Shirakawa I, Takahashi RN, Knobel E, Musty RE. Effects of delta9-tetrahydrocannabinol and cannabiniol in man. *Pharmacology*. 1975; 13 (6): 502-512.
12. Medicine AAoS. *International classification of sleep disorders*. 3rd Edition ed. Darien, IL: American Academy of Sleep Medicine; 2014.
13. Carney CE, Buysse DJ, Ancoli-Israel S, et al. The consensus sleep diary: standardizing prospective sleep self-monitoring. *Sleep*. 2012; 35 (2): 287-302.
14. Berry RB, C.L. A, Harding SM, Medicine. eaftAAoS. *The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications. Version 2.5*. Darien, IL: American Academy of Sleep Medicine; 2018.

15. Moscucci M, Byrne L, Weintraub M, Cox C. Blinding, unblinding, and the placebo effect: an analysis of patients' guesses of treatment assignment in a double-blind clinical trial. *Clinical pharmacology and therapeutics*. 1987; 41 (3): 259-265.
16. Yang M, Morin CM, Schaefer K, Wallenstein GV. Interpreting score differences in the Insomnia Severity Index: using health-related outcomes to define the minimally important difference. *Curr Med Res Opin*. 2009; 25 (10): 2487-2494.
17. Bastien CH, Vallieres A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep medicine*. 2001; 2 (4): 297-307.
18. Monti JM, Alvarino F, Monti D. Conventional and power spectrum analysis of the effects of zolpidem on sleep EEG in patients with chronic primary insomnia. *Sleep*. 2000; 23 (8): 1075-1084.
19. Sawilowsky SS. New effect sizes rules of thumb. *Journal of Modern Applied Statistical Methods*. 2009; 8 (2): 597-599.
20. Sateia MJ, Buysse DJ, Krystal AD, Neubauer DN, Heald JL. Clinical Practice Guideline for the Pharmacologic Treatment of Chronic Insomnia in Adults: An American Academy of Sleep Medicine Clinical Practice Guideline. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*. 2017; 13 (2): 307-349.
21. Wang T, Collet JP, Shapiro S, Ware MA. Adverse effects of medical cannabinoids: a systematic review. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2008; 178 (13): 1669-1678.
22. MacCallum CA, Russo EB. Practical considerations in medical cannabis administration and dosing. *Eur J Intern Med*. 2018; 49: 12-19.
23. Mercer JD, Bootzin RR, Lack LC. Insomniacs' perception of wake instead of sleep. *Sleep*. 2002; 25 (5): 564-571.
24. Consensus Conference P, Watson NF, Badr MS, et al. Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society on the Recommended Amount of Sleep for a Healthy Adult: Methodology and Discussion. *Sleep*. 2015; 38 (8): 1161-1183.
25. Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV. Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep*. 2004; 27 (7): 1255-1273.
26. Martin JL, Hakim AD. Wrist actigraphy. *Chest*. 2011; 139 (6): 1514-1527.
27. Medicine AAoS. *International classification of sleep disorders, 3rd ed*. Carien, IL: American Academy of Sleep Medicine; 2014.
28. Roth T, Zorick F, Sicklesteel J, Stepanski E. Effects of benzodiazepines on sleep and wakefulness. *British journal of clinical pharmacology*. 1981; 11 Suppl 1: 31S-35S.
29. McCall WV, Benca RM, Rumble ME, Krystal AD. Blinding and bias in a hypnotic clinical trial. *Hum Psychopharmacol*. 2021; 36 (1): 1-5.

30. Casarett D. The Achilles Heel of Medical Cannabis Research-Inadequate Blinding of Placebo-Controlled Trials. *JAMA Intern Med.* 2018; 178 (1): 9-10.
31. Fernandez-Trapero M, Perez-Diaz C, Espejo-Porras F, de Lago E, Fernandez-Ruiz J. Pharmacokinetics of Sativex(R) in Dogs: Towards a Potential Cannabinoid-Based Therapy for Canine Disorders. *Biomolecules.* 2020; 10 (2).
32. Karschner EL, Darwin WD, Goodwin RS, Wright S, Huestis MA. Plasma cannabinoid pharmacokinetics following controlled oral delta9-tetrahydrocannabinol and oromucosal cannabis extract administration. *Clin Chem.* 2011; 57 (1): 66-75.
33. Althouse AD. Adjust for Multiple Comparisons? It's Not That Simple. *The Annals of thoracic surgery.* 2016; 101 (5): 1644-1645.

Accepted Manuscript

List of Figures

Figure 1. CONSORT flow diagram of participants at each phase of the trial

Figure 2. Individual (A) and group mean (\pm SD) (B) Insomnia Severity Index scores for all participants (n=23) at baseline and during treatment with ZTL-101 and placebo.

Figure 3. Mean plasma concentrations (\pm SE) for (A) CBD, (B) CBN, (C) THC and (D) THC-COOH over time after administration of a single dose (n=9) of ZTL-101.

Hours since time of administration (Hours) and mean time of day (hh:mm) are shown on the x-axis.

CBD, cannabidiol; CBN, cannabinol; THC, delta-9-tetrahydrocannabinol; THC-COOH, carboxy-THC.

List of Tables

Table 1. Adverse events reported during ZTL-101 and placebo dosing ^a

Table 2. Group mean^a measures of insomnia symptoms and sleep quantity and quality at baseline and during ZTL-101 and placebo dosing (n=23).

Accepted Manuscript

Table 1. Adverse events reported during ZTL-101 and placebo dosing ^a

Adverse Event	ZTL-101 (n=24)	Placebo (n=23)	Sensitivity (n=24)
Headache	4 (16.7)	2 (8.7)	0 (0)
Xerostomia	8 (33.3)	0 (0)	1 (4.2)
Dizziness	6 (25.0)	1 (4.2)	0 (0)
Feeling abnormal	4 (16.7)	0 (0)	0 (0)
Dry eye	2 (8.3)	0 (0)	0 (0)
Palpitations	2 (8.3)	0 (0)	0 (0)
Nausea	2 (8.3)	0 (0)	0 (0)
Ataxia	2 (8.3)	0 (0)	0 (0)
Mood variable	1 (4.2)	1 (4.2)	0 (0)
Tachyphrenia	1 (4.2)	0 (0)	0 (0)
Auditory hallucinations	1 (4.2)	0 (0)	0 (0)
Visual hallucinations	1 (4.2)	0 (0)	0 (0)
Oral hypoesthesia	1 (4.2)	0 (0)	0 (0)
Swollen tongue	1 (4.2)	0 (0)	0 (0)

^a All values expressed as number and (percentage) of participants

Table 2. Group mean^a measures of insomnia symptoms and sleep quantity and quality at baseline and during ZTL-101 and placebo dosing (n=23).

	Baseline	ZTL-101	Placebo	p-value ^b	Effect size ^c <i>d</i>
Insomnia Symptoms					
ISI (score)	18.0 ± 3.7	12.9 ± 5.3	18.0 ± 4.3	0.0001	0.94
Self-reported measures – Sleep Diary					
sSOL (mins)	47.6 ± 38.0	38.1 ± 28.0	46.9 ± 34.4	0.0369	0.44
sTST (mins)	300.5 ± 82.6	366.7 ± 74.5	304.1 ± 81.0	<0.001	1.27
sSQ ^d	2.5 ± 0.5	3.2 ± 0.6	2.5 ± 0.5	<0.001	1.37
Rested on Waking ^e	1.2 ± 0.4	1.8 ± 0.6	1.2 ± 0.5	0.0007	0.83
Actigraphy					
aSOL (mins)	4.7 ± 1.3	4.8 ± 0.9	5.2 ± 1.9	0.2825	0.22
aWASO (mins)	84.3 ± 23.2	71.8 ± 23.9	82.3 ± 27.7	0.0021	0.68
aTST (mins)	398.9 ± 33.8	424.7 ± 34.7	391.2 ± 35.5	<0.001	1.44
aSE (%)	81.8 ± 4.3	84.8 ± 4.3	81.9 ± 4.9	<0.001	1.28
aAI (n)	24.8 ± 5.8	23.4 ± 6.0	23.6 ± 6.2	0.1236	0.03
Polysomnography					
SOL (mins)	12.3 ± 11.4	25.5 ± 22.0	25.4 ± 52.7	0.9743	0.00
WASO (mins)	108.8 ± 49.5	96.2 ± 65.9	82.0 ± 51.0	0.1872	0.26
TST (mins)	375.0 ± 68.4	384.3 ± 82.0	386.9 ± 72.0	0.8396	0.03
SE (%)	75.6 ± 10.6	75.8 ± 14.2	78.0 ± 12.7	0.4124	0.15
AI (n)	23.0 ± 8.7	22.7 ± 8.2	18.4 ± 5.6	0.0709	0.40

ISI, Insomnia Severity Index (ISI); TST, total sleep time; SE, sleep efficiency; SOL, sleep onset latency; WASO, wake after sleep onset; AI, awakening index; sQS, self-reported quality of sleep.

^a unadjusted mean ± SD

^b p-value based on linear mixed model analysis of treatment responses for ZTL-101 vs placebo

^c Effect size for ZTL-101 vs placebo

^d Rating of sleep quality 0=very poor, 1=poor, 2=fair, 3=good, 4=very good

^e Rating of feeling rested/refreshed upon waking 0=not at all rested, 1=slightly rested, 2=somewhat rested, 3=well-rested, 4=very well-rested

Figure 1






Figure 2



Accepted Manuscript

Figure 3



Acc

191