

RUNNING HEADER: High AQ adults do not see angry faces in a RSVP

Threatening faces fail to guide attention for adults with autistic-like traits.

Michael C.W. English¹, Murray T. Maybery¹, Troy A.W. Visser¹

¹School of Psychology, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

Correspondence concerning this article should be addressed to:

Michael C.W. English.

School of Psychology, The University of Western Australia, Crawley, WA 6009, Australia.

Email: michael.english@uwa.edu.au. Telephone: +61 8 6488 1421.

Number of text pages: 10 pages (22 pages inc. references)

Number of Tables: 3

Number of Figures: 2

Grant sponsor: Australian Research Council; Grant Number(s): DP120102313, DP120104713

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/aur.1658](https://doi.org/10.1002/aur.1658).

RUNNING HEADER: High AQ adults do not see angry faces in a RSVP

Threatening faces fail to guide attention for adults with autistic-like traits.

Michael C.W. English, Murray T. Maybery, Troy A.W. Visser

Scientific Abstract

Individuals diagnosed with autistic spectrum conditions often show deficits in processing emotional faces relative to neurotypical peers. However, little is known about whether similar deficits exist in neurotypical individuals who show high-levels of autistic-like traits. To address this question, we compared performance on an attentional blink task in a large sample of adults who showed low- or high-levels of autistic-like traits on the Autism Spectrum Quotient. We found that threatening faces inserted as the second target in a rapid serial visual presentation were identified more accurately amongst individuals with low- compared to high-levels of autistic-like traits. This is the first study to show that attentional blink abnormalities seen in autism extend to the neurotypical population with autistic-like traits, adding to the growing body of research suggesting that autistic-related patterns of behaviours extend into a subset of the neurotypical population.

Lay Abstract

The majority of the neurotypical population detect threatening stimuli, such as angry faces, faster and more accurately than faces with neutral or positive emotional expressions. The ability for the threatening stimuli to more efficiently guide attention relative to less threatening stimuli is believed to reflect an evolutionary advantage, such that individuals can swiftly respond to threatening or dangerous situations. However, there is evidence to suggest that individuals diagnosed with autistic spectrum conditions fail to show this 'anger superiority effect' when the stimuli are presented for very brief periods of time (less than 20ms). In our study, we found that neurotypical adults with few

RUNNING HEADER: High AQ adults do not see angry faces in a RSVP

autistic-like traits (measured with the Autism Spectrum Quotient) were able to detect angry faces more often than neutral faces, whereas neurotypical individuals who reported higher levels of autistic-like traits did not detect angry faces any more often than neutral faces in the most difficult condition. This is further evidence in a growing field of research that suggests that some attentional abnormalities expressed by individuals with autism spectrum conditions may also be present in a subset of the neurotypical population.

Keywords

autistic-like traits; attentional blink; emotion; faces

Accepted Article

Introduction

Because resource limitations prevent us from simultaneously processing all of the stimuli available to sensory systems, attention mechanisms must be employed to select a subset of information based on endogenous and exogenous factors. One particularly important exogenous factor in guiding attention is the level of emotional arousal (both positive and negative) elicited by incoming information. An abundance of evidence suggests arousing stimuli are responded to more quickly and accurately than neutral stimuli (e.g., Algom, Chajut, & Lev, 2004; Anderson, 2005; Mather & Nesmith, 2008; Mathews & MacLeod, 1985; Schimmack, 2005; Vogt, De Houwer, Koster, Van Damme, & Crombez, 2008), indicating they are more rapidly attended. Moreover, compared to neutral images, emotionally-arousing visual images reliably increase activation in areas such as the amygdala, prefrontal cortex (PFC) and occipital regions (Costafreda, Brammer, David, & Fu, 2008; Fusar-Poli et al., 2009; Phan, Wager, Taylor, & Liberzon, 2002).

Faces are a common source of both positively- and negatively-valenced emotional arousal, and have been reliably shown to guide attention, most commonly illustrated by faster reaction times to identifying a target emotional face amongst neutral expression distractors (Frischen, Eastwood, & Smilek, 2008). For example, in the well-known “anger superiority effect” (Hansen & Hansen, 1988), participants reliably identify the presence of an individual with an angry face within an array of individuals with neutral or happy expressions faster than they identify the presence of a happy face within an array of angry or neutral faces (Eastwood, Smilek, & Merikle, 2001; Fox et al., 2000; Hansen & Hansen, 1988; Hershler & Hochstein, 2005; Öhman, Lundqvist, & Esteves, 2001). As angry or threatening faces are a critical signal of potential social conflicts, the ability to attend and respond to them swiftly is often theorised to benefit the observing individual in attempting to avoid the conflict (Öhman & Dimberg, 1984).

Individuals diagnosed with neurological disorders such as autism spectrum disorders (ASD) often exhibit relatively greater difficulty with facial and emotional recognition than their

High AQ adults do not see angry faces in a RSVP

2

neurotypical counterparts (Adolphs, Sears, & Piven, 2001; Bölte & Poustka, 2003; Celani, Battacchi, & Arcidiacono, 1999; Farran, Branson, & King, 2011; Grossman, Klin, Carter, & Volkmar, 2000; Howard et al., 2000; Humphreys, Minshew, Leonard, & Behrmann, 2007; Pelphrey et al., 2002; see Uljarevic & Hamilton, 2013, for a review), which may in part stem from a failure to attend to the more socially meaningful aspects of the face, such as the eyes (Baron-Cohen, Wheelwright, & Jolliffe, 1997; Corden, Chilvers, & Skuse, 2008a; Dalton et al., 2005; Jones, Carr, & Klin, 2008; Kirchner, Hatri, Heekeren, & Dziobek, 2011; Kliemann, Dziobek, Hatri, Baudewig, & Heekeren, 2012; Ami Klin, Jones, Schultz, Volkmar, & Cohen, 2002; Spezio, Adolphs, Hurley, & Piven, 2007). This is not to say that individuals with ASD are universally impaired when processing facial information. Krysko and Rutherford (2009), for example, demonstrated that ASD individuals show a relatively intact anger superiority effect, detecting angry faces in a visual search paradigm as quickly as their neurotypical peers (although with less accuracy). That said, however, even when performance is similar between ASD and neurotypical samples, there is evidence to suggest that compensatory mechanisms and atypical perceptual processing strategies are present in ASD participants (see Harms, Martin, & Wallace, 2010, for a review).

It is possible that the diminished ability to attend to emotional stimuli also manifests itself in parts of the neurotypical population. An increasing body of research suggests that behavioural characteristics generally associated with ASD are also present in neurotypical individuals who demonstrate above-average scores on measures of autistic-like traits, such as the Autism Spectrum Quotient (AQ) (Almeida, Dickinson, Maybery, Badcock, & Badcock, 2010a, 2010b, 2013; Baron-Cohen, Wheelwright, Skinner, Martin, & Clubley, 2001; Grinter, Van Beek, Maybery, & Badcock, 2009; Grinter, Maybery, et al., 2009; Rhodes, Jeffery, Taylor, & Ewing, 2013; Russell-Smith, Maybery, & Bayliss, 2010; Sutherland & Crewther, 2010). To date, however, relatively little research has examined whether similar overlap exists in the guidance of emotional attention. To our knowledge, the only autistic-trait based study to provide some suggestive results in this regard was conducted by Miu, Pană and Avram (2012), who found that a fearful (compared to neutral) facial expression

High AQ adults do not see angry faces in a RSVP

3

slowed the redirection of attention through gaze cueing in low-trait individuals, but had no differential effect in high-trait individuals. This finding parallels dissociations found in the processing of eye gaze cues in ASD and control groups (Uono, Sato, & Toichi, 2009). However, it is not known whether differences in the ability of emotional faces to capture attention found in ASD relative to non-ASD groups (Ashwin et al., 2006) are also generalizable to high-trait groups. It is also not known whether emotions other than fear, such as anger, guide attention in a similar manner across ASD and high-trait groups.

One way to examine how emotions guide attention is via a paradigm known as the attentional blink (AB). The AB refers to the reduced detection of the second of two targets presented within approximately 500ms of each other in a rapid serial visual presentation (RSVP; Broadbent & Broadbent, 1987; Raymond, Shapiro, & Arnell, 1992). The AB is typically attributed, either directly or indirectly, to the requirement to process the initial target (T1), which limits the availability of attentional resources for the second target (T2; Chun & Potter, 1995; Di Lollo, Kawahara, Ghorashi, & Enns, 2005; Jolicoeur & Dell'Acqua, 1998; Olivers & Meeter, 2008; Taatgen, Juvina, Schipper, Borst, & Martens, 2009; Visser, Merikle, & Di Lollo, 2005; Visser, 2007; Wyble, Bowman, & Nieuwenstein, 2009). Notably, the AB is attenuated (higher T2 accuracy) when T2 is an emotionally arousing stimulus (e.g. the word "RAPE") relative to a non-arousing, neutral stimulus (e.g. the word "TABLE"; Anderson, 2005; Keil & Ihssen, 2004). This has led to the suggestion that arousing stimuli have enhanced resistance to interference from competing, neutral stimuli, increasing the likelihood that the information will be consciously available (Anderson, 2005).

To date, evidence suggests that a similar AB occurs across both ASD and neurotypical individuals when neutral target stimuli are presented in the AB (Amirault et al., 2009; Rinehart, Tonge, Brereton, & Bradshaw, 2010). From this we can infer that the general attentional mechanisms responsible for processing rapid sensory inputs are comparable between the two groups. However, differences in AB performance arise when making comparisons between

High AQ adults do not see angry faces in a RSVP

4

emotionally arousing and neutral target stimuli. In neurotypical children and adults, when T2 is emotionally arousing, the AB is attenuated relative to when T2 is non-arousing (Anderson, 2005; de Oca, Villa, Cervantes, & Welbourne, 2012; Jong et al., 2009; Keil, Ihssen, & Heim, 2006; Maratos, Mogg, & Bradley, 2008; Schwabe et al., 2011; Yerys et al., 2013). In contrast, results in the ASD literature are mixed. Adults with ASD show no change in performance when T2 is an emotionally-arousing word stimulus (Corden, Chilvers, & Skuse, 2008b; Gaigg & Bowler, 2009). Conversely, children with ASD show an attenuated AB when T2 is an emotionally-arousing picture stimulus (Yerys et al., 2013).

In sum then, our study has two primary motivations. First, we wanted to investigate significant questions about the overlap between ASD and high autistic-like traits with respect to the emotional guidance of attention. To do this, we examined task performance on an attentional paradigm that has been examined in ASD individuals, but not in individuals with high levels of autistic-like traits – the attentional blink. Second, we wished to investigate whether emotional face stimuli would produce similar results to those obtained using emotional words with ASD adults (Corden et al., 2008b; Gaigg & Bowler, 2009). This would provide some suggestive evidence clarifying whether differences between earlier studies with ASD adults and children were attributable to participant age or the type of emotional stimuli (words vs. pictures). To address these questions, we recruited neurotypical adults who displayed low (Low AQ) or high (High AQ) levels of autistic-like traits and tested them using an AB paradigm in which an angry or neutral face was presented as T2 (see Yerys et al., 2013). We reasoned that if autistic-like behaviours are on a spectrum that extends to the neurotypical population then, when viewing angry faces in a RSVP, resulting attenuation of the AB should be greater in Low AQ participants relative to High AQ participants, mirroring the dichotomy between neurotypical and ASD adults reported in prior research (Corden et al., 2008b; Gaigg & Bowler, 2009).

High AQ adults do not see angry faces in a RSVP

5

Methods

Participants

Three-hundred-and-sixty-eight second year psychology students (122 male) at the University of Western Australia participated in the study to satisfy part of a course requirement.

Materials

Autism Spectrum Quotient (AQ)

The AQ, developed by Baron-Cohen et al. (2001), is a 50 item self-report questionnaire designed to measure autistic-like traits and behaviours exhibited in a neurotypical sample. The questionnaire uses a four-item forced-choice format and scoring followed the 1-4 method described by Austin (2005). Following the logic of Austin (2005), we did this to take advantage of the breadth of potentially useful information in each item and to increase variability in total AQ score.

Attentional Blink Task

Stimuli were presented and responses collected using Presentation® software (Version 16.3, Neurobehavioral Systems) running on a PC using Windows 7, connected to a 22" HP L2245wg monitor. Facial stimuli were identical to those used by Yerys et al. (2013), comprising colour headshot photographs, 4.05° high and 3.16° wide, selected from the NimStim set (Tottenham et al., 2009). Three types of stimuli were used in the task: a) two photographs of dogs' heads to be used as T1, b) 28 photographs of faces with the interior details (e.g. eyes, nose) scrambled and outline preserved to be used as distractors in the RSVP, and c) 14 photographs of faces with neutral and angry expressions (seven of each) to be used as T2. Of these 14, there were seven males and seven females, and seven were depicted with their mouths open, and seven with their mouths closed. The scrambled faces were generated by Yerys et al. (2013) using a method devised by Conway et al. (2008).

The task consisted of 216 trials divided into three equal blocks, separated by self-paced breaks. Each trial began with the presentation of a fixation cross for 750ms, followed by a RSVP of 15 images (see Figure 1). Successive images in the stream were presented at 100ms intervals (16ms stimulus; 84ms blank ISI). Nine different types of trials were presented during the AB task. Three of the types were catch trials with zero or one target, which included: a) 30 T1-only trials, where only the dog's head was present in the RSVP, b) 30 T2-only trials, where only the unscrambled face was present in the RSVP, and c) 30 trials where neither T1 or T2 were present in the RSVP. The purpose of these trials was to estimate levels of response bias. The remaining six trial types were the test trials that measured the AB and therefore contained both T1 and T2. The six types were formed by crossing three levels of T1-T2 lag with a neutral or angry T2 face.

[FIGURE 1 PLACEHOLDER]

The first target, if present, was positioned equally often as the fourth, fifth or sixth stimulus in the RSVP. The second target, if present, was positioned as either the second, fourth or eighth stimulus in the RSVP following T1 presentation (lag-2, lag-4 and lag-8 respectively; for T2-only catch trials, T2 was placed in a similar position as if T1 were present). Equal numbers of trials were presented at each lag, evenly divided between neutral and angry facial expressions (for test trials, 42 trials per lag; 21 per expression). Presentation order was random with the constraint that identical trial types did not occur more than twice in a row. Each block contained the same number of each trial type. Participants completed ten practice trials before commencing the test trials.

Procedure

Participants first completed a paper version of the AQ, with scores used to select Low- and High AQ groups for comparison. The AB task immediately followed, with participants seated approximately 500mm in front of the computer monitor. Participants were instructed to watch closely on each trial for the presentation of a dog (T1) and unscrambled face (T2) in the RSVP stream. Following the RSVP, participants were first prompted to respond if they had observed a dog in the

High AQ adults do not see angry faces in a RSVP

7

RSVP, and then to indicate whether they had observed an unscrambled face. Participants responded to both queries using a keyboard, pressing 'Y' to indicate that they had observed a dog/unscrambled face, or 'N' to indicate they had not. Participants were allowed 5000ms to respond to each of the questions, with no response being recorded as an error. The fixation cross marking the start of the next trial was presented immediately following the second response.

Results

Comparison groups were created by selecting participants in the top and bottom quintiles of the distribution of AQ scores. The Low AQ group consisted of students with an AQ score ≤ 95 ($n = 80$), and the high AQ group consisted of students with an AQ score ≥ 116 ($n = 75$). From these initial groups, nine participants (five Low AQ, four High AQ) scoring $\leq 50\%$ accuracy (i.e. chance level) on catch trials were omitted from subsequent analyses. The effect size for the separation in means for the Low and High AQ groups was $r = 0.94^1$. Descriptive statistics for the resulting groups are presented in Table 1.

[TABLE 1 PLACEHOLDER]

Catch trial accuracy was similar across groups (for a summary, see Table 2), with independent samples t-tests revealing no significant differences (all t 's ≤ 0.51 , all p 's ≥ 0.61). For trials containing both targets, to determine if T1 accuracy varied significantly between conditions, a three-way mixed factorial analysis of variance (ANOVA) was conducted on the means shown in Table 2, with AQ group as a between-groups factor (two levels: Low AQ and High AQ), Lag as a within-groups factor (three levels: 2, 4 and 8) and Emotion as a within-groups factor (two levels: neutral and

¹ In Ruzich et al.'s (2015) recent review of studies that have used the AQ, based on the original 0-1 scoring method, they report a mean AQ of 16.94 for non-clinical participants. In order to provide a useful comparison to this norm, we re-scored our participants using this method and found our Low AQ group to have a mean score of 8.93 (SD = 2.54) and the High AQ group to have a mean score of 24.54 (SD = 4.13). Single-sample t -tests revealed that both our groups had means significantly different to 16.94 (both t 's > 15.51 , both p 's $< .001$).

High AQ adults do not see angry faces in a RSVP

8

angry). No main effects or interactions were significant (all F 's ≤ 2.18 , all p 's ≥ 0.12 ; see Table 2 for descriptive statistics).

[TABLE 2 PLACEHOLDER]

Second-target accuracy was calculated using only trials for which T1 was correctly identified.

This is because the source of T2 errors is unknown if a T1 error is also made (Raymond et al., 1992).

Mean T2 accuracy separated by AQ group, lag and emotion type are presented in Figure 2. These T2 accuracy scores were submitted to a three-way mixed factorial ANOVA with the same factors used in the previous analysis of T1 accuracy. This analysis yielded a significant main effect of Lag, $F(2, 144) = 139.58, p < .001, \eta_p^2 = .49$, consistent with the AB, a marginal Lag x Emotion interaction, $F(2, 288) = 2.30, p = .10, \eta_p^2 = .02$, and an AQ Group x Lag x Emotion interaction, $F(2, 288) = 3.43, p = .03, \eta_p^2 = .02$. No other effects were significant (all p 's $> .31$, all $\eta_p^2 < .01$). We followed up the significant three-way interaction with post-hoc paired samples t -tests (see Table 3).

[TABLE 3 PLACEHOLDER]

As the paired samples t -tests revealed a discrepancy between AQ groups at Lag-2, follow-up independent samples t -tests were used to reveal that, though the two groups were comparable at detecting T2 in Lag-2 neutral trials, $t(144) = .36, p = .72, r = .03$, the Low AQ group was significantly better at reporting T2 in Lag-2 angry trials, $t(144) = 1.70, p = .05, r = .14$, one-tailed. Thus there was a facilitative effect of the angry stimuli at Lag-2 for Low AQ participants, but not for High AQ participants.

[FIGURE 2 PLACEHOLDER]

Discussion

This study examined whether deficits in guidance of attention by emotional stimuli, similar to those reported in ASD samples, were present within a high autistic-like trait neurotypical group. By comparing the performance of two groups of low- and high-trait individuals on an AB task that

High AQ adults do not see angry faces in a RSVP

9

used neutral and threatening faces as targets, we found a pattern of results comparable to those reported using similar paradigms with clinical ASD and neurotypical comparison groups. Specifically, our results mirror those reported by Corden (2008b) and Gaigg and Bowler (2009), who showed little-to-no enhancement for ASD adults in identifying emotionally arousing targets presented within a RSVP. Our findings on visual attention using the AB paradigm complement those reported by Miu et al. (2012) using a gaze-cueing paradigm: high levels of autistic-like traits were associated with reduced interference from emotional faces in Miu et al. (2012) and with reduced ability to capture attention in the present study. In both cases, this implies that the impact of emotional faces on attentional systems is altered in those with high levels of autistic-like traits.

Our findings significantly expand upon previous work by showing that the capture of attention by emotional stimuli is similar for ASD (Ashwin et al., 2006) and high-trait individuals. This outcome may be interpreted in at least two ways. One option is that the failure to obtain emotional guidance of attention in both ASD and high-trait individuals indicates that this deficit is not a unique or defining characteristic of ASD (see Gregory and Plaisted-Grant, 2013). On this option, instances of reduced emotional guidance in ASD reported in other studies (Ashwin et al., 2006; Corden et al., 2008b; Gaigg & Bowler, 2009) may not be indicative of an ASD-linked 'deficit', but rather a behavioural pattern that is also present in many neurotypical individuals. Another option is that our findings provide further support for a dimensional model of ASD, in which ASD and ASD-associated traits exist on a continuum that extends into the neurotypical population (e.g., see Constantino and Todd, 2003). This interpretation suggests that much may be learned about the mechanisms underlying ASD by studying performance in participants with high levels of autistic-like traits who are considerably greater in number and thus easier to recruit than individuals with an ASD diagnosis.

Importantly, our results also suggest that parallels between ASD and high-trait individuals in the guidance of attention by emotion can be seen in angry as well as fearful faces (Miu et al., 2012). This is significant because there are plausible reasons to predict that angry faces might be more

High AQ adults do not see angry faces in a RSVP

10

effective at guiding attention in high-trait/ASD participants. In particular, although both anger and fear are associated with threat, angry faces convey a more 'direct' threat, thus capturing attention more efficiently. For example, Palermo and Coltheart (2004) showed that the angry faces from the Nimstim database (Tottenham et al., 2009) are identified faster and more accurately than the fearful faces drawn from the same database [these faces are used here and in Miu et al. (2012)].

To the extent that there are parallels between High AQ and ASD individuals, our results can also comment on the source of earlier differences found in studies of emotional guidance in ASD reported by Corden et al. (2008b), Gaigg and Bowler (2009) and Yerys et al. (2013). These studies reported no attenuation of the AB for emotional words in adults with ASD (Corden et al., 2008b; Gaigg & Bowler, 2009), but intact attenuation of the AB in children with emotional pictures (Yerys et al., 2013). Assuming that our findings can be taken as a preliminary indication of the expected pattern of results for an adult ASD and control comparison study, ours and the Yerys et al. (2013) results together suggest there is a developmental change with aging in ASD that impacts the efficient guidance of attention arising from emotional information in faces. This follows from the fact that whereas we reported an absence of beneficial effects on attentional guidance by threatening faces in our High AQ adult group, Yerys et al. (2013) found no differences in attentional guidance between typically developing and autistic children using the same paradigm. Confirmation of this conjecture in a study replicating our methodology with an adult ASD sample would further bolster the suggestion for attentional-related developmental changes.

Given our findings show broad similarities in the way that emotional faces guide attention in ASD and in neurotypical individuals with autistic-like traits, an explanation for our results may usefully be made by looking at how faces are processed in ASD. Notably, the efficient processing of faces is believed to depend largely on the successful development of a holistic or configural processing style (De Sonneville et al., 2002). Whereas neurotypical individuals tend to develop a more configural processing style for face processing (Tanaka & Farah, 1993), ASD individuals often

High AQ adults do not see angry faces in a RSVP

11

show a preference for using more piecemeal processing styles across both facial and non-facial stimuli (Behrmann, Thomas, & Humphreys, 2006; Behrmann, Avidan, et al., 2006; Isomura, Ogawa, Yamada, Shibasaki, & Masataka, 2014a, see Happé & Frith (2006) for a review).

This is of particular relevance to the current study, as differences in configural versus piecemeal processing styles have been reported on numerous occasions when comparing neurotypical individuals with low and high AQ scores (Almeida et al., 2010a, 2010b, 2013; Grinter, Van Beek, et al., 2009; Grinter, Maybery, et al., 2009; Russell-Smith et al., 2010; Sutherland & Crewther, 2010). Plausibly, then, underlying differences in processing strategies potentially expressed early in childhood (Kaldy, Krapar, Carter, & Blaser, 2011) might lead to differences in the way that emotional faces serve to guide attention in Low and High AQ groups. Analogous to the distinction between holistic and piecemeal face processing styles, studies have found that adults with ASD rely on rule-based processing strategies when processing emotional stimuli to a greater degree than do neurotypical individuals, who employ prototype or template-based matching strategies (Rutherford & McIntosh, 2007; Walsh, Vida, & Rutherford, 2014). Further, using a rule-based processing style in adults is likely to be more time consuming (Celani et al., 1999). Thus, in situations such as the AB task, where stimuli must be evaluated quickly, we hypothesise that individuals with high levels of autistic-like traits process emotional information too slowly for it to effectively guide attention.

What neurological processes may underlie the differences in emotion-based attentional guidance in our Low and High AQ groups? The answer to this question may also be gleaned from existing literature on individuals with ASD and individuals with high levels of autistic-like traits. Amygdala dysfunction has been implicated as a key factor underlying relatively poorer processing of emotional stimuli in ASD (see Schultz, 2005, for a review), with the thought that there is a failure to develop the ability to efficiently assign significance to socially meaningful or arousing stimuli (Grelotti, Gauthier, & Schultz, 2002; Schultz, Romanski, & Tsatsanis, 2000; Schultz, 2005;

High AQ adults do not see angry faces in a RSVP

12

Waterhouse, Fein, & Modahl, 1996). A number of neuroimaging studies have supported this view, showing relatively reduced activation of the amygdala in ASD individuals when making non-emotional judgements of emotional faces (Baron-Cohen et al., 1999; Corbett et al., 2009; Critchley et al., 2000; Kleinhans et al., 2011; Kliemann et al., 2012; Nickl-Jockschat et al., 2015; Pelphrey, Morris, McCarthy, & LaBar, 2007; Pierce, Haist, Sedaghat, & Courchesne, 2004). As a result of this dysfunction, the failure to effectively code emotional information in faces may prevent them from efficiently guiding attention. Importantly, there is also evidence for amygdala abnormalities in neurotypical individuals with high levels of autistic-like traits. Iidaka, Miyakoshi, Harada and Nakai (2012) found an increased volume of white matter connecting the amygdala and the superior temporal gyrus for individuals high in autistic-like traits compared to those with low levels of these traits.

One potential objection to our conclusions might be made on the basis of the fact that we failed to find a main effect of emotion on T2 performance, as in earlier studies with ASD participants (Corden et al., 2008b; Gaigg & Bowler, 2009; Yerys et al., 2013). However, we suggest that this is in keeping with the relatively small AB magnitude observed in our study ($\approx 20\%$) compared to these earlier experiments ($\approx 45-60\%$). This left less room for emotional guidance to improve performance, and also likely reduced the size of our three-way interaction. With this in mind, it is possible that High AQ participants might have shown attentional guidance effects if overall accuracy was lower. This could be examined in future studies by increasing the difficulty of the T1 task in order to boost the magnitude of the AB (Visser, 2007). However, even if the AB were to be modulated in the High AQ group under these conditions, it is clear from the present findings that the effect of angry faces would still be reduced for High AQ participants relative to Low AQ participants.

In conclusion, we found that the typical modulation of attention by emotional face stimuli is not intact in high autistic-like trait adults in the AB paradigm. This finding is further evidence that ASD-like patterns of behaviour are seen in individuals in the neurotypical population who report high

High AQ adults do not see angry faces in a RSVP

13

levels of autistic-like traits. Notably, to our knowledge, no study has directly investigated emotional modulation of attention in an AB paradigm in conjunction with neuroimaging techniques in ASD or High AQ samples. Such data would aid greatly in clarifying underlying neural mechanisms.

Furthermore, the AB studies conducted so far have been limited to the use of negatively valenced stimuli (Corden et al., 2008b; Gaigg & Bowler, 2009). It would be useful to determine whether positive emotions lead to similar differences in patterns of attentional orienting between ASD/High AQ and control/Low AQ individuals.

Acknowledgements

This research was supported by Australia Research Council Discovery Project grants to TAWV (DP120102313) and MTM (DP120104713). Funding was also provided to MCWE by The University of

Western Australia. The authors have no conflicts of interest to declare.

References

- Adolphs, R., Sears, L., & Piven, J. (2001). Abnormal Processing of Social Information from Faces in Autism. *Journal of Cognitive Neuroscience*, *13*(2), 232–240.
<http://doi.org/10.1162/089892901564289>
- Algom, D., Chajut, E., & Lev, S. (2004). A rational look at the emotional stroop phenomenon: A generic slowdown, not a stroop effect. *Journal of Experimental Psychology: General*, *133*(3), 323–38. <http://doi.org/10.1037/0096-3445.133.3.323>
- Almeida, R. A., Dickinson, J. E., Maybery, M. T., Badcock, J. C., & Badcock, D. R. (2010a). A new step towards understanding Embedded Figures Test performance in the autism spectrum: the radial frequency search task. *Neuropsychologia*, *48*(2), 374–81.
<http://doi.org/10.1016/j.neuropsychologia.2009.09.024>
- Almeida, R. A., Dickinson, J. E., Maybery, M. T., Badcock, J. C., & Badcock, D. R. (2010b). Visual search

High AQ adults do not see angry faces in a RSVP

14

performance in the autism spectrum II: the radial frequency search task with additional segmentation cues. *Neuropsychologia*, 48(14), 4117–24.

<http://doi.org/10.1016/j.neuropsychologia.2010.10.009>

Almeida, R. A., Dickinson, J. E., Maybery, M. T., Badcock, J. C., & Badcock, D. R. (2013). Visual Search Targeting Either Local or Global Perceptual Processes Differs as a Function of Autistic-Like Traits in the Typically Developing Population. *Journal of Autism and Developmental Disorders*, 43(6), 1272–1286. <http://doi.org/10.1007/s10803-012-1669-7>

Amirault, M., Etchegoyhen, K., Delord, S., Mendizabal, S., Kraushaar, C., Hesling, I., ... Mayo, W. (2009). Alteration of Attentional Blink in High Functioning Autism: A Pilot Study. *Journal of Autism and Developmental Disorders*, 39(11), 1522–1528. <http://doi.org/10.1007/s10803-009-0821-5>

Anderson, A. K. (2005). Affective influences on the attentional dynamics supporting awareness. *Journal of Experimental Psychology: General*, 134(2), 258–81. <http://doi.org/10.1037/0096-3445.134.2.258>

Ashwin, C., Wheelwright, S., & Baron-Cohen, S. (2006). Finding a face in the crowd: Testing the anger superiority effect in Asperger Syndrome. *Brain and Cognition*, 61(1), 78–95. <http://doi.org/10.1016/j.bandc.2005.12.008>

Austin, E. J. (2005). Personality correlates of the broader autism phenotype as assessed by the Autism Spectrum Quotient (AQ). *Personality and Individual Differences*, 38(2), 451–460. <http://doi.org/10.1016/j.paid.2004.04.022>

Baron-Cohen, S., Ring, H. A., Wheelwright, S., Bullmore, E. T., Brammer, M. J., Simmons, A., & Williams, S. C. R. (1999). Social intelligence in the normal and autistic brain: an fMRI study. *European Journal of Neuroscience*, 11(6), 1891–1898. <http://doi.org/10.1046/j.1460-9568.1999.00621.x>

High AQ adults do not see angry faces in a RSVP

15

Baron-Cohen, S., Wheelwright, S., & Jolliffe, T. (1997). Is There a “Language of the Eyes”? Evidence from Normal Adults, and Adults with Autism or Asperger Syndrome. *Visual Cognition*, 4(3), 311–331. <http://doi.org/10.1080/713756761>

Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning Autism, Males and Females, Scientists and Mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5–17. <http://doi.org/10.1023/A:1005653411471>

Behrmann, M., Avidan, G., Leonard, G. L., Kimchi, R., Luna, B., Humphreys, K., & Minshew, N. (2006). Configural processing in autism and its relationship to face processing. *Neuropsychologia*, 44(1), 110–129. <http://doi.org/10.1016/j.neuropsychologia.2005.04.002>

Behrmann, M., Thomas, C., & Humphreys, K. (2006). Seeing it differently: visual processing in autism. *Trends in Cognitive Sciences*, 10(6), 258–264. <http://doi.org/10.1016/j.tics.2006.05.001>

Bölte, S., & Poustka, F. (2003). The recognition of facial affect in autistic and schizophrenic subjects and their first-degree relatives. *Psychological Medicine*, 33(5), 907–915. <http://doi.org/10.1017/S0033291703007438>

Broadbent, D. E., & Broadbent, M. H. (1987). From detection to identification: response to multiple targets in rapid serial visual presentation. *Perception & Psychophysics*, 42(2), 105–13. <http://doi.org/10.3758/BF03210498>

Celani, G., Battacchi, M. W., & Arcidiacono, L. (1999). The understanding of the emotional meaning of facial expressions in people with autism. *Journal of Autism and Developmental Disorders*, 29(1), 57–66. <http://doi.org/10.1023/A:1025970600181>

Chun, M., & Potter, M. (1995). A two-stage model for multiple target detection in rapid serial visual presentation. *Journal of Experimental Psychology: Human Perception and Performance*, 21(1), 109–127. <http://doi.org/10.3758/BF03210498>

High AQ adults do not see angry faces in a RSVP

16

Constantino, J. N., & Todd, R. D. (2003). Autistic traits in the general population: A twin study.

Archives of General Psychiatry, 60(May), 524–530. <http://doi.org/10.1001/archpsyc.60.5.524>

Conway, C. A., Jones, B. C., DeBruine, L. M., Little, A. C., & Sahraie, A. (2008). Transient pupil constrictions to faces are sensitive to orientation and species. *Journal of Vision*, 8(3).

<http://doi.org/10.1167/8.3.17>

Corbett, B. A., Carmean, V., Ravizza, S., Wendelken, C., Henry, M. L., Carter, C., & Rivera, S. M. (2009). A functional and structural study of emotion and face processing in children with autism. *Psychiatry Research: Neuroimaging*, 173(3), 196–205.

<http://doi.org/10.1016/j.psychresns.2008.08.005>

Corden, B., Chilvers, R., & Skuse, D. (2008a). Avoidance of emotionally arousing stimuli predicts social-perceptual impairment in Asperger's syndrome. *Neuropsychologia*, 46(1), 137–47.

<http://doi.org/10.1016/j.neuropsychologia.2007.08.005>

Corden, B., Chilvers, R., & Skuse, D. (2008b). Emotional Modulation of Perception in Asperger's Syndrome. *Journal of Autism and Developmental Disorders*, 38(6), 1072–1080.

<http://doi.org/10.1007/s10803-007-0485-y>

Costafreda, S. G., Brammer, M. J., David, A. S., & Fu, C. H. Y. (2008). Predictors of amygdala activation during the processing of emotional stimuli: A meta-analysis of 385 PET and fMRI studies. *Brain Research Reviews*, 58(1), 57–70.

<http://doi.org/10.1016/j.brainresrev.2007.10.012>

Critchley, H. D., Daly, E. M., Bullmore, E. T., Williams, S. C. R., Van Amelsvoort, T., Robertson, D. M., ... Murphy, D. G. M. (2000). The functional neuroanatomy of social behaviour: Changes in cerebral blood flow when people with autistic disorder process facial expressions. *Brain*, 123(11), 2203–2212. <http://doi.org/10.1093/brain/123.11.2203>

Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., ...

High AQ adults do not see angry faces in a RSVP

17

Davidson, R. J. (2005). Gaze fixation and the neural circuitry of face processing in autism.

Nature Neuroscience, 8(4), 519–26. <http://doi.org/10.1038/nn1421>

de Oca, B. M., Villa, M., Cervantes, M., & Welbourne, T. (2012). Emotional modulation of the attentional blink by pleasant and unpleasant pictures. *The Journal of General Psychology*, 139(4), 289–314. <http://doi.org/10.1080/00221309.2012.708681>

De Sonneville, L. M. J., Verschoor, C. A., Njokiktjien, C., Op het Veld, V., Toorenaar, N., & Vranken, M. (2002). Facial identity and facial emotions: speed, accuracy, and processing strategies in children and adults. *Journal of Clinical and Experimental Neuropsychology*, 24(2), 200–13. <http://doi.org/10.1076/jcen.24.2.200.989>

Di Lollo, V., Kawahara, J., Ghorashi, S. M. S., & Enns, J. T. (2005). The attentional blink: resource depletion or temporary loss of control? *Psychological Research*, 69(3), 191–200. <http://doi.org/10.1007/s00426-004-0173-x>

Eastwood, J. D., Smilek, D., & Merikle, P. M. (2001). Differential attentional guidance by unattended faces expressing positive and negative emotion. *Perception & Psychophysics*, 63(6), 1004–1013. <http://doi.org/10.3758/BF03194519>

Farran, E. K., Branson, A., & King, B. J. (2011). Visual search for basic emotional expressions in autism; impaired processing of anger, fear and sadness, but a typical happy face advantage. *Research in Autism Spectrum Disorders*, 5(1), 455–462. <http://doi.org/10.1016/j.rasd.2010.06.009>

Fox, E., Lester, V., Russo, R., Bowles, R. J., Pichler, A., & Dutton, K. (2000). Facial Expressions of Emotion: Are Angry Faces Detected More Efficiently? *Cognition & Emotion*, 14(1), 61–92. <http://doi.org/10.1080/026999300378996>

Frischen, A., Eastwood, J. D., & Smilek, D. (2008). Visual search for faces with emotional expressions. *Psychological Bulletin*, 134(5), 662–676. <http://doi.org/10.1037/0033-2909.134.5.662>

High AQ adults do not see angry faces in a RSVP

18

Fusar-Poli, P., Placentino, A., Carletti, F., Landi, P., Allen, P., Surguladze, S., ... Politi, P. (2009).

Functional atlas of emotional faces processing: a voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *Journal of Psychiatry & Neuroscience*, 34(6), 418–32.

Gaigg, S. B., & Bowler, D. M. (2009). Brief report: Attenuated emotional suppression of the attentional blink in Autism Spectrum Disorder: another non-social abnormality? *Journal of Autism and Developmental Disorders*, 39(8), 1211–7. [http://doi.org/10.1007/s10803-009-0719-](http://doi.org/10.1007/s10803-009-0719-2)

2

Gregory, B. L., & Plaisted-Grant, K. C. (2013). The Autism-Spectrum Quotient and Visual Search: Shallow and Deep Autistic Endophenotypes. *Journal of Autism and Developmental Disorders*. <http://doi.org/10.1007/s10803-013-1951-3>

Grelotti, D. J., Gauthier, I., & Schultz, R. T. (2002). Social interest and the development of cortical face specialization: What autism teaches us about face processing. *Developmental Psychobiology*, 40, 213–225. <http://doi.org/10.1002/dev.10028>

Grinter, E. J., Maybery, M. T., Van Beek, P. L., Pellicano, E., Badcock, J. C., & Badcock, D. R. (2009). Global visual processing and self-rated autistic-like traits. *Journal of Autism and Developmental Disorders*, 39(9), 1278–90. <http://doi.org/10.1007/s10803-009-0740-5>

Grinter, E. J., Van Beek, P. L., Maybery, M. T., & Badcock, D. R. (2009). Brief report: visuospatial analysis and self-rated autistic-like traits. *Journal of Autism and Developmental Disorders*, 39(4), 670–7. <http://doi.org/10.1007/s10803-008-0658-3>

Grossman, J. B., Klin, A., Carter, A. S., & Volkmar, F. (2000). Verbal bias in recognition of facial emotions in children with Asperger syndrome. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 41(3), 369–379. <http://doi.org/10.1111/1469-7610.00621>

Hansen, C. H., & Hansen, R. D. (1988). Finding the face in the crowd: an anger superiority effect. *Journal of Personality and Social Psychology*, 54(6), 917–924. <http://doi.org/10.1037/0022->

High AQ adults do not see angry faces in a RSVP

19

3514.54.6.917

Happé, F. G. E., & Frith, U. (2006). The Weak Coherence Account: Detail-focused Cognitive Style in Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, *36*(1), 5–25.

<http://doi.org/10.1007/s10803-005-0039-0>

Harms, M. B., Martin, A., & Wallace, G. L. (2010). Facial emotion recognition in autism spectrum disorders: a review of behavioral and neuroimaging studies. *Neuropsychology Review*, *20*(3), 290–322. <http://doi.org/10.1007/s11065-010-9138-6>

Hershler, O., & Hochstein, S. (2005). At first sight: A high-level pop out effect for faces. *Vision Research*, *45*(13), 1707–1724. <http://doi.org/10.1016/j.visres.2004.12.021>

Howard, M. a, Cowell, P. E., Boucher, J., Broks, P., Mayes, a, Farrant, a, & Roberts, N. (2000). Convergent neuroanatomical and behavioural evidence of an amygdala hypothesis of autism. *Neuroreport*, *11*(13), 2931–2935. <http://doi.org/10.1097/00001756-200009110-00020>

Humphreys, K., Minshew, N., Leonard, G. L., & Behrmann, M. (2007). A fine-grained analysis of facial expression processing in high-functioning adults with autism. *Neuropsychologia*, *45*(4), 685–695. <http://doi.org/10.1016/j.neuropsychologia.2006.08.003>

Iidaka, T., Miyakoshi, M., Harada, T., & Nakai, T. (2012). White matter connectivity between superior temporal sulcus and amygdala is associated with autistic trait in healthy humans. *Neuroscience Letters*, *510*(2), 154–158. <http://doi.org/10.1016/j.neulet.2012.01.029>

Isomura, T., Ogawa, S., Yamada, S., Shibasaki, M., & Masataka, N. (2014). Preliminary evidence that different mechanisms underlie the anger superiority effect in children with and without Autism Spectrum Disorders. *Frontiers in Psychology*, *5*(May), 461.

<http://doi.org/10.3389/fpsyg.2014.00461>

Jolicoeur, P., & Dell'Acqua, R. (1998). The demonstration of short-term consolidation. *Cognitive Psychology*, *36*(2), 138–202. <http://doi.org/10.1006/cogp.1998.0684>

High AQ adults do not see angry faces in a RSVP

20

Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of approaching adults predicts level of social disability in 2-year-old toddlers with autism spectrum disorder.

Archives of General Psychiatry, 65(8), 946–54. <http://doi.org/10.1001/archpsyc.65.8.946>

Jong, P. J. De, Koster, E. H. W., Van, R., De Jong, P. J., Van Wees, R., & Martens, S. (2009). Emotional facial expressions and the attentional blink: Attenuated blink for angry and happy faces irrespective of social anxiety. *Cognition & Emotion*, 23(8), 1640–1652.

<http://doi.org/10.1080/02699930802490227>

Kaldy, Z., Kraper, C., Carter, A. S., & Blaser, E. (2011). Toddlers with Autism Spectrum Disorder are more successful at visual search than typically developing toddlers. *Developmental Science*, 14(5), 980–8. <http://doi.org/10.1111/j.1467-7687.2011.01053.x>

Keil, A., & Ihssen, N. (2004). Identification Facilitation for Emotionally Arousing Verbs During the Attentional Blink. *Emotion*, 4(1), 23–35. <http://doi.org/10.1037/1528-3542.4.1.23>

Keil, A., Ihssen, N., & Heim, S. (2006). Early cortical facilitation for emotionally arousing targets during the attentional blink. *BMC Biology*, 4, 23. <http://doi.org/10.1186/1741-7007-4-23>

Kirchner, J. C., Hatri, A., Heekeren, H. R., & Dziobek, I. (2011). Autistic symptomatology, face processing abilities, and eye fixation patterns. *Journal of Autism and Developmental Disorders*, 41(2), 158–67. <http://doi.org/10.1007/s10803-010-1032-9>

Kleinhans, N. M., Richards, T., Johnson, L. C., Weaver, K. E., Greenson, J., Dawson, G., & Aylward, E. (2011). fMRI evidence of neural abnormalities in the subcortical face processing system in ASD. *NeuroImage*, 54(1), 697–704. <http://doi.org/10.1016/j.neuroimage.2010.07.037>

Kliemann, D., Dziobek, I., Hatri, A., Baudewig, J., & Heekeren, H. R. (2012). The Role of the Amygdala in Atypical Gaze on Emotional Faces in Autism Spectrum Disorders. *Journal of Neuroscience*, 32(28), 9469–9476. <http://doi.org/10.1523/JNEUROSCI.5294-11.2012>

Klin, A., Jones, W., Schultz, R., Volkmar, F., & Cohen, D. (2002). Visual fixation patterns during

High AQ adults do not see angry faces in a RSVP

21

viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, 59(9), 809–16.

<http://doi.org/10.1001/archpsyc.59.9.809>

Krysko, K. M., & Rutherford, M. D. (2009). A threat-detection advantage in those with autism spectrum disorders. *Brain and Cognition*, 69(3), 472–80.

<http://doi.org/10.1016/j.bandc.2008.10.002>

Maratos, F. A., Mogg, K., & Bradley, B. P. (2008). Identification of angry faces in the attentional blink. *Cognition & Emotion*, 22(7), 1340–1352. <http://doi.org/10.1080/02699930701774218>

Mather, M., & Nesmith, K. (2008). Arousal-Enhanced Location Memory for Pictures. *Journal of Memory and Language*, 58(2), 449–464. <http://doi.org/10.1016/j.jml.2007.01.004>

Mathews, A., & MacLeod, C. (1985). Selective processing of threat cues in anxiety states. *Behaviour Research and Therapy*, 23(5), 563–569. [http://doi.org/10.1016/0005-7967\(85\)90104-4](http://doi.org/10.1016/0005-7967(85)90104-4)

Miu, A. C., Pană, S. E., & Avram, J. (2012). Emotional face processing in neurotypicals with autistic traits: Implications for the broad autism phenotype. *Psychiatry Research*, 198(3), 489–494.

<http://doi.org/10.1016/j.psychres.2012.01.024>

Nickl-Jockschat, T., Rottschy, C., Thommes, J., Schneider, F., Laird, A. R., Fox, P. T., & Eickhoff, S. B. (2015). Neural networks related to dysfunctional face processing in autism spectrum disorder. *Brain Structure and Function*, 220(4), 2355–2371. <http://doi.org/10.1007/s00429-014-0791-z>

Öhman, A., & Dimberg, U. (1984). An evolutionary perspective on human behavior. In W. M. Waid (Ed.), *Sociophysiology* (pp. 47–86). New York: Springer-Verlag.

Öhman, A., Lundqvist, D., & Esteves, F. (2001). The face in the crowd revisited: A threat advantage with schematic stimuli. *Journal of Personality and Social Psychology*, 80(3), 381–396.

<http://doi.org/10.1037/0022-3514.80.3.381>

Olivers, C. N. L., & Meeter, M. (2008). A boost and bounce theory of temporal attention.

High AQ adults do not see angry faces in a RSVP

22

Psychological Review, 115(4), 836–63. <http://doi.org/10.1037/a0013395>

Palermo, R., & Coltheart, M. (2004). Photographs of facial expression: accuracy, response times, and ratings of intensity. *Behavior Research Methods, Instruments, & Computers : A Journal of the Psychonomic Society, Inc*, 36(4), 634–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15641409>

Pelphrey, K. A., Morris, J. P., McCarthy, G., & LaBar, K. S. (2007). Perception of dynamic changes in facial affect and identity in autism. *Social Cognitive and Affective Neuroscience*, 2(2), 140–149. <http://doi.org/10.1093/scan/nsm010>

Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual Scanning of Faces in Autism. *Journal of Autism and Developmental Disorders*, 32(4), 249–261. <http://doi.org/10.1023/A:1016374617369>

Phan, K. L., Wager, T., Taylor, S. F., & Liberzon, I. (2002). Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage*, 16(2), 331–348. <http://doi.org/10.1006/nimg.2002.1087>

Pierce, K., Haist, F., Sedaghat, F., & Courchesne, E. (2004). The brain response to personally familiar faces in autism: findings of fusiform activity and beyond. *Brain*, 127(12), 2703–2716. <http://doi.org/10.1093/brain/awh289>

Raymond, J. E., Shapiro, K. L., & Arnell, K. M. (1992). Temporary suppression of visual processing in an RSVP task: An attentional blink? *Journal of Experimental Psychology: Human Perception and Performance*, 18(3), 849–860. <http://doi.org/10.1037/0096-1523.18.3.849>

Rhodes, G., Jeffery, L., Taylor, L., & Ewing, L. (2013). Autistic traits are linked to reduced adaptive coding of face identity and selectively poorer face recognition in men but not women. *Neuropsychologia*, 51(13), 2702–2708. <http://doi.org/10.1016/j.neuropsychologia.2013.08.016>

Rinehart, N., Tonge, B., Brereton, A., & Bradshaw, J. (2010). Attentional blink in young people with

High AQ adults do not see angry faces in a RSVP

23

high-functioning autism and Asperger's disorder. *Autism*, 14(1), 47–66.

<http://doi.org/10.1177/1362361309335718>

Russell-Smith, S. N., Maybery, M. T., & Bayliss, D. M. (2010). Are the autism and positive schizotypy spectra diametrically opposed in local versus global processing? *Journal of Autism and Developmental Disorders*, 40(8), 968–77. <http://doi.org/10.1007/s10803-010-0945-7>

Rutherford, M. D., & McIntosh, D. N. (2007). Rules versus prototype matching: strategies of perception of emotional facial expressions in the autism spectrum. *Journal of Autism and Developmental Disorders*, 37(2), 187–96. <http://doi.org/10.1007/s10803-006-0151-9>

Ruzich, E., Allison, C., Smith, P., Watson, P., Auyeung, B., Ring, H., & Baron-cohen, S. (2015). Measuring autistic traits in the general population : a systematic review of the Autism-Spectrum Quotient (AQ) in a nonclinical population sample of 6,900 typical adult males and females. *Molecular Autism*, 6(1), 1–12. <http://doi.org/10.1186/2040-2392-6-2>

Schimmack, U. (2005). Attentional interference effects of emotional pictures: threat, negativity, or arousal? *Emotion*, 5(1), 55–66. <http://doi.org/10.1037/1528-3542.5.1.55>

Schultz, R. T. (2005). Developmental deficits in social perception in autism: the role of the amygdala and fusiform face area. *International Journal of Developmental Neuroscience*, 23(2-3), 125–41. <http://doi.org/10.1016/j.ijdevneu.2004.12.012>

Schultz, R. T., Romanski, L. M., & Tsatsanis, K. D. (2000). Neurofunctional models of autistic disorder and Asperger syndrome: Clues from neuroimaging. (A. Klin, F. R. Volkmar, & S. S. Sparrow, Eds.) *Asperger Syndrome*. New York, NY, US: Guilford Press.

Schwabe, L., Merz, C. J., Walter, B., Vaitl, D., Wolf, O. T., & Stark, R. (2011). Emotional modulation of the attentional blink: the neural structures involved in capturing and holding attention. *Neuropsychologia*, 49(3), 416–25. <http://doi.org/10.1016/j.neuropsychologia.2010.12.037>

Spezio, M. L., Adolphs, R., Hurley, R. S. E., & Piven, J. (2007). Abnormal use of facial information in

High AQ adults do not see angry faces in a RSVP

24

high-functioning autism. *Journal of Autism and Developmental Disorders*, 37(5), 929–39.

<http://doi.org/10.1007/s10803-006-0232-9>

Sutherland, A., & Crewther, D. P. (2010). Magnocellular visual evoked potential delay with high autism spectrum quotient yields a neural mechanism for altered perception. *Brain*, 133(7), 2089–97. <http://doi.org/10.1093/brain/awq122>

Taatgen, N. A., Juvina, I., Schipper, M., Borst, J. P., & Martens, S. (2009). Too much control can hurt: a threaded cognition model of the attentional blink. *Cognitive Psychology*, 59(1), 1–29.

<http://doi.org/10.1016/j.cogpsych.2008.12.002>

Tanaka, J. W., & Farah, M. J. (1993). Parts and wholes in face recognition. *The Quarterly Journal of Experimental Psychology Section A*, 46(2), 225–245.

<http://doi.org/10.1080/14640749308401045>

Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., ... Nelson, C. (2009). The NimStim set of facial expressions: judgments from untrained research participants.

Psychiatry Research, 168(3), 242–9. <http://doi.org/10.1016/j.psychres.2008.05.006>

Uljarevic, M., & Hamilton, A. (2013). Recognition of emotions in autism: a formal meta-analysis.

Journal of Autism and Developmental Disorders, 43(7), 1517–26.

<http://doi.org/10.1007/s10803-012-1695-5>

Uono, S., Sato, W., & Toichi, M. (2009). Dynamic fearful gaze does not enhance attention orienting in individuals with Asperger's disorder. *Brain and Cognition*, 71(3), 229–233.

<http://doi.org/10.1016/j.bandc.2009.08.015>

Visser, T. A. W. (2007). Masking T1 difficulty: processing time and the attentional blink. *Journal of Experimental Psychology. Human Perception and Performance*, 33(2), 285–97.

<http://doi.org/10.1037/0096-1523.33.2.285>

Visser, T. A. W., Merikle, P., & Di Lollo, V. (2005). Priming in the attentional blink: Perception without

High AQ adults do not see angry faces in a RSVP

25

awareness? *Visual Cognition*. <http://doi.org/10.1080/13506280444000733>

Vogt, J., De Houwer, J., Koster, E. H. W., Van Damme, S., & Crombez, G. (2008). Allocation of spatial attention to emotional stimuli depends upon arousal and not valence. *Emotion*, 8(6), 880–5.

<http://doi.org/10.1037/a0013981>

Walsh, J. A., Vida, M. D., & Rutherford, M. D. (2014). Strategies for perceiving facial expressions in adults with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 44(5),

1018–26. <http://doi.org/10.1007/s10803-013-1953-1>

Waterhouse, L., Fein, D., & Modahl, C. (1996). Neurofunctional mechanisms in autism. *Psychological Review*, 103(3), 457–489. <http://doi.org/10.1037/0033-295X.103.3.457>

Wyble, B., Bowman, H., & Nieuwenstein, M. (2009). The attentional blink provides episodic

distinctiveness: sparing at a cost. *Journal of Experimental Psychology: Human Perception and Performance*, 35(3), 787–807. <http://doi.org/10.1037/a0013902>

Yerys, B. E., Ruiz, E., Strang, J., Sokoloff, J., Kenworthy, L., & Vaidya, C. J. (2013). Modulation of attentional blink with emotional faces in typical development and in autism spectrum

disorders. *Journal of Child Psychology and Psychiatry*, 54(6), 636–643.

<http://doi.org/10.1111/jcpp.12013>

RUNNING HEADER: High AQ adults do not see angry faces in a RSVP

Threatening faces fail to guide attention for adults with autistic-like traits.

Michael C.W. English¹, Murray T. Maybery¹, Troy A.W. Visser¹

¹School of Psychology, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

Correspondence concerning this article should be addressed to:

Michael C.W. English.

School of Psychology, The University of Western Australia, Crawley, WA 6009, Australia.

Email: michael.english@uwa.edu.au. Telephone: +61 8 6488 1421.

Number of text pages: 10 pages (22 pages inc. references)

Number of Tables: 3

Number of Figures: 2

Grant sponsor: Australian Research Council; Grant Number(s): DP120102313, DP120104713

RUNNING HEADER: High AQ adults do not see angry faces in a RSVP

Threatening faces fail to guide attention for adults with autistic-like

traits.

Michael C.W. English, Murray T. Maybery, Troy A.W. Visser

Scientific Abstract

Individuals diagnosed with autistic spectrum conditions often show deficits in processing emotional faces relative to neurotypical peers. However, little is known about whether similar deficits exist in neurotypical individuals who show high-levels of autistic-like traits. To address this question, we compared performance on an attentional blink task in a large sample of adults who showed low- or high-levels of autistic-like traits on the Autism Spectrum Quotient. We found that threatening faces inserted as the second target in a rapid serial visual presentation were identified more accurately amongst individuals with low- compared to high-levels of autistic-like traits. This is the first study to show that attentional blink abnormalities seen in autism extend to the neurotypical population with autistic-like traits, adding to the growing body of research suggesting that autistic-related patterns of behaviours extend into a subset of the neurotypical population.

Lay Abstract

The majority of the neurotypical population detect threatening stimuli, such as angry faces, faster and more accurately than faces with neutral or positive emotional expressions. The ability for the threatening stimuli to more efficiently guide attention relative to less threatening stimuli is believed to reflect an evolutionary advantage, such that individuals can swiftly respond to threatening or dangerous situations. However, there is evidence to suggest that individuals diagnosed with autistic spectrum conditions fail to show this 'anger superiority effect' when the stimuli are presented for very brief periods of time (less than 20ms). In our study, we found that neurotypical adults with few

RUNNING HEADER: High AQ adults do not see angry faces in a RSVP

autistic-like traits (measured with the Autism Spectrum Quotient) were able to detect angry faces more often than neutral faces, whereas neurotypical individuals who reported higher levels of autistic-like traits did not detect angry faces any more often than neutral faces in the most difficult condition. This is further evidence in a growing field of research that suggests that some attentional abnormalities expressed by individuals with autism spectrum conditions may also be present in a subset of the neurotypical population.

Keywords

autistic-like traits; attentional blink; emotion; faces

Accepted Article

Introduction

Because resource limitations prevent us from simultaneously processing all of the stimuli available to sensory systems, attention mechanisms must be employed to select a subset of information based on endogenous and exogenous factors. One particularly important exogenous factor in guiding attention is the level of emotional arousal (both positive and negative) elicited by incoming information. An abundance of evidence suggests arousing stimuli are responded to more quickly and accurately than neutral stimuli (e.g., Algom, Chajut, & Lev, 2004; Anderson, 2005; Mather & Nesmith, 2008; Mathews & MacLeod, 1985; Schimmack, 2005; Vogt, De Houwer, Koster, Van Damme, & Crombez, 2008), indicating they are more rapidly attended. Moreover, compared to neutral images, emotionally-arousing visual images reliably increase activation in areas such as the amygdala, prefrontal cortex (PFC) and occipital regions (Costafreda, Brammer, David, & Fu, 2008; Fusar-Poli et al., 2009; Phan, Wager, Taylor, & Liberzon, 2002).

Faces are a common source of both positively- and negatively-valenced emotional arousal, and have been reliably shown to guide attention, most commonly illustrated by faster reaction times to identifying a target emotional face amongst neutral expression distractors (Frischen, Eastwood, & Smilek, 2008). For example, in the well-known “anger superiority effect” (Hansen & Hansen, 1988), participants reliably identify the presence of an individual with an angry face within an array of individuals with neutral or happy expressions faster than they identify the presence of a happy face within an array of angry or neutral faces (Eastwood, Smilek, & Merikle, 2001; Fox et al., 2000; Hansen & Hansen, 1988; Hershler & Hochstein, 2005; Öhman, Lundqvist, & Esteves, 2001). As angry or threatening faces are a critical signal of potential social conflicts, the ability to attend and respond to them swiftly is often theorised to benefit the observing individual in attempting to avoid the conflict (Öhman & Dimberg, 1984).

Individuals diagnosed with neurological disorders such as autism spectrum disorders (ASD) often exhibit relatively greater difficulty with facial and emotional recognition than their

High AQ adults do not see angry faces in a RSVP

2

neurotypical counterparts (Adolphs, Sears, & Piven, 2001; Bölte & Poustka, 2003; Celani, Battacchi, & Arcidiacono, 1999; Farran, Branson, & King, 2011; Grossman, Klin, Carter, & Volkmar, 2000; Howard et al., 2000; Humphreys, Minshew, Leonard, & Behrmann, 2007; Pelphrey et al., 2002; see Uljarevic & Hamilton, 2013, for a review), which may in part stem from a failure to attend to the more socially meaningful aspects of the face, such as the eyes (Baron-Cohen, Wheelwright, & Jolliffe, 1997; Corden, Chilvers, & Skuse, 2008a; Dalton et al., 2005; Jones, Carr, & Klin, 2008; Kirchner, Hatri, Heekeren, & Dziobek, 2011; Kliemann, Dziobek, Hatri, Baudewig, & Heekeren, 2012; Ami Klin, Jones, Schultz, Volkmar, & Cohen, 2002; Spezio, Adolphs, Hurley, & Piven, 2007). This is not to say that individuals with ASD are universally impaired when processing facial information. Krysko and Rutherford (2009), for example, demonstrated that ASD individuals show a relatively intact anger superiority effect, detecting angry faces in a visual search paradigm as quickly as their neurotypical peers (although with less accuracy). That said, however, even when performance is similar between ASD and neurotypical samples, there is evidence to suggest that compensatory mechanisms and atypical perceptual processing strategies are present in ASD participants (see Harms, Martin, & Wallace, 2010, for a review).

It is possible that the diminished ability to attend to emotional stimuli also manifests itself in parts of the neurotypical population. An increasing body of research suggests that behavioural characteristics generally associated with ASD are also present in neurotypical individuals who demonstrate above-average scores on measures of autistic-like traits, such as the Autism Spectrum Quotient (AQ) (Almeida, Dickinson, Maybery, Badcock, & Badcock, 2010a, 2010b, 2013; Baron-Cohen, Wheelwright, Skinner, Martin, & Clubley, 2001; Grinter, Van Beek, Maybery, & Badcock, 2009; Grinter, Maybery, et al., 2009; Rhodes, Jeffery, Taylor, & Ewing, 2013; Russell-Smith, Maybery, & Bayliss, 2010; Sutherland & Crewther, 2010). To date, however, relatively little research has examined whether similar overlap exists in the guidance of emotional attention. To our knowledge, the only autistic-trait based study to provide some suggestive results in this regard was conducted by Miu, Pană and Avram (2012), who found that a fearful (compared to neutral) facial expression

High AQ adults do not see angry faces in a RSVP

3

slowed the redirection of attention through gaze cueing in low-trait individuals, but had no differential effect in high-trait individuals. This finding parallels dissociations found in the processing of eye gaze cues in ASD and control groups (Uono, Sato, & Toichi, 2009). However, it is not known whether differences in the ability of emotional faces to capture attention found in ASD relative to non-ASD groups (Ashwin et al., 2006) are also generalizable to high-trait groups. It is also not known whether emotions other than fear, such as anger, guide attention in a similar manner across ASD and high-trait groups.

One way to examine how emotions guide attention is via a paradigm known as the attentional blink (AB). The AB refers to the reduced detection of the second of two targets presented within approximately 500ms of each other in a rapid serial visual presentation (RSVP; Broadbent & Broadbent, 1987; Raymond, Shapiro, & Arnell, 1992). The AB is typically attributed, either directly or indirectly, to the requirement to process the initial target (T1), which limits the availability of attentional resources for the second target (T2; Chun & Potter, 1995; Di Lollo, Kawahara, Ghorashi, & Enns, 2005; Jolicoeur & Dell'Acqua, 1998; Olivers & Meeter, 2008; Taatgen, Juvina, Schipper, Borst, & Martens, 2009; Visser, Merikle, & Di Lollo, 2005; Visser, 2007; Wyble, Bowman, & Nieuwenstein, 2009). Notably, the AB is attenuated (higher T2 accuracy) when T2 is an emotionally arousing stimulus (e.g. the word "RAPE") relative to a non-arousing, neutral stimulus (e.g. the word "TABLE"; Anderson, 2005; Keil & Ihssen, 2004). This has led to the suggestion that arousing stimuli have enhanced resistance to interference from competing, neutral stimuli, increasing the likelihood that the information will be consciously available (Anderson, 2005).

To date, evidence suggests that a similar AB occurs across both ASD and neurotypical individuals when neutral target stimuli are presented in the AB (Amirault et al., 2009; Rinehart, Tonge, Brereton, & Bradshaw, 2010). From this we can infer that the general attentional mechanisms responsible for processing rapid sensory inputs are comparable between the two groups. However, differences in AB performance arise when making comparisons between

High AQ adults do not see angry faces in a RSVP

4

emotionally arousing and neutral target stimuli. In neurotypical children and adults, when T2 is emotionally arousing, the AB is attenuated relative to when T2 is non-arousing (Anderson, 2005; de Oca, Villa, Cervantes, & Welbourne, 2012; Jong et al., 2009; Keil, Ihssen, & Heim, 2006; Maratos, Mogg, & Bradley, 2008; Schwabe et al., 2011; Yerys et al., 2013). In contrast, results in the ASD literature are mixed. Adults with ASD show no change in performance when T2 is an emotionally-arousing word stimulus (Corden, Chilvers, & Skuse, 2008b; Gaigg & Bowler, 2009). Conversely, children with ASD show an attenuated AB when T2 is an emotionally-arousing picture stimulus (Yerys et al., 2013).

In sum then, our study has two primary motivations. First, we wanted to investigate significant questions about the overlap between ASD and high autistic-like traits with respect to the emotional guidance of attention. To do this, we examined task performance on an attentional paradigm that has been examined in ASD individuals, but not in individuals with high levels of autistic-like traits – the attentional blink. Second, we wished to investigate whether emotional face stimuli would produce similar results to those obtained using emotional words with ASD adults (Corden et al., 2008b; Gaigg & Bowler, 2009). This would provide some suggestive evidence clarifying whether differences between earlier studies with ASD adults and children were attributable to participant age or the type of emotional stimuli (words vs. pictures). To address these questions, we recruited neurotypical adults who displayed low (Low AQ) or high (High AQ) levels of autistic-like traits and tested them using an AB paradigm in which an angry or neutral face was presented as T2 (see Yerys et al., 2013). We reasoned that if autistic-like behaviours are on a spectrum that extends to the neurotypical population then, when viewing angry faces in a RSVP, resulting attenuation of the AB should be greater in Low AQ participants relative to High AQ participants, mirroring the dichotomy between neurotypical and ASD adults reported in prior research (Corden et al., 2008b; Gaigg & Bowler, 2009).

High AQ adults do not see angry faces in a RSVP

5

Methods

Participants

Three-hundred-and-sixty-eight second year psychology students (122 male) at the University of Western Australia participated in the study to satisfy part of a course requirement.

Materials

Autism Spectrum Quotient (AQ)

The AQ, developed by Baron-Cohen et al. (2001), is a 50 item self-report questionnaire designed to measure autistic-like traits and behaviours exhibited in a neurotypical sample. The questionnaire uses a four-item forced-choice format and scoring followed the 1-4 method described by Austin (2005). Following the logic of Austin (2005), we did this to take advantage of the breadth of potentially useful information in each item and to increase variability in total AQ score.

Attentional Blink Task

Stimuli were presented and responses collected using Presentation[®] software (Version 16.3, Neurobehavioral Systems) running on a PC using Windows 7, connected to a 22" HP L2245wg monitor. Facial stimuli were identical to those used by Yerys et al. (2013), comprising colour headshot photographs, 4.05° high and 3.16° wide, selected from the NimStim set (Tottenham et al., 2009). Three types of stimuli were used in the task: a) two photographs of dogs' heads to be used as T1, b) 28 photographs of faces with the interior details (e.g. eyes, nose) scrambled and outline preserved to be used as distractors in the RSVP, and c) 14 photographs of faces with neutral and angry expressions (seven of each) to be used as T2. Of these 14, there were seven males and seven females, and seven were depicted with their mouths open, and seven with their mouths closed. The scrambled faces were generated by Yerys et al. (2013) using a method devised by Conway et al. (2008).

The task consisted of 216 trials divided into three equal blocks, separated by self-paced breaks. Each trial began with the presentation of a fixation cross for 750ms, followed by a RSVP of 15 images (see Figure 1). Successive images in the stream were presented at 100ms intervals (16ms stimulus; 84ms blank ISI). Nine different types of trials were presented during the AB task. Three of the types were catch trials with zero or one target, which included: a) 30 T1-only trials, where only the dog's head was present in the RSVP, b) 30 T2-only trials, where only the unscrambled face was present in the RSVP, and c) 30 trials where neither T1 or T2 were present in the RSVP. The purpose of these trials was to estimate levels of response bias. The remaining six trial types were the test trials that measured the AB and therefore contained both T1 and T2. The six types were formed by crossing three levels of T1-T2 lag with a neutral or angry T2 face.

[FIGURE 1 PLACEHOLDER]

The first target, if present, was positioned equally often as the fourth, fifth or sixth stimulus in the RSVP. The second target, if present, was positioned as either the second, fourth or eighth stimulus in the RSVP following T1 presentation (lag-2, lag-4 and lag-8 respectively; for T2-only catch trials, T2 was placed in a similar position as if T1 were present). Equal numbers of trials were presented at each lag, evenly divided between neutral and angry facial expressions (for test trials, 42 trials per lag; 21 per expression). Presentation order was random with the constraint that identical trial types did not occur more than twice in a row. Each block contained the same number of each trial type. Participants completed ten practice trials before commencing the test trials.

Procedure

Participants first completed a paper version of the AQ, with scores used to select Low- and High AQ groups for comparison. The AB task immediately followed, with participants seated approximately 500mm in front of the computer monitor. Participants were instructed to watch closely on each trial for the presentation of a dog (T1) and unscrambled face (T2) in the RSVP stream. Following the RSVP, participants were first prompted to respond if they had observed a dog in the

High AQ adults do not see angry faces in a RSVP

7

RSVP, and then to indicate whether they had observed an unscrambled face. Participants responded to both queries using a keyboard, pressing 'Y' to indicate that they had observed a dog/unscrambled face, or 'N' to indicate they had not. Participants were allowed 5000ms to respond to each of the questions, with no response being recorded as an error. The fixation cross marking the start of the next trial was presented immediately following the second response.

Results

Comparison groups were created by selecting participants in the top and bottom quintiles of the distribution of AQ scores. The Low AQ group consisted of students with an AQ score ≤ 95 ($n = 80$), and the high AQ group consisted of students with an AQ score ≥ 116 ($n = 75$). From these initial groups, nine participants (five Low AQ, four High AQ) scoring $\leq 50\%$ accuracy (i.e. chance level) on catch trials were omitted from subsequent analyses. The effect size for the separation in means for the Low and High AQ groups was $r = 0.94^1$. Descriptive statistics for the resulting groups are presented in Table 1.

[TABLE 1 PLACEHOLDER]

Catch trial accuracy was similar across groups (for a summary, see Table 2), with independent samples t-tests revealing no significant differences (all t 's ≤ 0.51 , all p 's ≥ 0.61). For trials containing both targets, to determine if T1 accuracy varied significantly between conditions, a three-way mixed factorial analysis of variance (ANOVA) was conducted on the means shown in Table 2, with AQ group as a between-groups factor (two levels: Low AQ and High AQ), Lag as a within-groups factor (three levels: 2, 4 and 8) and Emotion as a within-groups factor (two levels: neutral and

¹ In Ruzich et al.'s (2015) recent review of studies that have used the AQ, based on the original 0-1 scoring method, they report a mean AQ of 16.94 for non-clinical participants. In order to provide a useful comparison to this norm, we re-scored our participants using this method and found our Low AQ group to have a mean score of 8.93 (SD = 2.54) and the High AQ group to have a mean score of 24.54 (SD = 4.13). Single-sample t -tests revealed that both our groups had means significantly different to 16.94 (both t 's > 15.51 , both p 's $< .001$).

High AQ adults do not see angry faces in a RSVP

8

angry). No main effects or interactions were significant (all F 's ≤ 2.18 , all p 's ≥ 0.12 ; see Table 2 for descriptive statistics).

[TABLE 2 PLACEHOLDER]

Second-target accuracy was calculated using only trials for which T1 was correctly identified.

This is because the source of T2 errors is unknown if a T1 error is also made (Raymond et al., 1992).

Mean T2 accuracy separated by AQ group, lag and emotion type are presented in Figure 2. These T2 accuracy scores were submitted to a three-way mixed factorial ANOVA with the same factors used in the previous analysis of T1 accuracy. This analysis yielded a significant main effect of Lag, $F(2, 144) = 139.58, p < .001, \eta_p^2 = .49$, consistent with the AB, a marginal Lag x Emotion interaction, $F(2, 288) = 2.30, p = .10, \eta_p^2 = .02$, and an AQ Group x Lag x Emotion interaction, $F(2, 288) = 3.43, p = .03, \eta_p^2 = .02$. No other effects were significant (all p 's $> .31$, all $\eta_p^2 < .01$). We followed up the significant three-way interaction with post-hoc paired samples t -tests (see Table 3).

[TABLE 3 PLACEHOLDER]

As the paired samples t -tests revealed a discrepancy between AQ groups at Lag-2, follow-up independent samples t -tests were used to reveal that, though the two groups were comparable at detecting T2 in Lag-2 neutral trials, $t(144) = .36, p = .72, r = .03$, the Low AQ group was significantly better at reporting T2 in Lag-2 angry trials, $t(144) = 1.70, p = .05, r = .14$, one-tailed. Thus there was a facilitative effect of the angry stimuli at Lag-2 for Low AQ participants, but not for High AQ participants.

[FIGURE 2 PLACEHOLDER]

Discussion

This study examined whether deficits in guidance of attention by emotional stimuli, similar to those reported in ASD samples, were present within a high autistic-like trait neurotypical group. By comparing the performance of two groups of low- and high-trait individuals on an AB task that

High AQ adults do not see angry faces in a RSVP

9

used neutral and threatening faces as targets, we found a pattern of results comparable to those reported using similar paradigms with clinical ASD and neurotypical comparison groups. Specifically, our results mirror those reported by Corden (2008b) and Gaigg and Bowler (2009), who showed little-to-no enhancement for ASD adults in identifying emotionally arousing targets presented within a RSVP. Our findings on visual attention using the AB paradigm complement those reported by Miu et al. (2012) using a gaze-cueing paradigm: high levels of autistic-like traits were associated with reduced interference from emotional faces in Miu et al. (2012) and with reduced ability to capture attention in the present study. In both cases, this implies that the impact of emotional faces on attentional systems is altered in those with high levels of autistic-like traits.

Our findings significantly expand upon previous work by showing that the capture of attention by emotional stimuli is similar for ASD (Ashwin et al., 2006) and high-trait individuals. This outcome may be interpreted in at least two ways. One option is that the failure to obtain emotional guidance of attention in both ASD and high-trait individuals indicates that this deficit is not a unique or defining characteristic of ASD (see Gregory and Plaisted-Grant, 2013). On this option, instances of reduced emotional guidance in ASD reported in other studies (Ashwin et al., 2006; Corden et al., 2008b; Gaigg & Bowler, 2009) may not be indicative of an ASD-linked 'deficit', but rather a behavioural pattern that is also present in many neurotypical individuals. Another option is that our findings provide further support for a dimensional model of ASD, in which ASD and ASD-associated traits exist on a continuum that extends into the neurotypical population (e.g., see Constantino and Todd, 2003). This interpretation suggests that much may be learned about the mechanisms underlying ASD by studying performance in participants with high levels of autistic-like traits who are considerably greater in number and thus easier to recruit than individuals with an ASD diagnosis.

Importantly, our results also suggest that parallels between ASD and high-trait individuals in the guidance of attention by emotion can be seen in angry as well as fearful faces (Miu et al., 2012). This is significant because there are plausible reasons to predict that angry faces might be more

High AQ adults do not see angry faces in a RSVP

10

effective at guiding attention in high-trait/ASD participants. In particular, although both anger and fear are associated with threat, angry faces convey a more 'direct' threat, thus capturing attention more efficiently. For example, Palermo and Coltheart (2004) showed that the angry faces from the Nimstim database (Tottenham et al., 2009) are identified faster and more accurately than the fearful faces drawn from the same database [these faces are used here and in Miu et al. (2012)].

To the extent that there are parallels between High AQ and ASD individuals, our results can also inform comment on the source of earlier differences found in studies of emotional guidance in ASD reported by Corden et al. (2008b), Gaigg and Bowler (2009) and Yerys et al. (2013). These studies reported no attenuation of the AB for emotional words in adults with ASD (Corden et al., 2008b; Gaigg & Bowler, 2009), but intact attenuation of the AB in children with emotional pictures (Yerys et al., 2013). Assuming that our findings can be taken as a preliminary indication of the expected pattern of results for an adult ASD and control comparison study, ours and the Yerys et al. (2013) results together suggest there is a developmental change with aging in ASD that impacts the efficient guidance of attention arising from emotional information in faces. This follows from the fact that whereas we reported an absence of beneficial effects on attentional guidance by threatening faces in our High AQ adult group, Yerys et al. (2013) found no differences in attentional guidance between typically developing and autistic children using the same paradigm. Confirmation of this conjecture in a study replicating our methodology with an adult ASD sample would further bolster the suggestion for attentional-related developmental changes.

Given our findings show broad similarities in the way that emotional faces guide attention in ASD and in neurotypical individuals with autistic-like traits, an explanation for our results may usefully be made by looking at how faces are processed in ASD. Notably, the efficient processing of faces is believed to depend largely on the successful development of a holistic or configural processing style (De Sonneville et al., 2002). Whereas neurotypical individuals tend to develop a more configural processing style for face processing (Tanaka & Farah, 1993), ASD individuals often

High AQ adults do not see angry faces in a RSVP

11

show a preference for using more piecemeal processing styles across both facial and non-facial stimuli (Behrmann, Thomas, & Humphreys, 2006; Behrmann, Avidan, et al., 2006; Isomura, Ogawa, Yamada, Shibasaki, & Masataka, 2014a, see Happé & Frith (2006) for a review).

This is of particular relevance to the current study, as differences in configural versus piecemeal processing styles have been reported on numerous occasions when comparing neurotypical individuals with low and high AQ scores (Almeida et al., 2010a, 2010b, 2013; Grinter, Van Beek, et al., 2009; Grinter, Maybery, et al., 2009; Russell-Smith et al., 2010; Sutherland & Crewther, 2010). Plausibly, then, underlying differences in processing strategies potentially expressed early in childhood (Kaldy, Kraper, Carter, & Blaser, 2011) might lead to differences in the way that emotional faces serve to guide attention in Low and High AQ groups. Analogous to the distinction between holistic and piecemeal face processing styles, studies have found that adults with ASD rely on rule-based processing strategies when processing emotional stimuli to a greater degree than do neurotypical individuals, who employ prototype or template-based matching strategies (Rutherford & McIntosh, 2007; Walsh, Vida, & Rutherford, 2014). Further, using a rule-based processing style in adults is likely to be more time consuming (Celani et al., 1999). Thus, in situations such as the AB task, where stimuli must be evaluated quickly, we hypothesise that individuals with high levels of autistic-like traits process emotional information too slowly for it to effectively guide attention.

What neurological processes may underlie the differences in emotion-based attentional guidance in our Low and High AQ groups? The answer to this question may also be gleaned from existing literature on individuals with ASD and individuals with high levels of autistic-like traits. Amygdala dysfunction has been implicated as a key factor underlying relatively poorer processing of emotional stimuli in ASD (see Schultz, 2005, for a review), with the thought that there is a failure to develop the ability to efficiently assign significance to socially meaningful or arousing stimuli (Grelotti, Gauthier, & Schultz, 2002; Schultz, Romanski, & Tsatsanis, 2000; Schultz, 2005;

High AQ adults do not see angry faces in a RSVP

12

Waterhouse, Fein, & Modahl, 1996). A number of neuroimaging studies have supported this view, showing relatively reduced activation of the amygdala in ASD individuals when making non-emotional judgements of emotional faces (Baron-Cohen et al., 1999; Corbett et al., 2009; Critchley et al., 2000; Kleinhans et al., 2011; Kliemann et al., 2012; Nickl-Jockschat et al., 2015; Pelphrey, Morris, McCarthy, & LaBar, 2007; Pierce, Haist, Sedaghat, & Courchesne, 2004). As a result of this dysfunction, the failure to effectively code emotional information in faces may prevent them from efficiently guiding attention. Importantly, there is also evidence for amygdala abnormalities in neurotypical individuals with high levels of autistic-like traits. Iidaka, Miyakoshi, Harada and Nakai (2012) found an increased volume of white matter connecting the amygdala and the superior temporal gyrus for individuals high in autistic-like traits compared to those with low levels of these traits.

One potential objection to our conclusions might be made on the basis of the fact that we failed to find a main effect of emotion on T2 performance, as in earlier studies with ASD participants (Corden et al., 2008b; Gaigg & Bowler, 2009; Yerys et al., 2013). However, we suggest that this is in keeping with the relatively small AB magnitude observed in our study ($\approx 20\%$) compared to these earlier experiments ($\approx 45\text{-}60\%$). This left less room for emotional guidance to improve performance, and also likely reduced the size of our three-way interaction. With this in mind, it is possible that High AQ participants might have shown attentional guidance effects if overall accuracy was lower. This could be examined in future studies by increasing the difficulty of the T1 task in order to boost the magnitude of the AB (Visser, 2007). However, even if the AB were to be modulated in the High AQ group under these conditions, it is clear from the present findings that the effect of angry faces would still be reduced for High AQ participants relative to Low AQ participants.

In conclusion, we found that the typical modulation of attention by emotional face stimuli is not intact in high autistic-like trait adults in the AB paradigm. This finding is further evidence that ASD-like patterns of behaviour are seen in individuals in the neurotypical population who report high

High AQ adults do not see angry faces in a RSVP

13

levels of autistic-like traits. Notably, to our knowledge, no study has directly investigated emotional modulation of attention in an AB paradigm in conjunction with neuroimaging techniques in ASD or High AQ samples. Such data would aid greatly in clarifying underlying neural mechanisms.

Furthermore, the AB studies conducted so far have been limited to the use of negatively valenced stimuli (Corden et al., 2008b; Gaigg & Bowler, 2009). It would be useful to determine whether positive emotions lead to similar differences in patterns of attentional orienting between ASD/High AQ and control/Low AQ individuals.

Acknowledgements

This research was supported by Australia Research Council Discovery Project grants to TAWV (DP120102313) and MTM (DP120104713). Funding was also provided to MCWE by The University of

Western Australia. The authors have no conflicts of interest to declare.

References

- Adolphs, R., Sears, L., & Piven, J. (2001). Abnormal Processing of Social Information from Faces in Autism. *Journal of Cognitive Neuroscience*, *13*(2), 232–240.
<http://doi.org/10.1162/089892901564289>
- Algom, D., Chajut, E., & Lev, S. (2004). A rational look at the emotional stroop phenomenon: A generic slowdown, not a stroop effect. *Journal of Experimental Psychology: General*, *133*(3), 323–38. <http://doi.org/10.1037/0096-3445.133.3.323>
- Almeida, R. A., Dickinson, J. E., Maybery, M. T., Badcock, J. C., & Badcock, D. R. (2010a). A new step towards understanding Embedded Figures Test performance in the autism spectrum: the radial frequency search task. *Neuropsychologia*, *48*(2), 374–81.
<http://doi.org/10.1016/j.neuropsychologia.2009.09.024>
- Almeida, R. A., Dickinson, J. E., Maybery, M. T., Badcock, J. C., & Badcock, D. R. (2010b). Visual search

performance in the autism spectrum II: the radial frequency search task with additional segmentation cues. *Neuropsychologia*, 48(14), 4117–24.

<http://doi.org/10.1016/j.neuropsychologia.2010.10.009>

Almeida, R. A., Dickinson, J. E., Maybery, M. T., Badcock, J. C., & Badcock, D. R. (2013). Visual Search Targeting Either Local or Global Perceptual Processes Differs as a Function of Autistic-Like Traits in the Typically Developing Population. *Journal of Autism and Developmental Disorders*, 43(6), 1272–1286. <http://doi.org/10.1007/s10803-012-1669-7>

Amirault, M., Etchegoyhen, K., Delord, S., Mendizabal, S., Kraushaar, C., Hesling, I., ... Mayo, W. (2009). Alteration of Attentional Blink in High Functioning Autism: A Pilot Study. *Journal of Autism and Developmental Disorders*, 39(11), 1522–1528. <http://doi.org/10.1007/s10803-009-0821-5>

Anderson, A. K. (2005). Affective influences on the attentional dynamics supporting awareness. *Journal of Experimental Psychology: General*, 134(2), 258–81. <http://doi.org/10.1037/0096-3445.134.2.258>

Ashwin, C., Wheelwright, S., & Baron-Cohen, S. (2006). Finding a face in the crowd: Testing the anger superiority effect in Asperger Syndrome. *Brain and Cognition*, 61(1), 78–95. <http://doi.org/10.1016/j.bandc.2005.12.008>

Austin, E. J. (2005). Personality correlates of the broader autism phenotype as assessed by the Autism Spectrum Quotient (AQ). *Personality and Individual Differences*, 38(2), 451–460. <http://doi.org/10.1016/j.paid.2004.04.022>

Baron-Cohen, S., Ring, H. A., Wheelwright, S., Bullmore, E. T., Brammer, M. J., Simmons, A., & Williams, S. C. R. (1999). Social intelligence in the normal and autistic brain: an fMRI study. *European Journal of Neuroscience*, 11(6), 1891–1898. <http://doi.org/10.1046/j.1460-9568.1999.00621.x>

High AQ adults do not see angry faces in a RSVP

15

Baron-Cohen, S., Wheelwright, S., & Jolliffe, T. (1997). Is There a "Language of the Eyes"? Evidence from Normal Adults, and Adults with Autism or Asperger Syndrome. *Visual Cognition*, 4(3), 311–331. <http://doi.org/10.1080/713756761>

Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning Autism, Males and Females, Scientists and Mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5–17. <http://doi.org/10.1023/A:1005653411471>

Behrmann, M., Avidan, G., Leonard, G. L., Kimchi, R., Luna, B., Humphreys, K., & Minshew, N. (2006). Configural processing in autism and its relationship to face processing. *Neuropsychologia*, 44(1), 110–129. <http://doi.org/10.1016/j.neuropsychologia.2005.04.002>

Behrmann, M., Thomas, C., & Humphreys, K. (2006). Seeing it differently: visual processing in autism. *Trends in Cognitive Sciences*, 10(6), 258–264. <http://doi.org/10.1016/j.tics.2006.05.001>

Bölte, S., & Poustka, F. (2003). The recognition of facial affect in autistic and schizophrenic subjects and their first-degree relatives. *Psychological Medicine*, 33(5), 907–915. <http://doi.org/10.1017/S0033291703007438>

Broadbent, D. E., & Broadbent, M. H. (1987). From detection to identification: response to multiple targets in rapid serial visual presentation. *Perception & Psychophysics*, 42(2), 105–13. <http://doi.org/10.3758/BF03210498>

Celani, G., Battacchi, M. W., & Arcidiacono, L. (1999). The understanding of the emotional meaning of facial expressions in people with autism. *Journal of Autism and Developmental Disorders*, 29(1), 57–66. <http://doi.org/10.1023/A:1025970600181>

Chun, M., & Potter, M. (1995). A two-stage model for multiple target detection in rapid serial visual presentation. *Journal of Experimental Psychology: Human Perception and Performance*, 21(1), 109–127. <http://doi.org/10.3758/BF03210498>

High AQ adults do not see angry faces in a RSVP

16

Constantino, J. N., & Todd, R. D. (2003). Autistic traits in the general population: A twin study.

Archives of General Psychiatry, 60(May), 524–530. <http://doi.org/10.1001/archpsyc.60.5.524>

Conway, C. A., Jones, B. C., DeBruine, L. M., Little, A. C., & Sahraie, A. (2008). Transient pupil constrictions to faces are sensitive to orientation and species. *Journal of Vision*, 8(3).

<http://doi.org/10.1167/8.3.17>

Corbett, B. A., Carmean, V., Ravizza, S., Wendelken, C., Henry, M. L., Carter, C., & Rivera, S. M. (2009). A functional and structural study of emotion and face processing in children with autism. *Psychiatry Research: Neuroimaging*, 173(3), 196–205.

<http://doi.org/10.1016/j.psychresns.2008.08.005>

Corden, B., Chilvers, R., & Skuse, D. (2008a). Avoidance of emotionally arousing stimuli predicts social-perceptual impairment in Asperger's syndrome. *Neuropsychologia*, 46(1), 137–47.

<http://doi.org/10.1016/j.neuropsychologia.2007.08.005>

Corden, B., Chilvers, R., & Skuse, D. (2008b). Emotional Modulation of Perception in Asperger's Syndrome. *Journal of Autism and Developmental Disorders*, 38(6), 1072–1080.

<http://doi.org/10.1007/s10803-007-0485-y>

Costafreda, S. G., Brammer, M. J., David, A. S., & Fu, C. H. Y. (2008). Predictors of amygdala activation during the processing of emotional stimuli: A meta-analysis of 385 PET and fMRI studies. *Brain Research Reviews*, 58(1), 57–70.

<http://doi.org/10.1016/j.brainresrev.2007.10.012>

Critchley, H. D., Daly, E. M., Bullmore, E. T., Williams, S. C. R., Van Amelsvoort, T., Robertson, D. M., ... Murphy, D. G. M. (2000). The functional neuroanatomy of social behaviour: Changes in cerebral blood flow when people with autistic disorder process facial expressions. *Brain*, 123(11), 2203–2212. <http://doi.org/10.1093/brain/123.11.2203>

Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., ...

High AQ adults do not see angry faces in a RSVP

17

Davidson, R. J. (2005). Gaze fixation and the neural circuitry of face processing in autism.

Nature Neuroscience, 8(4), 519–26. <http://doi.org/10.1038/nn1421>

de Oca, B. M., Villa, M., Cervantes, M., & Welbourne, T. (2012). Emotional modulation of the attentional blink by pleasant and unpleasant pictures. *The Journal of General Psychology*, 139(4), 289–314. <http://doi.org/10.1080/00221309.2012.708681>

De Sonneville, L. M. J., Verschoor, C. A., Njokiktjien, C., Op het Veld, V., Toorenaar, N., & Vranken, M. (2002). Facial identity and facial emotions: speed, accuracy, and processing strategies in children and adults. *Journal of Clinical and Experimental Neuropsychology*, 24(2), 200–13. <http://doi.org/10.1076/jcen.24.2.200.989>

Di Lollo, V., Kawahara, J., Ghorashi, S. M. S., & Enns, J. T. (2005). The attentional blink: resource depletion or temporary loss of control? *Psychological Research*, 69(3), 191–200. <http://doi.org/10.1007/s00426-004-0173-x>

Eastwood, J. D., Smilek, D., & Merikle, P. M. (2001). Differential attentional guidance by unattended faces expressing positive and negative emotion. *Perception & Psychophysics*, 63(6), 1004–1013. <http://doi.org/10.3758/BF03194519>

Farran, E. K., Branson, A., & King, B. J. (2011). Visual search for basic emotional expressions in autism; impaired processing of anger, fear and sadness, but a typical happy face advantage. *Research in Autism Spectrum Disorders*, 5(1), 455–462. <http://doi.org/10.1016/j.rasd.2010.06.009>

Fox, E., Lester, V., Russo, R., Bowles, R. J., Pichler, A., & Dutton, K. (2000). Facial Expressions of Emotion: Are Angry Faces Detected More Efficiently? *Cognition & Emotion*, 14(1), 61–92. <http://doi.org/10.1080/026999300378996>

Frischen, A., Eastwood, J. D., & Smilek, D. (2008). Visual search for faces with emotional expressions. *Psychological Bulletin*, 134(5), 662–676. <http://doi.org/10.1037/0033-2909.134.5.662>

High AQ adults do not see angry faces in a RSVP

18

Fusar-Poli, P., Placentino, A., Carletti, F., Landi, P., Allen, P., Surguladze, S., ... Politi, P. (2009).

Functional atlas of emotional faces processing: a voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *Journal of Psychiatry & Neuroscience*, 34(6), 418–32.

Gaigg, S. B., & Bowler, D. M. (2009). Brief report: Attenuated emotional suppression of the attentional blink in Autism Spectrum Disorder: another non-social abnormality? *Journal of Autism and Developmental Disorders*, 39(8), 1211–7. <http://doi.org/10.1007/s10803-009-0719-2>

Gregory, B. L., & Plaisted-Grant, K. C. (2013). The Autism-Spectrum Quotient and Visual Search: Shallow and Deep Autistic Endophenotypes. *Journal of Autism and Developmental Disorders*. <http://doi.org/10.1007/s10803-013-1951-3>

Grelotti, D. J., Gauthier, I., & Schultz, R. T. (2002). Social interest and the development of cortical face specialization: What autism teaches us about face processing. *Developmental Psychobiology*, 40, 213–225. <http://doi.org/10.1002/dev.10028>

Grinter, E. J., Maybery, M. T., Van Beek, P. L., Pellicano, E., Badcock, J. C., & Badcock, D. R. (2009). Global visual processing and self-rated autistic-like traits. *Journal of Autism and Developmental Disorders*, 39(9), 1278–90. <http://doi.org/10.1007/s10803-009-0740-5>

Grinter, E. J., Van Beek, P. L., Maybery, M. T., & Badcock, D. R. (2009). Brief report: visuospatial analysis and self-rated autistic-like traits. *Journal of Autism and Developmental Disorders*, 39(4), 670–7. <http://doi.org/10.1007/s10803-008-0658-3>

Grossman, J. B., Klin, A., Carter, A. S., & Volkmar, F. (2000). Verbal bias in recognition of facial emotions in children with Asperger syndrome. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 41(3), 369–379. <http://doi.org/10.1111/1469-7610.00621>

Hansen, C. H., & Hansen, R. D. (1988). Finding the face in the crowd: an anger superiority effect. *Journal of Personality and Social Psychology*, 54(6), 917–924. <http://doi.org/10.1037/0022->

High AQ adults do not see angry faces in a RSVP

19

3514.54.6.917

Happé, F. G. E., & Frith, U. (2006). The Weak Coherence Account: Detail-focused Cognitive Style in Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 36(1), 5–25.

<http://doi.org/10.1007/s10803-005-0039-0>

Harms, M. B., Martin, A., & Wallace, G. L. (2010). Facial emotion recognition in autism spectrum disorders: a review of behavioral and neuroimaging studies. *Neuropsychology Review*, 20(3), 290–322. <http://doi.org/10.1007/s11065-010-9138-6>

Hershler, O., & Hochstein, S. (2005). At first sight: A high-level pop out effect for faces. *Vision Research*, 45(13), 1707–1724. <http://doi.org/10.1016/j.visres.2004.12.021>

Howard, M. a, Cowell, P. E., Boucher, J., Broks, P., Mayes, a, Farrant, a, & Roberts, N. (2000). Convergent neuroanatomical and behavioural evidence of an amygdala hypothesis of autism. *Neuroreport*, 11(13), 2931–2935. <http://doi.org/10.1097/00001756-200009110-00020>

Humphreys, K., Minshew, N., Leonard, G. L., & Behrmann, M. (2007). A fine-grained analysis of facial expression processing in high-functioning adults with autism. *Neuropsychologia*, 45(4), 685–695. <http://doi.org/10.1016/j.neuropsychologia.2006.08.003>

Iidaka, T., Miyakoshi, M., Harada, T., & Nakai, T. (2012). White matter connectivity between superior temporal sulcus and amygdala is associated with autistic trait in healthy humans. *Neuroscience Letters*, 510(2), 154–158. <http://doi.org/10.1016/j.neulet.2012.01.029>

Isomura, T., Ogawa, S., Yamada, S., Shibasaki, M., & Masataka, N. (2014). Preliminary evidence that different mechanisms underlie the anger superiority effect in children with and without Autism Spectrum Disorders. *Frontiers in Psychology*, 5(May), 461. <http://doi.org/10.3389/fpsyg.2014.00461>

Jolicoeur, P., & Dell'Acqua, R. (1998). The demonstration of short-term consolidation. *Cognitive Psychology*, 36(2), 138–202. <http://doi.org/10.1006/cogp.1998.0684>

High AQ adults do not see angry faces in a RSVP

20

Jones, W., Carr, K., & Klin, A. (2008). Absence of preferential looking to the eyes of approaching adults predicts level of social disability in 2-year-old toddlers with autism spectrum disorder.

Archives of General Psychiatry, 65(8), 946–54. <http://doi.org/10.1001/archpsyc.65.8.946>

Jong, P. J. De, Koster, E. H. W., Van, R., De Jong, P. J., Van Wees, R., & Martens, S. (2009). Emotional facial expressions and the attentional blink: Attenuated blink for angry and happy faces irrespective of social anxiety. *Cognition & Emotion*, 23(8), 1640–1652.

<http://doi.org/10.1080/02699930802490227>

Kaldy, Z., Kraper, C., Carter, A. S., & Blaser, E. (2011). Toddlers with Autism Spectrum Disorder are more successful at visual search than typically developing toddlers. *Developmental Science*, 14(5), 980–8. <http://doi.org/10.1111/j.1467-7687.2011.01053.x>

Keil, A., & Ihssen, N. (2004). Identification Facilitation for Emotionally Arousing Verbs During the Attentional Blink. *Emotion*, 4(1), 23–35. <http://doi.org/10.1037/1528-3542.4.1.23>

Keil, A., Ihssen, N., & Heim, S. (2006). Early cortical facilitation for emotionally arousing targets during the attentional blink. *BMC Biology*, 4, 23. <http://doi.org/10.1186/1741-7007-4-23>

Kirchner, J. C., Hatri, A., Heekeren, H. R., & Dziobek, I. (2011). Autistic symptomatology, face processing abilities, and eye fixation patterns. *Journal of Autism and Developmental Disorders*, 41(2), 158–67. <http://doi.org/10.1007/s10803-010-1032-9>

Kleinhans, N. M., Richards, T., Johnson, L. C., Weaver, K. E., Greenson, J., Dawson, G., & Aylward, E. (2011). fMRI evidence of neural abnormalities in the subcortical face processing system in ASD. *NeuroImage*, 54(1), 697–704. <http://doi.org/10.1016/j.neuroimage.2010.07.037>

Kliemann, D., Dziobek, I., Hatri, A., Baudewig, J., & Heekeren, H. R. (2012). The Role of the Amygdala in Atypical Gaze on Emotional Faces in Autism Spectrum Disorders. *Journal of Neuroscience*, 32(28), 9469–9476. <http://doi.org/10.1523/JNEUROSCI.5294-11.2012>

Klin, A., Jones, W., Schultz, R., Volkmar, F., & Cohen, D. (2002). Visual fixation patterns during

High AQ adults do not see angry faces in a RSVP

21

viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, 59(9), 809–16.

<http://doi.org/10.1001/archpsyc.59.9.809>

Krysko, K. M., & Rutherford, M. D. (2009). A threat-detection advantage in those with autism spectrum disorders. *Brain and Cognition*, 69(3), 472–80.

<http://doi.org/10.1016/j.bandc.2008.10.002>

Maratos, F. A., Mogg, K., & Bradley, B. P. (2008). Identification of angry faces in the attentional blink. *Cognition & Emotion*, 22(7), 1340–1352. <http://doi.org/10.1080/02699930701774218>

Mather, M., & Nesmith, K. (2008). Arousal-Enhanced Location Memory for Pictures. *Journal of Memory and Language*, 58(2), 449–464. <http://doi.org/10.1016/j.jml.2007.01.004>

Mathews, A., & MacLeod, C. (1985). Selective processing of threat cues in anxiety states. *Behaviour Research and Therapy*, 23(5), 563–569. [http://doi.org/10.1016/0005-7967\(85\)90104-4](http://doi.org/10.1016/0005-7967(85)90104-4)

Miu, A. C., Pană, S. E., & Avram, J. (2012). Emotional face processing in neurotypicals with autistic traits: Implications for the broad autism phenotype. *Psychiatry Research*, 198(3), 489–494.

<http://doi.org/10.1016/j.psychres.2012.01.024>

Nickl-Jockschat, T., Rottschy, C., Thommes, J., Schneider, F., Laird, A. R., Fox, P. T., & Eickhoff, S. B. (2015). Neural networks related to dysfunctional face processing in autism spectrum disorder. *Brain Structure and Function*, 220(4), 2355–2371. <http://doi.org/10.1007/s00429-014-0791-z>

Öhman, A., & Dimberg, U. (1984). An evolutionary perspective on human behavior. In W. M. Waid (Ed.), *Sociophysiology* (pp. 47–86). New York: Springer-Verlag.

Öhman, A., Lundqvist, D., & Esteves, F. (2001). The face in the crowd revisited: A threat advantage with schematic stimuli. *Journal of Personality and Social Psychology*, 80(3), 381–396.

<http://doi.org/10.1037/0022-3514.80.3.381>

Olivers, C. N. L., & Meeter, M. (2008). A boost and bounce theory of temporal attention.

High AQ adults do not see angry faces in a RSVP

22

Psychological Review, 115(4), 836–63. <http://doi.org/10.1037/a0013395>

Palermo, R., & Coltheart, M. (2004). Photographs of facial expression: accuracy, response times, and ratings of intensity. *Behavior Research Methods, Instruments, & Computers : A Journal of the Psychonomic Society, Inc*, 36(4), 634–8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15641409>

Pelphrey, K. A., Morris, J. P., McCarthy, G., & LaBar, K. S. (2007). Perception of dynamic changes in facial affect and identity in autism. *Social Cognitive and Affective Neuroscience*, 2(2), 140–149. <http://doi.org/10.1093/scan/nsm010>

Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual Scanning of Faces in Autism. *Journal of Autism and Developmental Disorders*, 32(4), 249–261. <http://doi.org/10.1023/A:1016374617369>

Phan, K. L., Wager, T., Taylor, S. F., & Liberzon, I. (2002). Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage*, 16(2), 331–348. <http://doi.org/10.1006/nimg.2002.1087>

Pierce, K., Haist, F., Sedaghat, F., & Courchesne, E. (2004). The brain response to personally familiar faces in autism: findings of fusiform activity and beyond. *Brain*, 127(12), 2703–2716. <http://doi.org/10.1093/brain/awh289>

Raymond, J. E., Shapiro, K. L., & Arnell, K. M. (1992). Temporary suppression of visual processing in an RSVP task: An attentional blink? *Journal of Experimental Psychology: Human Perception and Performance*, 18(3), 849–860. <http://doi.org/10.1037/0096-1523.18.3.849>

Rhodes, G., Jeffery, L., Taylor, L., & Ewing, L. (2013). Autistic traits are linked to reduced adaptive coding of face identity and selectively poorer face recognition in men but not women. *Neuropsychologia*, 51(13), 2702–2708. <http://doi.org/10.1016/j.neuropsychologia.2013.08.016>

Rinehart, N., Tonge, B., Brereton, A., & Bradshaw, J. (2010). Attentional blink in young people with

High AQ adults do not see angry faces in a RSVP

23

high-functioning autism and Asperger's disorder. *Autism*, 14(1), 47–66.

<http://doi.org/10.1177/1362361309335718>

Russell-Smith, S. N., Maybery, M. T., & Bayliss, D. M. (2010). Are the autism and positive schizotypy spectra diametrically opposed in local versus global processing? *Journal of Autism and Developmental Disorders*, 40(8), 968–77. <http://doi.org/10.1007/s10803-010-0945-7>

Rutherford, M. D., & McIntosh, D. N. (2007). Rules versus prototype matching: strategies of perception of emotional facial expressions in the autism spectrum. *Journal of Autism and Developmental Disorders*, 37(2), 187–96. <http://doi.org/10.1007/s10803-006-0151-9>

Ruzich, E., Allison, C., Smith, P., Watson, P., Auyeung, B., Ring, H., & Baron-cohen, S. (2015). Measuring autistic traits in the general population : a systematic review of the Autism-Spectrum Quotient (AQ) in a nonclinical population sample of 6,900 typical adult males and females. *Molecular Autism*, 6(1), 1–12. <http://doi.org/10.1186/2040-2392-6-2>

Schimmack, U. (2005). Attentional interference effects of emotional pictures: threat, negativity, or arousal? *Emotion*, 5(1), 55–66. <http://doi.org/10.1037/1528-3542.5.1.55>

Schultz, R. T. (2005). Developmental deficits in social perception in autism: the role of the amygdala and fusiform face area. *International Journal of Developmental Neuroscience*, 23(2-3), 125–41. <http://doi.org/10.1016/j.ijdevneu.2004.12.012>

Schultz, R. T., Romanski, L. M., & Tsatsanis, K. D. (2000). Neurofunctional models of autistic disorder and Asperger syndrome: Clues from neuroimaging. (A. Klin, F. R. Volkmar, & S. S. Sparrow, Eds.) *Asperger Syndrome*. New York, NY, US: Guilford Press.

Schwabe, L., Merz, C. J., Walter, B., Vaitl, D., Wolf, O. T., & Stark, R. (2011). Emotional modulation of the attentional blink: the neural structures involved in capturing and holding attention. *Neuropsychologia*, 49(3), 416–25. <http://doi.org/10.1016/j.neuropsychologia.2010.12.037>

Spezio, M. L., Adolphs, R., Hurley, R. S. E., & Piven, J. (2007). Abnormal use of facial information in

High AQ adults do not see angry faces in a RSVP

24

high-functioning autism. *Journal of Autism and Developmental Disorders*, 37(5), 929–39.

<http://doi.org/10.1007/s10803-006-0232-9>

Sutherland, A., & Crewther, D. P. (2010). Magnocellular visual evoked potential delay with high autism spectrum quotient yields a neural mechanism for altered perception. *Brain*, 133(7), 2089–97. <http://doi.org/10.1093/brain/awq122>

Taatgen, N. A., Juvina, I., Schipper, M., Borst, J. P., & Martens, S. (2009). Too much control can hurt: a threaded cognition model of the attentional blink. *Cognitive Psychology*, 59(1), 1–29.

<http://doi.org/10.1016/j.cogpsych.2008.12.002>

Tanaka, J. W., & Farah, M. J. (1993). Parts and wholes in face recognition. *The Quarterly Journal of Experimental Psychology Section A*, 46(2), 225–245.

<http://doi.org/10.1080/14640749308401045>

Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., ... Nelson, C. (2009). The NimStim set of facial expressions: judgments from untrained research participants.

Psychiatry Research, 168(3), 242–9. <http://doi.org/10.1016/j.psychres.2008.05.006>

Uljarevic, M., & Hamilton, A. (2013). Recognition of emotions in autism: a formal meta-analysis.

Journal of Autism and Developmental Disorders, 43(7), 1517–26.

<http://doi.org/10.1007/s10803-012-1695-5>

Uono, S., Sato, W., & Toichi, M. (2009). Dynamic fearful gaze does not enhance attention orienting in individuals with Asperger's disorder. *Brain and Cognition*, 71(3), 229–233.

<http://doi.org/10.1016/j.bandc.2009.08.015>

Visser, T. A. W. (2007). Masking T1 difficulty: processing time and the attentional blink. *Journal of Experimental Psychology. Human Perception and Performance*, 33(2), 285–97.

<http://doi.org/10.1037/0096-1523.33.2.285>

Visser, T. A. W., Merikle, P., & Di Lollo, V. (2005). Priming in the attentional blink: Perception without

High AQ adults do not see angry faces in a RSVP

25

awareness? *Visual Cognition*. <http://doi.org/10.1080/13506280444000733>

Vogt, J., De Houwer, J., Koster, E. H. W., Van Damme, S., & Crombez, G. (2008). Allocation of spatial attention to emotional stimuli depends upon arousal and not valence. *Emotion*, *8*(6), 880–5.

<http://doi.org/10.1037/a0013981>

Walsh, J. A., Vida, M. D., & Rutherford, M. D. (2014). Strategies for perceiving facial expressions in adults with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *44*(5),

1018–26. <http://doi.org/10.1007/s10803-013-1953-1>

Waterhouse, L., Fein, D., & Modahl, C. (1996). Neurofunctional mechanisms in autism. *Psychological Review*, *103*(3), 457–489. <http://doi.org/10.1037/0033-295X.103.3.457>

Wyble, B., Bowman, H., & Nieuwenstein, M. (2009). The attentional blink provides episodic

distinctiveness: sparing at a cost. *Journal of Experimental Psychology: Human Perception and Performance*, *35*(3), 787–807. <http://doi.org/10.1037/a0013902>

Yerys, B. E., Ruiz, E., Strang, J., Sokoloff, J., Kenworthy, L., & Vaidya, C. J. (2013). Modulation of attentional blink with emotional faces in typical development and in autism spectrum

disorders. *Journal of Child Psychology and Psychiatry*, *54*(6), 636–643.

<http://doi.org/10.1111/jcpp.12013>

Table 1.

Characteristics of the Low-AQ and High-AQ comparison groups (standard deviation in parentheses).

	Low AQ (n = 75, 17 male)	High AQ (n = 71, 27 male)
Mean Age (years)	21.35 (6.82)	20.77 (3.90)
Mean AQ	87.72 (6.36)	124.59 (7.47)

Accepted Article

Table 2.

Mean catch trial and Target 1 accuracy (proportion correct; standard deviation in parentheses).

		<i>Low AQ</i>	<i>High AQ</i>
<i>Mean Catch Trial Accuracy</i>			
No Target		91.29 (10.40)	91.13 (9.98)
T1-only		94.22 (5.96)	94.41 (6.64)
T2-only		86.00 (10.46)	86.90 (10.71)
<i>Mean Target 1 Accuracy for T1-T2 Trials</i>			
Lag-2	Neutral	94.79 (6.12)	94.97 (5.51)
	Angry	95.68 (5.65)	94.77 (6.14)
Lag-4	Neutral	95.05 (7.28)	94.63 (7.76)
	Angry	95.68 (5.81)	95.44 (5.72)
Lag-8	Neutral	96.25 (5.09)	96.04 (6.78)
	Angry	95.81 (6.91)	95.17 (6.90)

Table 3.

Results for paired samples *t*-tests comparing emotion at each lag for each AQ group (*p* values in parentheses).

	<i>Lag-2</i>	<i>Lag-4</i>	<i>Lag-8</i>
Low AQ	2.80 (<.01)**	1.26 (.21)	0.27 (.79)
High AQ	0.01 (1.00)	0.22 (.82)	0.98 (.33)

** Significant at the Bonferroni corrected 0.008 level

Accepted Article




Figure 1. Example of a RSVP with both T1 and T2 stimuli.
371x233mm (96 x 96 DPI)

Accepte




Figure 2. Mean percentage of T2 responses correct given T1 correct, as a function of AQ Group (Low and High), T2 Emotion (neutral and angry) and Lag (2, 4 and 8). Error bars represent standard error of the mean.

450x287mm (96 x 96 DPI)

Accept