

**The efficacy of Botulinum Toxin A on improving ease
of care in the upper and lower limbs: a systematic
review and meta-analysis using the GRADE
approach**

Authors: Jennifer A Baker jennyjas1@yahoo.co.uk

Gavin Pereira gpereira@ichr.uwa.edu.au

**Correspondence to: JA
Baker jennyjas1@yahoo.co.uk**

South Hams Physio Ltd

15 Allens Road

The Watermark

Ivybridge

Devon

PL21 0PW

Tel: +441752 891930

ABSTRACT

Objectives: A systematic review and meta analysis using the Grades of Recommendation, Assessment, Development and Evaluation (GRADE) approach; evaluating Botulinum Toxin type A efficacy on improving ease of care in the upper/lower limb.

Data Sources: Pubmed, Cinahl, Amed, Embase and Cochrane databases. English Language. Search to July 2014.

Review Methods: All randomized, placebo controlled trials on adults with difficulty in caring for the upper/lower limb resulting from spasticity of any origin and treated with a single dose of Botulinum Toxin A. Evidence quality was assessed by GRADE.

Results: Thirty-two studies were reviewed. Meta analysis was carried out on eleven upper limb and three lower limb studies. Evidence quality for the upper limb was moderate. A significant result for Botulinum Toxin A was found at four to twelve weeks for the upper limb (SMD 0.80 CI 0.55, 1.06, $p < 0.0001$) The effects were maintained for up to six months (SMD 0.48 CI 0.34, 0.62, $p < 0.0001$).

Evidence quality was very low for the lower limb. Meta analysis was only possible for global assessment of benefit. No significant effect was found. (Patient: RR 1.37 CI (0.94, 2.00) $p = 0.11$, Clinician: RR= 1.06 (0.84, 1.34) $p = 0.60$.)

Conclusion: Botulinum Toxin A improves ease of care in the upper limb for up to six months. No conclusion can be drawn for the lower limb.

INTRODUCTION

This is the second paper of three, which examine the efficacy of Botulinum Toxin A using the Grades of Assessment, Development and Evaluation (GRADE) approach.¹ The first review focused on the topics of pain and spasticity and found very low quality evidence with no significant effects on pain for the upper and lower limbs, while evidence quality was moderate with significant effects in favour of Botulinum Toxin A for reducing spasticity².

This review looks at the effect of Botulinum Toxin A on improving ease of care. Tasks performed for the patient by a carer or by the patient's unaffected limb are often referred to as passive function or self care activities³. Problems with these tasks can arise from the complications of spasticity and soft tissue changes leading to; skin breakdown, mal odour, abnormal limb position and difficulty washing and dressing the limb⁴. This all has an adverse effect on caregiver burden⁴.

Botulinum Toxin blocks the release of acetylcholine from nerve endings and can selectively and reversibly induce muscle weakness and help reduce spasticity⁵. This temporary relief of muscle overactivity can allow easier extrinsic stretching of soft tissue structures and interrupt the muscle overactivity, which exacerbates tendon shortening⁶. This can in turn allow access to improved hygiene and dressing of the limb.

Several reviews and three Meta analyses have already reported that Botulinum Toxin A is effective in improving ease of care in stroke patients⁷⁻¹². The Meta analyses have generally reviewed the global assessment of benefit scales and looked at the upper limb primarily.⁷⁻⁹ Passive range of movement has been studied in a few reviews but with less significant outcomes.¹³⁻¹⁴ A recent review found a moderate treatment effect

using the disability assessment scale but this was combined with other measures that looked at more active or global constructs.¹⁵ No reviews have looked at goal attainment or lower limb studies separately.

This is the first systematic review of Botulinum Toxin A using ease of care outcomes and GRADE. The GRADE approach is becoming more widely recognised, is recommended for systematic reviews and guideline development, and is now used by Cochrane reviewers among other clinical groups¹⁶.

The following clinical questions will be addressed; Should Botulinum Toxin A be used to:

- improve ease of care in the upper limb?
- improve ease of care in the lower limb?

A subsequent paper will address active movement and quality of life.

METHODOLOGY

This study defined Botulinum Toxin as any standard preparation of the Type A toxin for clinical use to reduce spasticity.

Studies were included in the review if they were randomised controlled trials, included the use of Botulinum Toxin A versus a placebo/saline injection/control group, on either upper or lower limb in adult inpatients or outpatients, with outcome measures relating to ease of care. Muscle spasticity of any origin was considered; this allows for a comprehensive review and is more representative of the clinical setting.

Studies were excluded if they lacked a control group, were observational studies, on paediatrics or used other preparations of Botulinum Toxin besides Type A. Repeated injection studies were also excluded unless data from the initial injection could be used independently. All non English studies were excluded from this review.

The outcomes considered were passive range of movement, global assessment of benefit scales (also called clinical global impression, global assessment scale), disability assessment scale, carer burden scales and goals/goal attainment scale. These outcomes were selected from the recommendations in the Royal College of Physicians spasticity guidelines¹⁷ and from initial literature searches. Outcomes were rated from highest to lowest in terms of patient importance as per the GRADE protocol. Goals and patient rating scales were rated highest, followed by carer burden scales and passive range of movement was rated lowest: as improvement may or may not directly help the patient.

A literature search was carried out for all relevant studies from 1989 (date of approval of Botulinum Toxin for clinical use)⁸ up to 31st July 2014. The primary database searched was Pubmed using the MeSH terms of “muscle spasticity” and “Botulinum toxins”. The limits and exclusions set were: adults, humans, English language and randomised controlled trials. (See Appendix I) The search was repeated adding all individual outcome measure names but no new, relevant studies were found.

Searches with the same terms and limitations were carried out for Embase, Amed, Cinahl, and Cochrane.

Study quality was established using the GRADE approach as detailed in the previous paper². The following definition was used: “quality of evidence” relates to the extent which one can be confident that an estimate of effect is correct.¹ Initial summary of findings were carried out for qualitative syntheses and then where possible a meta analysis was carried out on subgroups with appropriate data. Data from the relevant studies was extracted by the author using a preprepared form. The data was reviewed and cross checked for anomalies. Decisions on quality were made using guidance from GRADE publications and the website.^{1,18-28}

Studies were chosen for Meta analysis if they provided sufficient data as a group in either dichotomous form (when data allowed results to be divided into improved versus no change/worse) or as means and standard deviations. Data was used from between one and six months post intervention; results were then analysed for four to twelve weeks (to analyse the effect over the active time for Botulinum Toxin) and for twelve to twenty four weeks (to gauge any significant lasting effects of treatment). Results given for week twelve could be used in either analysis but only once for each study, i.e. not duplicated.

Dichotomous data was analysed using the Mantel-Haenszel method to provide risk ratios (RR). Continuous data was analysed using the inverse-variance method to give a Weighted Mean Difference (WMD) for individual outcome measures where possible. The standardised mean difference (SMD) was used to pool the results of all outcome measures together as it allows for a variety of measurement methods. Random effects models were used in the presence of significant unexplained heterogeneity.

For dose ranging studies, treatment groups were combined to give a single pair wise comparison, to avoid potential bias in choosing results for analysis and as recommended by the Cochrane Handbook for systematic reviewers²⁹. When multiple joint results were presented the most commonly measured joint was used and any variations noted. Significance was set at $p < 0.05$.

Data was analysed using the statistical pack Revman 5.2 from the Cochrane Collaboration.

RESULTS

Qualitative analysis

In the upper limb, fourteen studies looked at passive range of movement³⁰⁻⁴³, four at the disability assessment scale⁴³⁻⁴⁶, nine at the global assessment of benefit^{32,37,38,44,46-50}, six at carer burden scales^{30,41,47,51-53} and six at goal setting^{32,37,39,41,47,51}. The majority of trials used stroke patients with a few including acquired brain injury or using mixed neurological diagnoses. Methodology and results presentation varied greatly across trials.

The Disability Assessment Scale was the most consistently applied tool. Goal setting in particular was often poorly explained, making it difficult to ascertain whether the nature of the goals was impairment, activity or participation based. Only two studies used the validated Goal Attainment Scale and one used the Canadian Occupational Performance Measure.

Significant improvement for treatment groups was found for all studies using the global assessment of benefit and the disability assessment scale, eight using passive range of movement and three who used Carer burden scales. Both trials using the Goal Attainment Scale found significant improvements with Botulinum Toxin. Other goal setting studies were less positive; two studies noted improvements but no difference was found between groups (these studies included targeted additional therapy in their trials) and two studies failed to present their results.

In the lower limb, five trials examined passive range of movement^{37,54-57}, seven measured global assessment of benefit^{37,54,55,57-60}, two looked at carer burden with the hygiene score^{54,61} and only one trial set goals³⁷. Treatment was either aimed at the hip adductors or the triceps surae. The patient groups were more varied in the lower limb; stroke, acquired brain injury, multiple sclerosis and cerebral palsy. Again, the methodology and presentation of the results varied widely across trials making comparison difficult.

Results were less notable for the treatment groups in the lower limb; two trials found improvements in passive range of movement but failed to reach significance, four trials found significant improvements in global assessment of benefit. The results for goal setting were not presented. Both trials that used the hygiene score found a

significant improvement for Botulinum Toxin groups but, unfortunately, their results were not suitable for Meta analysis.

Quantitative Analysis

A Meta analysis of the upper limb results for weeks four to twelve, demonstrated a significant effect for all outcomes in favour of Botulinum Toxin with moderate quality of evidence, with the exception of the risk ratio analysis of clinician rated scores; these were of a lower quality due to small event numbers^{31-33,35,38,41, 44-47,49}. (See Tables 1 and 2)

Overall pooled results gave a standardised mean difference of 0.80 (CI 0.55, 1.06), $Z=6.17$, $p<0.0001$. This equates to a large effect size using Cohen's descriptions.⁶² (See Figure 1)

The risk ratio was also calculated for the global assessment of benefit measures in the upper limb to calculate the number needed to treat (NNT). Patient rated = RR 2.21, (CI 1.67, 2.93), $P<0.0001$, NNT=5. Clinician rated = RR 2.51, (CI 1.21, 5.20), $P=0.01$, NNT=6.

A Meta analysis of upper limb outcomes for weeks 12 to 24, demonstrated a continued significant effect in favour of Botulinum Toxin for individual outcomes except the goal attainment scale where analysis was not possible. (See Table 2)

Pooled analysis of outcomes gave a continued significant effect but with a moderate/low effect size; SMD 0.48 (CI 0.34, 0.62), $Z= 6.62$, $P<0.0001$.

Meta analysis of the lower limb studies was only possible for the global assessment of benefit and only in dichotomous form^{54,55,58}. Both the patient and clinician rated scores failed to demonstrate a significant effect and were rated as low and very low

quality evidence respectively. (See Figures 2a and 2b) (Full details of grading can be found in Appendices II and III)

DISCUSSION

In this review significant results in favour of Botulinum Toxin A were found for the upper limb for all outcomes. The lower limb is less investigated in the literature and the results were not significant or were not suitable for a Meta analysis.

The quality of evidence was moderate overall for the upper limb. Most studies were graded down for inconsistencies or lack of clarity on randomisation, blinding, methodology and presentation of results.

Previous Meta analyses^{8,9} on global assessment of benefit concur with the findings of this review and although the secondary analysis of dichotomous outcomes was a lower quality of evidence due to low event numbers, the numbers needed to treat are promising.

Passive range of movement has not been found to be significantly altered in other reviews^{13,14}, with the exception of shoulder external rotation at four to six weeks.¹⁴ This review included a greater number of studies in the analysis, didn't use median values and chose change from baseline scores rather than final values; which may explain the different result here. Although passive range of movement does not measure ease of care directly, the carers ability to passively open fingers or move a shoulder or hip does impact on the effort required for washing and dressing and as an outcome measure is used frequently in the clinical setting. For these reasons it has been included in this review.

Foley et al¹⁵ also found a significant effect using the Disability Assessment Scale but with only a moderate effect size of 0.53. However their study included both the Barthel and Action Research Arm test as outcome measures, which look at global daily functions and active upper limb movements respectively. The greater effect size here may be attributed to using outcomes more specific to ease of care.

The Disability Assessment Scale is a valuable outcome to use clinically and for future trials; it has been validated in stroke, covers areas of self care that are important to the patient and includes the patient choosing a goal for their target area. It was also developed primarily to detect a change from a focal treatment such as Botulinum Toxin⁶³.

The Goal Attainment Scale needs further investigation as it also showed a significant result but with small sample sizes; it remains a useful clinical tool, as it is an objective and measurable way of setting goals with patient involvement. The setting of a patient tailored goal is extremely important and ensures that the treatment is focussed on the individual and their priorities⁶⁴. Ascertaining the nature of the goals set was difficult as few studies provided details. The majority of goals in the two studies reviewed here were around self care and clinically goals are often set around hygiene and self care; therefore the results have been included in this review. However an argument could easily be made for inclusion in a review of active outcomes in some cases.

A more consistent approach to the use of carer burden scales would allow this important outcome measure to be used in future Meta analyses.

Any official guideline recommendation for the use of Botulinum Toxin A to reduce problems with ease of care will have to balance the positive outcomes against costs and any possible harm from its use. An interesting outcome in this review was the

potential longevity of the toxin's effects. A significant effect was demonstrated for up to six months for most outcomes. Although further trials are needed to confirm the length of efficacy as the small sample sizes reduce the quality of evidence.

While these results do suggest that Botulinum Toxin A could be a potentially cost effective option for improving ease of care in the upper limb; this may be further enhanced by concurrent targeted therapy¹². In this review, only two trials used targeted therapy in conjunction with Botulinum Toxin; both found significant results. It is likely that the addition of targeted physiotherapy and occupational therapy with the injection would produce enhanced results, as it allows for stretching and splinting of the relaxed soft tissue structures, while electrical stimulation has been postulated to increase the uptake of the toxin⁶. A review of observational studies may also support this, as rehabilitative therapies are more often used in conjunction with Botulinum Toxin A outside the artificial constraints of a controlled trial.

The lower limb evidence suffers from fewer trials to review. Where studies have been carried out the methodology is often lacking. It is worth noting that in one study, which was highly scored for quality, patient rated scales showed significant improvement for Botulinum Toxin A⁵⁸. Further studies are needed with clear parameters for measuring care outcomes. The absence of goal setting methods in the lower limb was particularly evident. Carer burden scales would also provide useful information. The studies that did use a hygiene scale found positive results but the results were not suitable for Meta analysis. As the overall quality of evidence is very low, no conclusions can be drawn for the use of Botulinum Toxin A on improving ease of care in the lower limb. Clinically, treatment should be in conjunction with targeted goal setting and the use of carer burden scales to ensure the reduction of spasticity is meaningful for the patient¹⁷.

A drawback of many studies was that ease of care limitations were not a primary focus of the trial or the inclusion criteria. Often studies had equivocal results, which could have been attributed to the selection of higher functioning patients with minimal care restrictions. Most of the studies focused on measuring spasticity as their primary measure; this affected their inclusion criteria and subject population, and explains why often results were not reported fully. It can be noted that in studies where the focus was on care measures such as the disability assessment scale then the inclusion criteria was more appropriate and better results were gained.

A major limitation recognised by this review is the use of only one author to carry out the search and critique the articles. PRISMA recommends that the use of at least two investigators reduces the risk of rejecting relevant reports⁶⁵. Every attempt was made to minimise bias by the use of pre-drawn up forms, criteria and by double checking results. A second author was used for statistical analysis and robustness. It is a possibility that relevant studies have been missed in the search or that unwitting bias has been allowed into some decisions. Other criticisms may be directed at some of the subjective decisions on grading the quality of evidence. The decisions are based on uniform criteria and any diversions are fully explained. As documented by GRADE; as long as decisions are justified with clear reasoning then different recommendations can be valid and must be judged by the clinician on this basis¹. The decision to group dose ranging studies may also be an area for debate. A Meta analysis in this area is often problematic. Trials look at differing indications, injection techniques, doses and muscles as well as a wide variety of outcome measurement; this is often alongside the use of ordinal measures with parametric statistical tests.⁶⁶ Therefore any results and conclusions must take into account these limitations.

Randomised controlled trials need to carry out studies with carefully selected outcomes, inclusion criteria and methodology to enable a comprehensive Meta analysis with high quality evidence in the future. Lower limb studies are especially needed to provide clear data for analysis. Focusing on carer burden scales and goal setting would provide more meaningful results for patients and clinicians.

Observational studies need to be reviewed and may provide additional information especially on the effect of additional therapies.

This review supports the use of Botulinum Toxin A for improving ease of care in the upper limb; demonstrated by the global assessment of benefit, disability assessment scale, goal attainment scale and passive range of movement outcome measures. The benefit may last for up to six months in some cases. No decision can be made for the lower limb.

Clinical Messages

- Botulinum Toxin A can be recommended for improving ease of care in the upper limb; effects are evident for up to six months.
- No conclusions can be drawn for the lower limb.

Acknowledgements: None

Author Contributions: JB carried out literature search, grading of evidence and writing of main paper. GP advised and contributed to the statistical analysis and presentation of results.

Competing interests: None declared.

Funding support: Funding was provided by the Australian National Health and Medical Research Council (NHMRC, <http://www.nhmrc.gov.au>) NHMRC Early career fellowship (1052236 for GP)

References.

1. GRADE working group. Grading quality of evidence and strength of recommendations. *BMJ* 2004; **328**: 1490-1497
2. Baker J, Pereira G. The efficacy of Botulinum toxin A for spasticity and pain: a systematic review and meta analysis using the GRADE approach. *Clin Rehabil* ??
3. Sheean G.L. Botulinum treatment of spasticity: why is it so difficult to show a functional benefit? *Curr Opinion Neurol* 2001; **14**: 771-776
4. Elovic E.P., Brashear A, Kaelin D et al. Repeated treatments with botulinum toxin type A produce sustained decreases in the limitations associated with focal upper limb poststroke spasticity for caregivers and patients. *Arch Phys Med Rehabil* 2008; **89**: 799-806
5. Barnes M Botulinum toxin- mechanisms of action and clinical use in spasticity *J Rehabil Med* 2003; **41**: 56-59
6. Gracies J-M. Physiological effects of botulinum toxin in spasticity. *Mov Disord* 2004; **19** (S8): S120-S128
7. Francis HP, Wade DT, Turner-Stokes, Kingswell RS et al. Does reducing spasticity translate into functional benefit? An exploratory meta-analysis. *J Neurol Neurosurg Psychiatry* 2004; **75**: 1547-1551
8. Cardoso E, Rodrigues B, Lucena R et al. Botulinum toxin type A for the treatment of the upper limb spasticity after stroke- a meta analysis. *Arq Neuropsiquiatr* 2005; **63** (1): 30-33
9. Rosales RL and Chua-Yap AS. Evidence based systematic review on the efficacy and safety of botulinum toxin A therapy in post stroke spasticity. *J Neural Transm* 2008; **115**: 617-623
10. Simpson DM, Gracies J-M, Graham HK et al. Assessment: Botulinum neurotoxin for the treatment of spasticity (an evidence based review): Report of the therapeutics and technology assessment subcommittee of the American Academy of Neurology. *Neurology* 2008; **70**: 1691-1698
11. Wissel J, Ward A, Erztgaard P et al European consensus table on the use of botulinum toxin type A in adult spasticity *J Rehabil Med* 2009; **41**: 13-25
12. Sheean G, Lannin NA, Turner-Stokes L, Rawicki B, Snow BJ. Botulinum toxin assessment, intervention and after-care for upper limb hypertonicity in adults: international consensus statement. *Eur J Neurol* 2010; **17** (S2) : 74-93
13. Garces K, McCormick A, McGahan L, Skidmore B. *Botulinum toxin A upper and lower limb spasticity: a systematic review* [Technology report no 51]. Ottawa: Canadian Coordinating Office for Health Technology Assessment; 2005.
14. Singh JA and Fitzgerald PM. Botulinum toxin for shoulder pain (review). *The Cochrane Collaboration*. 2011: Issue 1: 1-51
15. Foley N, Pereira S, Salter K et al. Treatment with Botulinum Toxin improves Upper-Extremity function post stroke: A systematic review and meta-analysis. *Arch Phys Med Rehabil* 2013; **94** : 977-989
16. Guyatt GH, Oxman AD, Vist GE et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008; **336**: 924-926
17. Royal College of Physicians, British Society of Rehabilitation Medicine, Chartered Society of Physiotherapy, Association of Chartered Physiotherapists Interested in Neurology. *Spasticity in adults: management using botulinum toxin. National guidelines*. London: RCP, 2009.
18. Harbour R and Miller J. A new system for grading recommendations in evidence based guidelines. *BMJ* 2001; **323**: 334-336
19. Guyatt GH, Oxman AD, Kunz R et al. GRADE guidelines:2. Framing the question and deciding on important outcomes. *J Clin Epidemiol* 2011; **64**: 395-400
20. Guyatt GH, Oxman AD, Akl EA et al. GRADE guidelines:1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011; **64**: 383-394

21. Balshem H, Helfand M, Schunemann HJ et al. GRADE guidelines:3. Rating the quality of evidence. *J Clin Epidemiol* 2011; **64**: 401-406.
22. Guyatt GH, Oxman AD, Vist G et al. GRADE guidelines:4. Rating the quality of evidence-study limitations (risk of bias). *J Clin Epidemiol* 2011; **64**: 407-415
23. Guyatt GH, Oxman AD, Montori V et al. GRADE guidelines: 5. Rating the quality of evidence- publication bias. *J Clin Epidemiol* 2011; **64**: 1277-1282
24. Guyatt GH, Oxman AD, Kunz R et al. GRADE guidelines: 6. Rating the quality of evidence- imprecision. *J Clin Epidemiol* 2011; **64**: 1283-1293
25. Guyatt GH, Oxman AD, Kunz R et al. GRADE guideline: 7. Rating the quality of evidence- inconsistency. *J Clin Epidemiol* 2011; **64**: 1294-1302
26. Guyatt GH, Oxman AD, Kunz R et al. GRADE guidelines: 8. Rating the quality of evidence-indirectness. *J Clin Epidemiol* 2011; **64**: 1303-1310
27. Guyatt GH, Oxman AD, Sultan S et al. GRADE guidelines: 9. Rating up the quality of evidence. *J Clin Epidemiol* 2011; **64**: 1311-1316
28. GRADE working group. <http://www.gradeworkinggroup.org/index.htm> (last accessed May 2012)
29. Higgins JPT, Green S (editors). *Cochrane Handbook for systematic reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. www.cochrane-handbook.org
30. Bhakta BB, Cozens JA, Chamberlain MA et al. Impact of botulinum toxin Type A on disability and carer burden due to arm spasticity after stroke: a randomised double blind placebo controlled trial. *J Neurol Neurosurg Psychiatry* 2000; **69** : 217-221
31. Bakheit AMO, Thilman AF, Ward AB et al. A randomised, double-blind, placebo-controlled, dose-ranging study to compare the efficacy and safety of three doses of botulinum toxin Type A (Dysport) with placebo in upper limb spasticity after stroke. *Stroke* 2000; **31** : 2402-2406
32. Bakheit AMO, Pittock S, Moore AP et al. A randomized, double blind, placebo controlled study of the efficacy and safety of botulinum toxin type A in upper limb spasticity in patients with stroke. *J Neurol* 2001; **8**: 559-565
33. De Boer KS, Arwert HJ, De Groot JH et al. Shoulder pain and external rotation in spastic hemiplegia do not improve by injection of botulinum toxin A into the subscapular muscle. *J Neurol Neurosurg Psychiatry* 2008; **79**: 581-583
34. Yelnik AP, Colle FM, Bonan IV, Vicout E. Treatment of shoulder pain in spastic hemiplegia by reducing spasticity of the subscapular muscle: a randomised, double blind, placebo controlled study of botulinum toxin A. *J Neurol Neurosurg Psychiatry* 2007; **78**: 845-848
35. Marco E, Duarte E, Vila J et al. Is botulinum toxin type A effective in the treatment of spastic shoulder pain in patients after stroke? A double blind randomized clinical trial. *J Rehabil Med* 2007; **39**: 440-447
36. Kong K-H, Neo J-J, Chua SG A randomized controlled study of botulinum toxin A in the treatment of hemiplegic shoulder pain associated with spasticity. *Clin Rehabil* 2007; **21**: 28- 35
37. Richardson D, Sheean G, Werring D et al Evaluating the role of botulinum toxin in the management of focal hypertonia in adults. *J Neurol Neurosurg Psychiatry* 2000; **69**: 499-506.
38. Smith SJ, Ellis E, White S et al. A double-blind, placebo-controlled study of botulinum toxin in upper limb spasticity after stroke or head injury. *Clin Rehabil* 2000; **14** : 5-13
39. Meythaler JM, Vogtle L, Brunner RC. A preliminary assessment of the benefits of the addition of botulinum toxin A to a conventional therapy program on the function of people with longstanding stroke. *Arch Phys Med Rehabil* 2009; **90**: 1453-1461
40. Cousins E, Ward A, Roffe C, Rimington L et al. Does low dose botulinum toxin help the recovery of arm function when given early after stroke? A phase II randomised controlled pilot study to estimate effect size. *Clin Rehabil* 2010; **24**: 501-513

41. Lam K, Lau KK, So KK et al. Can botulinum toxin decrease carer burden in long term care residents with upper limb spasticity? A randomised controlled study. *J Am Med Dir Assoc* 2012; **13**: 477-484
42. Rosales RL, Kong KH, Goh KJ et al. Botulinum Toxin Injection for Hypertonicity of the upper extremity within 12 weeks after stroke: A randomised controlled trial. *J Neurorehabil Neural Repair* 2012; **26** (7): 812-821
43. Marciniak CM, Harvey RL, Gagnon CM et al. Does Botulinum Toxin Type A decrease pain and lessen disability in hemiplegic survivors of stroke with shoulder pain and spasticity? *Am J Phys Med Rehabil* 2012; **91**: 1007-1019
44. Brashear A, Gordon MF, Elovic E et al. Intramuscular injection of botulinum toxin for the treatment of wrist and finger spasticity after a stroke. *N Engl J Med* 2002; **347** (6) : 395-400
45. Simpson DM, Gracies JM, Yablon SA, Barbano R, Brashear A. Botulinum neurotoxin versus tizanidine in upper limb spasticity: a placebo controlled study. *J Neurol Neurosurg Psychiatry* 2009; **80**: 380-385
46. Kaji R, Osako Y, Suyama K et al. Botulinum toxin type A in post-stroke upper limb spasticity. *Curr Med Res Opinion* 2010; **26** (8) : 1983-1992
47. McCrory P, Turner-Stokes L, Baguley IJ et al. Botulinum toxin A for treatment of upper limb spasticity following stroke: a multi centre randomised placebo controlled study of the effects on quality of life and other person-centred outcomes. *J Rehabil Med* 2009; **41**: 536-544
48. Childers MK, Brashear A, Jozefczyk P et al. Dose-dependent response to intramuscular botulinum toxin type A for upper limb spasticity in patients after stroke. *Arch Phys Med Rehabil* 2004; **85**: 1063-1069
49. Simpson DM, Alexander DN, O'Brien CF et al. Botulinum toxin type A in the treatment of upper extremity spasticity: a randomised, double blind, placebo controlled trial. *Neurology* 1996; **46**: 1306-1310
50. Bhakta BB, O'Connor RJ, Cozens JA. Associated Reactions after stroke: a randomised controlled trial of the effect of botulinum toxin type A. *J Rehabil Med* 2008; **40** : 36-41
51. Hesse S, Mach H, Behrend S, Werner C, Melzer I. An early botulinum toxin A treatment in subacute stroke patients may prevent a disabling finger flexor stiffness six months later: a randomised controlled trial. *Clin Rehabil* 2011; **26** (3): 239-245
52. Rodgers H, Shaw L, Price C et al. BoTULS trial: a randomised controlled trial to evaluate the clinical effect and cost effectiveness of treating upper limb spasticity due to stroke with botulinum toxin type A. *Health Technol Assess* 2010; **14**(26): 1-141
53. Hesse S, Reiter F, Konrad M, Jahnke MT. Botulinum toxin type A and short term electrical stimulation in the treatment of upper limb flexor spasticity after stroke: a randomised, double blind, placebo controlled trial. *Clin Rehabil* 1998; **12**: 381-388
54. Hyman N, Barnes M, Bhakta B et al. Botulinum toxin (Dysport) treatment of hip adductor spasticity in multiple sclerosis: a prospective, randomized, double blind, placebo controlled, dose ranging study. *J Neurol Neurosurg Psychiatry* 2000; **68**: 707-712
55. Pittock SJ, Moore AP, Hardiman O et al. A double blind randomized placebo controlled evaluation of three doses of botulinum toxin type A (Dysport) in the treatment of spastic equinovarus deformity after stroke. *Cerebrovasc Dis* 2003; **15**: 289-300
56. Verplancke D, Snape S, Salisbury CF et al. A randomized controlled trial of botulinum toxin on lower limb spasticity following acute acquired severe brain injury. *Clin Rehabil* 2005; **19**: 117-125
57. Gusev YI, Banach M, Simonow A et al. Efficacy and safety of Botulinum toxin Type A Toxin in Adductor spasticity due to Multiple Sclerosis. *J Musculoskelet Pain* 2008; **16** (3): 175-188

58. Maanum G, Jahnsen R, Stanghelle JK et al. Effects of botulinum toxin A in ambulant adults with spastic cerebral palsy: a randomised double blind placebo controlled trial. *J Rehabil Med* 2011; **43**: 338-347
59. Kaji R, Osako Y, Suyama K et al. Botulinum toxin type A in post stroke lower limb spasticity: a multicenter, double blind, placebo controlled trial. *J Neurol* 2010; **257**: 1330-1337
60. Burbaud P, Wiart L, Dubos JL et al. A randomised, double blind, placebo controlled trial of botulinum toxin in the treatment of spastic foot in hemiparetic patients. *J Neurol Neurosurg Psychiatry* 1996; **61 (3)**: 265-269
61. Snow BJ, Tsui JKC, Bhatt MH, Varelans M, Hashimoto SA, Calne DB. Treatment of spasticity with botulinum toxin: a double blind study. *Annals Neurol* 1990; **28 (4)**: 512-515
62. Cohen J. *Statistical power analysis for the behavioural sciences*. 2nd ed New York: Academic Press 1988
63. Brashear A and Lambeth K. Spasticity *Curr Treat Options Neurol* 2009; **11**: 153-161
64. Stevenson VL. Spasticity Management. *Clin Rehabil* 2010; **24**: 293-304
65. Liberati A, Altman D, Tetzlaff J et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med* 2009; **151 (4)**: W65-W94
66. Moore AP. Botulinum toxin A (BoNT-A) for spasticity in adults. What is the evidence? *Eur J Neurol* 2002; **9 (S1)**: 42-47

Table 1. Grades of evidence for the efficacy of Botulinum Toxin A on improving ease of self care in the upper and lower limb.

	PROM UL	DAS UL	GAS UL	GAB Pt UL (SMD)	GAB Cl UL (SMD)	GAB Pt UL (RR)	GAB Cl UL (RR)	Carer burden	PROM LL	Goals LL	GAB Pt LL	GAB Cl LL	Hygiene scale
Type of study	RCT (n=6)	RCT (n=3)	RCT (n=2)	RCT (n=4)	RCT (n=3)	RCT (n=4)	RCT (n=3)	RCT (n=5)	RCT (n=3)	RCT (n=1)	RCT (n=3)	RCT (n=2)	RCT (n=2)
Study Quality	-1	-1	0	-1	-1	-1	-1	-1	-1	N/A	-1	-2	-1
Study Consistency	0	0	0	0	0	0	0	0	0	N/A	0	0	0
Study Directness	0	0	0	0	0	0	0	0	0	N/A	0	0	0
Study precision	0	0	-1	0	0	0	-1	-1	-2	N/A	-1	-2	-2
Upgrades	None	None	None	None	None	None	None	None	None		None	None	None
Overall Grade	Mod	Mod	Mod	Mod	Mod	Mod	Low	Low	V Low		Low Low	V	V Low

PROM, passive range of movement; UL, upper limb; DAS, disability assessment scale; GAS, goal attainment scale; GAB, global assessment of benefit; Pt, patient rated; Cl, clinician rated; SMD, standardised mean difference; RR, risk ratio; LL, lower limb; Mod, moderate; V Low, very low; N/A, not possible to grade due to lack of evidence.

